



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:45 AM UTC

PDB ID : 5EEL / pdb_00005eel
Title : Grb7 SH2 with bicyclic peptide inhibitor
Authors : Watson, G.M.; Gunzburg, M.J.; Wilce, M.C.J.; Wilce, J.A.
Deposited on : 2015-10-23
Resolution : 2.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	120	7% 80% 7% 13%
1	B	120	2% 85% 6% 9%
1	C	120	8% 78% 10% 12%
1	D	120	4% 79% 6% 15%
1	E	120	4% 81% • 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	120	
2	L	9	
2	M	9	
2	N	9	
2	P	9	
2	Q	9	
2	R	9	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FMT	A	602	-	-	X	-
5	NA	C	603	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Growth factor receptor-bound protein 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	104	Total	C	N	O	S	0	0	0
			816	519	148	144	5			
1	B	109	Total	C	N	O	S	0	0	0
			866	550	162	149	5			
1	C	106	Total	C	N	O	S	0	0	0
			836	531	156	144	5			
1	D	102	Total	C	N	O	S	0	0	0
			817	520	150	142	5			
1	E	102	Total	C	N	O	S	0	0	0
			808	514	146	143	5			
1	F	104	Total	C	N	O	S	0	0	0
			837	531	156	145	5			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	413	GLY	-	expression tag	UNP Q14451
A	414	SER	-	expression tag	UNP Q14451
B	413	GLY	-	expression tag	UNP Q14451
B	414	SER	-	expression tag	UNP Q14451
C	413	GLY	-	expression tag	UNP Q14451
C	414	SER	-	expression tag	UNP Q14451
D	413	GLY	-	expression tag	UNP Q14451
D	414	SER	-	expression tag	UNP Q14451
E	413	GLY	-	expression tag	UNP Q14451
E	414	SER	-	expression tag	UNP Q14451
F	413	GLY	-	expression tag	UNP Q14451
F	414	SER	-	expression tag	UNP Q14451

- Molecule 2 is a protein called Bicyclic Peptide Inhibitor.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	9	Total	C	N	O	S	0	0	0
			78	48	11	18	1			
2	M	9	Total	C	N	O	S	0	0	0
			78	48	11	18	1			
2	N	9	Total	C	N	O	S	0	0	0
			78	48	11	18	1			
2	P	8	Total	C	N	O		0	0	0
			68	43	9	16				
2	Q	9	Total	C	N	O	S	0	0	0
			78	48	11	18	1			
2	R	9	Total	C	N	O	S	0	0	0
			78	48	11	18	1			

- Molecule 3 is MALONIC ACID (CCD ID: MLA) (formula: $C_3H_4O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	3	4		
3	C	1	Total	C	O	0	0
			7	3	4		
3	D	1	Total	C	O	0	0
			7	3	4		
3	E	1	Total	C	O	0	0
			7	3	4		
3	F	1	Total	C	O	0	0
			7	3	4		

- Molecule 4 is FORMIC ACID (CCD ID: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	C	1	Total	C	O	0	0
			3	1	2		
4	D	1	Total	C	O	0	0
			3	1	2		
4	E	1	Total	C	O	0	0
			3	1	2		
4	F	1	Total	C	O	0	0
			3	1	2		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		
5	B	1	Total	Na	0	0
			1	1		
5	C	1	Total	Na	0	0
			1	1		
5	D	1	Total	Na	0	0
			1	1		
5	E	1	Total	Na	0	0
			1	1		
5	F	1	Total	Na	0	0
			1	1		



- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	6	Total 6	O 6	0	0
6	B	22	Total 22	O 22	0	0
6	C	1	Total 1	O 1	0	0
6	D	6	Total 6	O 6	0	0
6	E	10	Total 10	O 10	0	0
6	F	22	Total 22	O 22	0	0
6	L	4	Total 4	O 4	0	0
6	N	3	Total 3	O 3	0	0
6	R	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Growth factor receptor-bound protein 7




- Molecule 1: Growth factor receptor-bound protein 7





- Molecule 1: Growth factor receptor-bound protein 7





- Molecule 1: Growth factor receptor-bound protein 7




- Molecule 1: Growth factor receptor-bound protein 7



- Molecule 1: Growth factor receptor-bound protein 7




- Molecule 2: Bicyclic Peptide Inhibitor




- Molecule 2: Bicyclic Peptide Inhibitor



- Molecule 2: Bicyclic Peptide Inhibitor



- Molecule 2: Bicyclic Peptide Inhibitor



- Molecule 2: Bicyclic Peptide Inhibitor



- Molecule 2: Bicyclic Peptide Inhibitor



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	95.23Å 95.23Å 241.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.72 – 2.47 46.72 – 2.47	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.72-2.47) 99.9 (46.72-2.47)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.48 (at 2.48Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.185 , 0.232 0.198 , 0.240	Depositor DCC
R_{free} test set	2050 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	37.9	Xtriage
Anisotropy	0.134	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5572	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, FMT, 73C, MLA, AEA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/833	0.84	3/1125 (0.3%)
1	B	0.45	0/884	0.78	2/1193 (0.2%)
1	C	0.41	0/853	0.77	0/1152
1	D	0.39	0/834	0.80	2/1125 (0.2%)
1	E	0.48	0/825	0.80	2/1114 (0.2%)
1	F	0.60	0/854	0.91	6/1151 (0.5%)
2	L	0.58	0/59	0.72	0/78
2	M	0.40	0/59	0.57	0/78
2	N	0.39	0/59	0.56	0/78
2	P	0.22	0/59	0.44	0/78
2	Q	0.45	0/59	0.71	0/78
2	R	0.47	0/59	0.59	0/78
All	All	0.47	0/5437	0.81	15/7328 (0.2%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	487	GLU	N-CA-C	-7.79	97.94	110.32
1	A	427	ARG	N-CA-C	6.99	118.90	111.28
1	A	463	ASN	CA-C-N	6.72	128.24	119.84
1	A	463	ASN	C-N-CA	6.72	128.24	119.84
1	E	463	ASN	CA-C-N	6.31	125.80	119.24
1	E	463	ASN	C-N-CA	6.31	125.80	119.24
1	F	501	ARG	N-CA-C	6.06	119.35	109.59
1	F	483	LEU	CA-C-N	-5.98	114.45	120.31
1	F	483	LEU	C-N-CA	-5.98	114.45	120.31
1	D	463	ASN	CA-C-N	5.83	127.13	119.84
1	D	463	ASN	C-N-CA	5.83	127.13	119.84
1	B	463	ASN	CA-C-N	5.77	127.05	119.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	463	ASN	C-N-CA	5.77	127.05	119.84
1	F	463	ASN	CA-C-N	5.69	126.07	119.47
1	F	463	ASN	C-N-CA	5.69	126.07	119.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	816	0	777	4	0
1	B	866	0	845	5	0
1	C	836	0	815	7	0
1	D	817	0	800	3	0
1	E	808	0	778	2	0
1	F	837	0	821	3	0
2	L	78	0	48	1	0
2	M	78	0	48	0	0
2	N	78	0	48	1	0
2	P	68	0	43	0	0
2	Q	78	0	48	1	0
2	R	78	0	49	1	0
3	A	7	0	2	0	0
3	C	7	0	2	1	0
3	D	7	0	2	0	0
3	E	7	0	2	0	0
3	F	7	0	2	0	0
4	A	3	0	1	2	0
4	B	3	0	1	0	0
4	C	3	0	1	0	0
4	D	3	0	1	0	0
4	E	3	0	1	0	0
4	F	3	0	1	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
6	A	6	0	0	0	0
6	B	22	0	0	0	0
6	C	1	0	0	0	0
6	D	6	0	0	0	0
6	E	10	0	0	0	0
6	F	22	0	0	0	0
6	L	4	0	0	1	0
6	N	3	0	0	1	0
6	R	1	0	0	0	0
All	All	5572	0	5136	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:448:GLN:HB2	1:D:454:LEU:HD21	1.72	0.71
1:B:435:ARG:NH1	1:B:461:GLN:OE1	2.28	0.67
1:A:512:HIS:NE2	4:A:602:FMT:H	2.10	0.67
2:Q:7:ASN:N	2:Q:7:ASN:OD1	2.31	0.62
1:A:458:ARG:NH1	1:A:470:SER:OG	2.33	0.61
1:B:445:ILE:HG23	1:B:454:LEU:HD23	1.84	0.58
1:E:485:SER:HB2	1:E:492:TYR:CZ	2.42	0.55
1:B:435:ARG:HH12	1:B:461:GLN:CD	2.17	0.53
1:B:482:ILE:HG12	1:B:495:MET:HG2	1.91	0.52
1:C:432:PHE:O	1:C:529:ARG:HD3	2.10	0.52
1:C:481:LEU:HD21	1:C:483:LEU:HD21	1.94	0.50
2:N:9:AEA:N2	6:N:101:HOH:O	2.33	0.49
1:C:482:ILE:HG12	1:C:495:MET:HG2	1.95	0.48
1:C:460:SER:OG	3:C:601:MLA:O1A	2.31	0.47
2:R:1:SER:HB3	2:R:7:ASN:HB2	1.97	0.46
1:A:502:PHE:HD1	1:A:507:GLN:HG2	1.80	0.46
1:C:452:ASP:OD1	1:C:473:HIS:ND1	2.51	0.43
1:D:482:ILE:HG12	1:D:495:MET:HG2	2.00	0.43
1:F:433:HIS:NE2	1:F:527:CYS:O	2.38	0.42
1:C:427:ARG:HE	1:C:505:LEU:HD12	1.84	0.42
1:A:512:HIS:NE2	4:A:602:FMT:C	2.81	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:435:ARG:HH12	1:B:461:GLN:NE2	2.17	0.41
1:C:427:ARG:HG3	1:C:432:PHE:CD2	2.55	0.41
1:F:490:ARG:H	1:F:490:ARG:HG2	1.58	0.41
2:L:9:AEA:N2	6:L:101:HOH:O	2.37	0.41
1:F:512:HIS:HA	1:F:515:ASN:O	2.20	0.41
1:E:433:HIS:NE2	1:E:527:CYS:O	2.54	0.41
1:D:512:HIS:HA	1:D:515:ASN:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/120 (85%)	102 (100%)	0	0	100	100
1	B	107/120 (89%)	107 (100%)	0	0	100	100
1	C	104/120 (87%)	102 (98%)	2 (2%)	0	100	100
1	D	100/120 (83%)	99 (99%)	1 (1%)	0	100	100
1	E	100/120 (83%)	100 (100%)	0	0	100	100
1	F	102/120 (85%)	102 (100%)	0	0	100	100
2	L	6/9 (67%)	6 (100%)	0	0	100	100
2	M	6/9 (67%)	5 (83%)	1 (17%)	0	100	100
2	N	6/9 (67%)	6 (100%)	0	0	100	100
2	P	6/9 (67%)	6 (100%)	0	0	100	100
2	Q	6/9 (67%)	6 (100%)	0	0	100	100
2	R	6/9 (67%)	5 (83%)	1 (17%)	0	100	100
All	All	651/774 (84%)	646 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/106 (80%)	84 (99%)	1 (1%)	63	81
1	B	91/106 (86%)	91 (100%)	0	100	100
1	C	88/106 (83%)	85 (97%)	3 (3%)	32	57
1	D	88/106 (83%)	88 (100%)	0	100	100
1	E	86/106 (81%)	86 (100%)	0	100	100
1	F	90/106 (85%)	87 (97%)	3 (3%)	33	58
2	L	6/6 (100%)	6 (100%)	0	100	100
2	M	6/6 (100%)	6 (100%)	0	100	100
2	N	6/6 (100%)	6 (100%)	0	100	100
2	P	6/6 (100%)	6 (100%)	0	100	100
2	Q	6/6 (100%)	4 (67%)	2 (33%)	0	0
2	R	6/6 (100%)	6 (100%)	0	100	100
All	All	564/672 (84%)	555 (98%)	9 (2%)	55	77

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	497	ASP
1	C	427	ARG
1	C	439	GLU
1	C	500	THR
1	F	487	GLU
1	F	491	LEU
1	F	507	GLN
2	Q	3	GLU
2	Q	7	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	448	GLN
1	A	473	HIS
1	B	442	GLN
1	B	448	GLN
1	C	525	HIS
1	D	442	GLN
1	D	515	ASN
1	E	507	GLN
1	F	442	GLN
1	F	447	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

11 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	AEA	Q	9	2	9,9,10	0.61	0	8,10,12	0.48	0
2	AEA	M	9	2	9,9,10	0.69	0	8,10,12	0.68	0
2	AEA	R	9	2	9,9,10	0.78	0	8,10,12	0.62	0
2	73C	P	8	2	8,9,10	1.22	1 (12%)	5,9,11	1.98	1 (20%)
2	73C	M	8	2	8,9,10	1.29	1 (12%)	5,9,11	1.10	0
2	73C	N	8	2	8,9,10	1.26	1 (12%)	5,9,11	1.37	1 (20%)
2	AEA	N	9	2	9,9,10	0.62	0	8,10,12	0.92	1 (12%)
2	73C	R	8	2	8,9,10	1.26	1 (12%)	5,9,11	1.60	1 (20%)
2	AEA	L	9	2	9,9,10	0.65	0	8,10,12	0.82	0
2	73C	L	8	2	8,9,10	1.31	1 (12%)	5,9,11	1.14	0
2	73C	Q	8	2	8,9,10	1.29	1 (12%)	5,9,11	1.79	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AEA	Q	9	2	-	4/8/9/10	-
2	AEA	M	9	2	-	4/8/9/10	-
2	AEA	R	9	2	-	7/8/9/10	-
2	73C	P	8	2	-	3/6/8/10	-
2	73C	M	8	2	-	1/6/8/10	-
2	73C	N	8	2	-	1/6/8/10	-
2	AEA	N	9	2	-	3/8/9/10	-
2	73C	R	8	2	-	1/6/8/10	-
2	AEA	L	9	2	-	3/8/9/10	-
2	73C	L	8	2	-	2/6/8/10	-
2	73C	Q	8	2	-	2/6/8/10	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	R	8	73C	C3-C2	-3.25	1.31	1.51
2	L	8	73C	C3-C2	-3.21	1.31	1.51
2	Q	8	73C	C3-C2	-3.20	1.31	1.51
2	M	8	73C	C3-C2	-3.20	1.31	1.51
2	P	8	73C	C3-C2	-3.17	1.32	1.51
2	N	8	73C	C3-C2	-3.14	1.32	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	8	73C	OG-C1-C2	3.91	130.87	110.32
2	Q	8	73C	OG-C1-C2	-3.26	93.19	110.32
2	R	8	73C	OG-C1-C2	-2.48	97.28	110.32
2	N	8	73C	OG-C1-C2	-2.13	99.11	110.32
2	N	9	AEA	O1-C5-C4	-2.04	116.03	125.44

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	L	9	AEA	N1-C1-C3-S1
2	M	9	AEA	N1-C1-C3-S1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	M	9	AEA	N1-C1-C2-N2
2	M	9	AEA	N1-C1-C2-O3
2	N	9	AEA	N1-C1-C3-S1
2	N	9	AEA	N1-C1-C2-O3
2	Q	9	AEA	N1-C1-C3-S1
2	Q	9	AEA	C2-C1-C3-S1
2	Q	9	AEA	N1-C1-C2-N2
2	Q	9	AEA	N1-C1-C2-O3
2	R	9	AEA	N1-C1-C3-S1
2	R	9	AEA	C2-C1-C3-S1
2	R	9	AEA	N1-C1-C2-N2
2	R	9	AEA	C3-C1-C2-N2
2	R	9	AEA	C3-C1-C2-O3
2	Q	8	73C	OG-C1-C2-C3
2	N	8	73C	OG-C1-C2-C3
2	P	8	73C	OG-C1-C2-C3
2	P	8	73C	C1-C2-C3-C4
2	R	8	73C	OG-C1-C2-C3
2	L	8	73C	OG-C1-C2-C3
2	M	8	73C	OG-C1-C2-C3
2	M	9	AEA	C2-C1-C3-S1
2	N	9	AEA	C2-C1-C3-S1
2	R	9	AEA	N1-C1-C2-O3
2	L	9	AEA	N1-C1-C2-O3
2	Q	8	73C	C2-C1-OG-CB
2	P	8	73C	C2-C1-OG-CB
2	R	9	AEA	C1-C3-S1-C4
2	L	9	AEA	C2-C1-C3-S1
2	L	8	73C	C2-C1-OG-CB

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	N	9	AEA	1	0
2	L	9	AEA	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 17 ligands modelled in this entry, 6 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MLA	F	601	-	6,6,6	1.20	0	7,7,7	1.30	0
4	FMT	C	602	5	2,2,2	0.06	0	1,1,1	0.46	0
4	FMT	D	602	5	2,2,2	0.19	0	1,1,1	0.42	0
4	FMT	F	602	5	2,2,2	0.07	0	1,1,1	0.35	0
3	MLA	E	601	-	6,6,6	1.10	0	7,7,7	1.30	0
3	MLA	C	601	-	6,6,6	1.11	0	7,7,7	1.27	0
4	FMT	B	601	5	2,2,2	0.19	0	1,1,1	0.57	0
3	MLA	A	601	-	6,6,6	1.11	0	7,7,7	1.30	0
4	FMT	A	602	5	2,2,2	0.10	0	1,1,1	0.32	0
4	FMT	E	602	5	2,2,2	0.20	0	1,1,1	0.47	0
3	MLA	D	601	-	6,6,6	1.15	0	7,7,7	1.31	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLA	F	601	-	-	2/4/4/4	-
3	MLA	E	601	-	-	2/4/4/4	-
3	MLA	C	601	-	-	0/4/4/4	-
3	MLA	A	601	-	-	4/4/4/4	-
3	MLA	D	601	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	601	MLA	O1A-C1-C2	2.03	120.80	114.51

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	601	MLA	O1A-C1-C2-C3
3	A	601	MLA	C1-C2-C3-O3B
3	E	601	MLA	C1-C2-C3-O3A
3	D	601	MLA	O1B-C1-C2-C3
3	E	601	MLA	C1-C2-C3-O3B
3	A	601	MLA	O1A-C1-C2-C3
3	A	601	MLA	O1B-C1-C2-C3
3	A	601	MLA	C1-C2-C3-O3A
3	F	601	MLA	C1-C2-C3-O3B
3	F	601	MLA	C1-C2-C3-O3A

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	601	MLA	1	0
4	A	602	FMT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	104/120 (86%)	0.24	8 (7%) 19 17	32, 50, 78, 97	0
1	B	109/120 (90%)	-0.26	3 (2%) 55 51	23, 36, 63, 79	0
1	C	106/120 (88%)	0.49	10 (9%) 14 12	38, 56, 88, 111	0
1	D	102/120 (85%)	0.01	5 (4%) 35 31	32, 43, 75, 89	0
1	E	102/120 (85%)	-0.12	5 (4%) 35 31	25, 40, 64, 108	0
1	F	104/120 (86%)	-0.20	6 (5%) 29 25	23, 32, 70, 91	0
2	L	7/9 (77%)	-0.54	0 100 100	30, 32, 39, 45	0
2	M	7/9 (77%)	-0.65	0 100 100	32, 34, 40, 43	0
2	N	7/9 (77%)	-0.91	0 100 100	26, 29, 34, 36	0
2	P	7/9 (77%)	1.58	1 (14%) 6 5	74, 77, 94, 96	0
2	Q	7/9 (77%)	0.09	0 100 100	40, 53, 73, 76	0
2	R	7/9 (77%)	-0.15	0 100 100	35, 41, 50, 67	0
All	All	669/774 (86%)	0.02	38 (5%) 29 26	23, 43, 79, 111	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	427	ARG	6.8
1	A	426	HIS	6.1
1	F	428	THR	6.0
1	A	428	THR	5.1
1	A	529	ARG	4.7
1	E	488	GLU	4.1
1	D	529	ARG	4.0
1	B	421	LEU	3.9
1	C	488	GLU	3.8
1	E	429	GLN	3.7
1	A	528	THR	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	532	LEU	3.4
1	F	530	VAL	3.3
1	C	489	GLY	3.2
1	C	427	ARG	3.2
1	C	487	GLU	3.2
2	P	1	SER	3.2
1	D	428	THR	2.9
1	C	486	GLU	2.8
1	E	489	GLY	2.8
1	A	427	ARG	2.8
1	B	529	ARG	2.8
1	A	465	GLN	2.7
1	D	488	GLU	2.7
1	F	488	GLU	2.7
1	E	486	GLU	2.7
1	F	486	GLU	2.6
1	B	528	THR	2.6
1	D	486	GLU	2.5
1	A	488	GLU	2.5
1	E	530	VAL	2.4
1	C	428	THR	2.3
1	F	491	LEU	2.3
1	C	464	PRO	2.3
1	A	489	GLY	2.2
1	D	489	GLY	2.2
1	C	499	GLN	2.0
1	C	490	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	AEA	Q	9	10/11	0.72	0.18	55,69,75,101	0
2	73C	P	8	10/11	0.73	0.21	39,93,104,113	0
2	AEA	M	9	10/11	0.91	0.10	34,46,55,56	0
2	73C	Q	8	10/11	0.91	0.12	35,58,71,75	0
2	AEA	L	9	10/11	0.92	0.11	31,45,48,51	0
2	73C	M	8	10/11	0.93	0.11	30,42,68,70	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	AEA	R	9	10/11	0.93	0.13	29,41,50,54	0
2	73C	N	8	10/11	0.94	0.07	20,34,41,44	0
2	AEA	N	9	10/11	0.96	0.08	28,40,45,47	0
2	73C	L	8	10/11	0.96	0.08	19,36,48,48	0
2	73C	R	8	10/11	0.96	0.09	16,33,56,57	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NA	C	603	1/1	0.72	0.43	83,83,83,83	0
5	NA	B	602	1/1	0.79	0.35	80,80,80,80	0
5	NA	D	603	1/1	0.83	0.25	59,59,59,59	0
5	NA	F	603	1/1	0.83	0.22	55,55,55,55	0
5	NA	E	603	1/1	0.85	0.21	55,55,55,55	0
5	NA	A	603	1/1	0.87	0.24	74,74,74,74	0
4	FMT	A	602	3/3	0.87	0.18	44,44,47,51	0
3	MLA	D	601	7/7	0.90	0.10	34,42,45,49	0
3	MLA	C	601	7/7	0.90	0.12	54,57,59,61	0
4	FMT	D	602	3/3	0.90	0.23	55,55,58,60	0
4	FMT	C	602	3/3	0.92	0.21	77,77,77,78	0
4	FMT	E	602	3/3	0.93	0.15	46,46,49,53	0
4	FMT	F	602	3/3	0.93	0.12	45,45,45,46	0
3	MLA	E	601	7/7	0.93	0.09	36,40,44,47	0
4	FMT	B	601	3/3	0.93	0.21	41,41,45,48	0
3	MLA	F	601	7/7	0.94	0.08	33,39,41,41	0
3	MLA	A	601	7/7	0.97	0.09	26,29,39,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.