



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:40 AM UTC

PDB ID : 6EEQ / pdb_00006eeq
Title : Crystal structure of Rhodiola rosea 4-hydroxyphenylacetaldehyde synthase
Authors : Torrens-Spence, M.P.; Chiang, Y.; Smith, T.; Vicent, M.A.; Wang, Y.; Weng, J.K.
Deposited on : 2018-08-15
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	490	6% 72% 23% 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-hydroxyphenylacetaldehyde synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	467	Total	C	N	O	P	S	0	3	0
			3694	2345	631	692	1	25			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	11	Total	O	0	0
			11	11		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-hydroxyphenylacetaldehyde synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	118.36Å 118.36Å 67.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.69 – 2.60 83.69 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (83.69-2.60) 97.5 (83.69-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.60 (at 2.62Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.211 , 0.269 0.220 , 0.275	Depositor DCC
R_{free} test set	1518 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å ²)	58.5	Xtriage
Anisotropy	0.407	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3705	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/3747	0.41	0/5076

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3694	0	3661	79	0
2	A	11	0	0	0	0
All	All	3705	0	3661	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:LYS:H	1:A:50:LYS:HD3	1.44	0.81
1:A:284:ALA:CB	1:A:314:LLP:H2'2	2.10	0.81

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:284:ALA:HB2	1:A:314:LLP:H2'2	1.67	0.74
1:A:28:LEU:HD13	1:A:70:GLU:HG2	1.73	0.71
1:A:192:TYR:HE1	1:A:247:PRO:HB3	1.58	0.69
1:A:476:ASN:OD1	1:A:479:ARG:NH2	2.27	0.67
1:A:438:MET:HE2	1:A:451:GLY:HA3	1.79	0.64
1:A:32:PHE:HE2	1:A:77:ASN:HB2	1.61	0.63
1:A:39:THR:HG21	1:A:43:ARG:HH21	1.63	0.63
1:A:161:ASN:OD1	1:A:164:GLU:N	2.24	0.61
1:A:403:ASP:N	1:A:403:ASP:OD1	2.35	0.60
1:A:49:VAL:HG11	1:A:53:PHE:CG	2.36	0.60
1:A:39:THR:HG21	1:A:43:ARG:NH2	2.16	0.59
1:A:50:LYS:H	1:A:50:LYS:CD	2.13	0.58
1:A:31:ASP:HA	1:A:34:THR:HG22	1.86	0.58
1:A:289:ALA:HA	1:A:385:ILE:HD13	1.86	0.57
1:A:76:VAL:HA	1:A:80:ILE:HD12	1.87	0.57
1:A:67:GLU:HG3	1:A:71:VAL:HG11	1.87	0.56
1:A:142:LEU:HB2	1:A:144:LEU:HD12	1.87	0.56
1:A:360:LEU:HD23	1:A:360:LEU:H	1.71	0.56
1:A:269:GLY:HA2	1:A:279:LEU:HD12	1.88	0.56
1:A:327:LEU:HD12	1:A:330:PRO:HA	1.89	0.55
1:A:404:ARG:NH1	1:A:486:GLN:HG2	2.22	0.54
1:A:101:SER:HB3	1:A:317:LEU:HD22	1.90	0.53
1:A:449:LEU:HD11	1:A:477:LEU:HD11	1.90	0.53
1:A:143:ASN:ND2	1:A:296:GLN:OE1	2.33	0.52
1:A:162:THR:H	1:A:314:LLP:P	2.33	0.52
1:A:198:HIS:HB3	1:A:200:THR:HG22	1.90	0.52
1:A:130:LEU:HA	1:A:133:VAL:HG12	1.91	0.52
1:A:142:LEU:HB2	1:A:144:LEU:CD1	2.42	0.50
1:A:265:ILE:H	1:A:265:ILE:HD12	1.77	0.49
1:A:230:PRO:HG2	1:A:267:GLU:HB3	1.95	0.49
1:A:168:CYS:HB3	1:A:333:LEU:HD11	1.95	0.48
1:A:299:LEU:O	1:A:302:VAL:HB	2.14	0.48
1:A:35:GLN:O	1:A:38:GLN:HG2	2.13	0.48
1:A:466:SER:O	1:A:468:LEU:N	2.43	0.48
1:A:31:ASP:O	1:A:34:THR:HG22	2.13	0.47
1:A:41:GLU:OE2	1:A:473:HIS:NE2	2.35	0.47
1:A:320:LEU:HA	1:A:321:ASP:HA	1.66	0.47
1:A:282:ASP:OD2	1:A:314:LLP:C2'	2.63	0.47
1:A:313:HIS:HB2	1:A:320:LEU:HD22	1.97	0.47
1:A:39:THR:HG22	1:A:43:ARG:HB2	1.96	0.46
1:A:165:ALA:O	1:A:169:THR:HG23	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:GLY:HA2	1:A:330:PRO:HD3	1.96	0.46
1:A:229:CYS:HB3	1:A:232:ASP:OD1	2.16	0.46
1:A:227:LYS:HB3	1:A:264:PRO:HD3	1.97	0.45
1:A:138:MET:O	1:A:142:LEU:HG	2.16	0.45
1:A:131:GLU:OE2	1:A:362:ARG:NH1	2.48	0.45
1:A:315:TRP:CD1	1:A:385:ILE:HG12	2.52	0.45
1:A:49:VAL:HG12	1:A:50:LYS:N	2.31	0.45
1:A:119:PHE:HB3	1:A:360:LEU:O	2.16	0.44
1:A:161:ASN:HB2	1:A:314:LLP:OP1	2.18	0.44
1:A:192:TYR:HB2	1:A:250:VAL:HG22	2.00	0.44
1:A:170:LEU:HB3	1:A:208:ILE:HD13	1.98	0.44
1:A:69:MET:HE2	1:A:73:LEU:HG	2.00	0.43
1:A:21:GLU:O	1:A:25:GLU:HG3	2.19	0.43
1:A:135:VAL:HG13	1:A:326:TRP:NE1	2.34	0.43
1:A:164:GLU:HG2	1:A:358:ILE:HD12	2.01	0.43
1:A:74:SER:O	1:A:78:GLU:HG2	2.19	0.43
1:A:80:ILE:O	1:A:84:LEU:HG	2.19	0.42
1:A:40:LEU:HB3	1:A:89:SER:HA	2.02	0.42
1:A:141:MET:HE3	1:A:378:VAL:HG22	2.00	0.42
1:A:49:VAL:HG13	1:A:50:LYS:HD3	2.01	0.42
1:A:415:LEU:HD13	1:A:463:VAL:HG22	2.02	0.42
1:A:376:TYR:O	1:A:380[B]:ASN:ND2	2.53	0.42
1:A:177:ALA:O	1:A:181:VAL:HG22	2.20	0.41
1:A:17:MET:O	1:A:19:LEU:HD22	2.21	0.41
1:A:267:GLU:OE1	1:A:270:LYS:HE2	2.20	0.41
1:A:284:ALA:CB	1:A:314:LLP:C2'	2.91	0.41
1:A:96:PHE:CE1	1:A:257:THR:HA	2.55	0.41
1:A:192:TYR:CE1	1:A:247:PRO:HB3	2.47	0.41
1:A:282:ASP:OD2	1:A:314:LLP:H2'3	2.22	0.40
1:A:167:LEU:HD11	1:A:207:LEU:HD13	2.03	0.40
1:A:188:LYS:HG2	1:A:245:LEU:HD23	2.04	0.40
1:A:266:LYS:O	1:A:270:LYS:HG3	2.21	0.40
1:A:96:PHE:HE1	1:A:257:THR:HA	1.86	0.40
1:A:284:ALA:HB1	1:A:314:LLP:C2'	2.52	0.40
1:A:145:PRO:HD2	1:A:148:PHE:HD1	1.86	0.40
1:A:289:ALA:HB2	1:A:315:TRP:CZ3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	465/490 (95%)	442 (95%)	19 (4%)	4 (1%)	14	30

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	315	TRP
1	A	465	GLY
1	A	124	SER
1	A	455	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	407/423 (96%)	407 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	ASN
1	A	231	ASN
1	A	357	GLN
1	A	384	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	314	1	23,24,25	2.51	5 (21%)	25,32,34	1.50	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	314	1	-	9/16/17/19	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	314	LLP	C3-C2	7.61	1.48	1.41
1	A	314	LLP	C4'-NZ	5.44	1.45	1.27
1	A	314	LLP	C4-C5	4.91	1.48	1.42
1	A	314	LLP	C4-C3	4.45	1.48	1.41
1	A	314	LLP	C4-C4'	2.27	1.51	1.46

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	314	LLP	C4-C3-C2	-4.70	117.50	120.14
1	A	314	LLP	C4-C4'-NZ	-3.15	109.52	124.04
1	A	314	LLP	C6-N1-C2	2.22	123.22	119.20

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	314	LLP	C4-C4'-NZ-CE
1	A	314	LLP	C5'-OP4-P-OP1
1	A	314	LLP	C5'-OP4-P-OP2
1	A	314	LLP	C5'-OP4-P-OP3
1	A	314	LLP	C-CA-CB-CG
1	A	314	LLP	CG-CD-CE-NZ
1	A	314	LLP	C3-C4-C4'-NZ
1	A	314	LLP	CD-CE-NZ-C4'
1	A	314	LLP	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	314	LLP	8	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	466/490 (95%)	0.67	28 (6%) 27 22	21, 64, 95, 138	3 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	163	CYS	4.5
1	A	339	ALA	3.5
1	A	337	LEU	3.4
1	A	199	PHE	3.1
1	A	336	SER	2.9
1	A	326	TRP	2.9
1	A	322	CYS	2.9
1	A	265	ILE	2.7
1	A	318	SER	2.7
1	A	96	PHE	2.6
1	A	298	TYR	2.6
1	A	238	ARG	2.6
1	A	361	SER	2.5
1	A	117	ILE	2.5
1	A	327	LEU	2.5
1	A	166	VAL	2.5
1	A	155	GLY	2.4
1	A	202[A]	HIS	2.3
1	A	466	SER	2.3
1	A	164	GLU	2.3
1	A	360	LEU	2.3
1	A	358	ILE	2.3
1	A	372	VAL	2.3
1	A	354	LYS	2.2
1	A	366	ALA	2.2
1	A	60	ASP	2.1
1	A	123	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	353	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q < 0.9
1	LLP	A	314	24/25	0.77	0.20	70,106,120,122	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.