



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:01 PM UTC

PDB ID : 3EF1 / pdb_00003ef1
Title : The Structure of Fcp1, an essential RNA polymerase II CTD phosphatase
Authors : Ghosh, A.; Lima, C.D.
Deposited on : 2008-09-07
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	442	14% 56% 25% • 16%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3143 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA polymerase II subunit A C-terminal domain phosphatase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	372	Total	Be	C	F	N	O	S	0	0	0
			2979	1	1890	3	513	563	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	139	SER	-	expression tag	UNP Q9P376

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	163	Total	O	0	0
			163	163		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RNA polymerase II subunit A C-terminal domain phosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.66Å 88.44Å 113.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.15 30.00 – 2.15	Depositor EDS
% Data completeness (in resolution range)	95.7 (30.00-2.15) 95.7 (30.00-2.15)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 2.16Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.211 , 0.234 0.209 , 0.205	Depositor DCC
R_{free} test set	1448 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	38.0	Xtriage
Anisotropy	0.246	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3143	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BFD, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3023	0.92	14/4086 (0.3%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	467	ILE	N-CA-C	-7.89	104.89	111.91
1	A	209	GLU	N-CA-C	7.04	119.64	109.14
1	A	490	VAL	N-CA-C	6.85	117.36	110.23
1	A	265	GLN	CB-CA-C	-6.62	108.95	116.63
1	A	579	LEU	N-CA-C	-5.99	106.51	113.88
1	A	180	VAL	N-CA-C	-5.57	107.30	112.43
1	A	462	GLU	N-CA-C	-5.49	104.70	111.40
1	A	239	LEU	N-CA-C	5.46	119.14	109.96
1	A	508	VAL	N-CA-C	5.36	115.49	110.30
1	A	162	GLU	N-CA-C	-5.19	106.95	113.28
1	A	293	VAL	N-CA-C	5.19	115.32	107.80
1	A	196	TYR	N-CA-C	5.12	116.55	111.07
1	A	210	GLY	CA-C-N	5.08	126.18	119.84
1	A	210	GLY	C-N-CA	5.08	126.18	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2979	0	3004	90	0
2	A	1	0	0	0	0
3	A	163	0	0	3	0
All	All	3143	0	3004	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:ASN:HD21	1:A:422:GLN:HE21	1.04	0.94
1:A:180:VAL:H	1:A:412:GLN:HE21	1.22	0.86
1:A:294:VAL:HG22	1:A:479:VAL:HG21	1.57	0.84
1:A:180:VAL:H	1:A:412:GLN:NE2	1.80	0.78
1:A:478:ASN:ND2	1:A:480:GLY:H	1.83	0.76
1:A:553:ASN:HD22	1:A:553:ASN:H	1.34	0.75
1:A:467:ILE:HG12	1:A:475:LYS:O	1.89	0.73
1:A:195:ASN:HD21	1:A:422:GLN:NE2	1.87	0.65
1:A:478:ASN:C	1:A:478:ASN:HD22	2.06	0.64
1:A:153:SER:O	1:A:157:LYS:HG2	1.98	0.64
1:A:478:ASN:HD22	1:A:480:GLY:H	1.46	0.64
1:A:488:GLN:O	1:A:492:LYS:HD3	1.98	0.63
1:A:464:GLU:OE1	1:A:476:HIS:HD2	1.84	0.60
1:A:247:LYS:HE3	1:A:251:LYS:CE	2.33	0.59
1:A:541:ARG:HG2	1:A:541:ARG:HH11	1.66	0.59
1:A:195:ASN:ND2	1:A:422:GLN:HE21	1.88	0.59
1:A:265:GLN:HG3	1:A:267:ARG:HH12	1.68	0.59
1:A:553:ASN:HD22	1:A:553:ASN:N	1.97	0.58
1:A:240:HIS:CE1	1:A:267:ARG:HD2	2.39	0.58
1:A:481:LEU:O	1:A:485:LYS:HG2	2.04	0.57
1:A:459:VAL:O	1:A:463:GLU:HG3	2.04	0.57
1:A:247:LYS:HE3	1:A:251:LYS:HE3	1.85	0.57
1:A:158:ARG:O	1:A:162:GLU:HG3	2.05	0.57
1:A:397:ALA:HA	1:A:400:GLU:OE1	2.05	0.56
1:A:236:LEU:CD1	1:A:454:LYS:HD3	2.36	0.56
1:A:433:THR:HB	1:A:434:PRO:HD3	1.88	0.56
1:A:504:LEU:O	1:A:504:LEU:HD23	2.05	0.56
1:A:221:LYS:HG2	1:A:317:PHE:CG	2.42	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:247:LYS:HG2	1:A:251:LYS:HE3	1.88	0.54
1:A:507:ASP:HB3	1:A:510:SER:OG	2.07	0.54
1:A:422:GLN:HE22	1:A:438:LEU:H	1.57	0.53
1:A:450:GLU:O	1:A:454:LYS:HG2	2.08	0.53
1:A:553:ASN:H	1:A:553:ASN:ND2	2.04	0.53
1:A:151:LEU:HA	3:A:615:HOH:O	2.09	0.52
1:A:429:GLU:H	1:A:429:GLU:CD	2.18	0.52
1:A:461:TYR:HA	1:A:464:GLU:HB3	1.92	0.52
1:A:488:GLN:HE21	1:A:521:GLY:HA2	1.74	0.51
1:A:236:LEU:HD12	1:A:454:LYS:HD3	1.90	0.51
1:A:200:ARG:HG2	1:A:200:ARG:HH11	1.76	0.51
1:A:271:ARG:NH2	1:A:402:ASN:OD1	2.37	0.51
1:A:410:GLU:HA	1:A:410:GLU:OE1	2.11	0.50
1:A:472:GLY:H	1:A:474:HIS:HD2	1.59	0.50
1:A:409:LEU:O	1:A:413:LYS:HG3	2.11	0.50
1:A:265:GLN:HE21	1:A:265:GLN:HA	1.77	0.50
1:A:402:ASN:O	1:A:406:VAL:HG23	2.12	0.49
1:A:431:LYS:HB3	1:A:432:PRO:HD3	1.95	0.48
1:A:478:ASN:ND2	1:A:480:GLY:N	2.59	0.48
1:A:297:ASP:C	1:A:312:VAL:HG12	2.39	0.48
1:A:207:LEU:O	1:A:215:THR:HA	2.14	0.48
1:A:207:LEU:HD11	1:A:249:TYR:HA	1.96	0.48
1:A:483:ILE:HB	1:A:484:PRO:HD3	1.96	0.48
1:A:528:PHE:O	1:A:529:SER:C	2.56	0.48
1:A:399:GLU:HA	1:A:402:ASN:HD22	1.79	0.47
1:A:221:LYS:HG2	1:A:317:PHE:CD2	2.50	0.46
1:A:208:GLN:HA	1:A:214:TYR:O	2.15	0.46
1:A:155:ASN:HB3	3:A:732:HOH:O	2.15	0.46
1:A:228:GLN:NE2	1:A:228:GLN:HA	2.31	0.45
1:A:497:LEU:O	1:A:535:LEU:HD12	2.15	0.45
1:A:294:VAL:CG2	1:A:479:VAL:HG21	2.36	0.45
1:A:139:SER:C	1:A:141:LEU:H	2.24	0.45
1:A:151:LEU:HG	3:A:616:HOH:O	2.15	0.45
1:A:417:PRO:O	1:A:421:GLN:HG3	2.17	0.45
1:A:478:ASN:HD22	1:A:479:VAL:N	2.14	0.44
1:A:429:GLU:CD	1:A:429:GLU:N	2.75	0.44
1:A:538:ALA:O	1:A:540:ILE:HG13	2.17	0.44
1:A:197:ASP:HA	1:A:200:ARG:HD3	2.00	0.44
1:A:422:GLN:NE2	1:A:438:LEU:H	2.16	0.43
1:A:431:LYS:H	1:A:432:PRO:CD	2.32	0.43
1:A:204:SER:HA	1:A:218:TYR:O	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:478:ASN:ND2	1:A:478:ASN:C	2.73	0.43
1:A:141:LEU:O	1:A:145:LEU:HG	2.19	0.43
1:A:312:VAL:HG22	1:A:313:VAL:N	2.32	0.43
1:A:169:VAL:HG22	1:A:296:ILE:HD11	2.01	0.43
1:A:191:PRO:HA	1:A:196:TYR:CG	2.54	0.42
1:A:428:ASP:O	1:A:429:GLU:C	2.61	0.42
1:A:469:SER:O	1:A:470:ARG:HB2	2.20	0.42
1:A:155:ASN:HD22	1:A:155:ASN:HA	1.61	0.41
1:A:180:VAL:N	1:A:412:GLN:HE21	2.03	0.41
1:A:295:VAL:HG11	1:A:303:TRP:CE2	2.55	0.41
1:A:177:HIS:CE1	1:A:318:PHE:HB2	2.56	0.41
1:A:228:GLN:NE2	1:A:232:LYS:NZ	2.69	0.41
1:A:233:ILE:HG23	1:A:239:LEU:HD21	2.02	0.41
1:A:244:MET:HE1	1:A:277:LEU:HA	2.03	0.41
1:A:309:LEU:O	1:A:487:LYS:NZ	2.49	0.41
1:A:426:LEU:C	1:A:428:ASP:H	2.29	0.41
1:A:432:PRO:O	1:A:436:HIS:ND1	2.54	0.41
1:A:264:PHE:O	1:A:265:GLN:HB2	2.22	0.40
1:A:528:PHE:CE2	1:A:547:LYS:HG2	2.56	0.40
1:A:467:ILE:O	1:A:467:ILE:HG23	2.20	0.40
1:A:460:TYR:CE1	1:A:477:ALA:HB3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	367/442 (83%)	334 (91%)	24 (6%)	9 (2%)	4 1

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	211	PRO
1	A	469	SER
1	A	210	GLY
1	A	468	SER
1	A	552	GLY
1	A	140	ASP
1	A	472	GLY
1	A	431	LYS
1	A	432	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	332/401 (83%)	319 (96%)	13 (4%)	28	28

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	151	LEU
1	A	203	ARG
1	A	211	PRO
1	A	231	GLN
1	A	234	SER
1	A	265	GLN
1	A	410	GLU
1	A	418	LEU
1	A	429	GLU
1	A	466	ASP
1	A	488	GLN
1	A	553	ASN
1	A	571	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	155	ASN
1	A	206	ASN
1	A	208	GLN
1	A	228	GLN
1	A	240	HIS
1	A	265	GLN
1	A	279	GLN
1	A	412	GLN
1	A	422	GLN
1	A	423	ASN
1	A	441	ASN
1	A	474	HIS
1	A	476	HIS
1	A	478	ASN
1	A	488	GLN
1	A	553	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	BFD	A	170	1	8,11,12	3.41	3 (37%)	2,15,17	2.28	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	BFD	A	170	1	-	0/5/11/13	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	170	BFD	F2-BE	-6.25	1.39	1.54
1	A	170	BFD	F3-BE	-5.58	1.41	1.54
1	A	170	BFD	F1-BE	-4.31	1.44	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	170	BFD	OD2-CG-CB	-3.22	117.16	124.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	371/442 (83%)	0.76	62 (16%) 4 5	25, 44, 99, 117	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	467	ILE	8.1
1	A	432	PRO	7.6
1	A	468	SER	6.1
1	A	580	TYR	5.9
1	A	474	HIS	5.9
1	A	213	GLY	5.8
1	A	143	VAL	5.2
1	A	473	ASN	4.7
1	A	471	SER	4.6
1	A	579	LEU	4.6
1	A	397	ALA	4.6
1	A	406	VAL	4.3
1	A	212	SER	3.7
1	A	211	PRO	3.7
1	A	433	THR	3.6
1	A	141	LEU	3.6
1	A	144	SER	3.5
1	A	326	SER	3.5
1	A	431	LYS	3.5
1	A	503	PRO	3.4
1	A	434	PRO	3.4
1	A	578	LEU	3.2
1	A	407	THR	3.1
1	A	530	VAL	3.0
1	A	466	ASP	3.0
1	A	148	ALA	2.9
1	A	139	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	147	GLU	2.8
1	A	462	GLU	2.8
1	A	145	LEU	2.8
1	A	504	LEU	2.8
1	A	208	GLN	2.7
1	A	402	ASN	2.7
1	A	430	GLY	2.7
1	A	151	LEU	2.7
1	A	214	TYR	2.7
1	A	325	ASN	2.7
1	A	465	ASN	2.6
1	A	398	LEU	2.6
1	A	401	GLN	2.5
1	A	472	GLY	2.5
1	A	215	THR	2.5
1	A	538	ALA	2.5
1	A	529	SER	2.4
1	A	475	LYS	2.4
1	A	492	LYS	2.3
1	A	441	ASN	2.3
1	A	405	ARG	2.3
1	A	288	CYS	2.3
1	A	507	ASP	2.2
1	A	505	GLY	2.2
1	A	153	SER	2.2
1	A	560	ASN	2.2
1	A	207	LEU	2.2
1	A	454	LYS	2.2
1	A	210	GLY	2.1
1	A	228	GLN	2.1
1	A	469	SER	2.1
1	A	470	ARG	2.1
1	A	403	LYS	2.0
1	A	539	LYS	2.0
1	A	321	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	BFD	A	170	12/13	0.94	0.09	23,27,28,28	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MG	A	1	1/1	0.98	0.13	34,34,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.