



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:49 PM UTC

PDB ID : 4EF4 / pdb\_00004ef4  
Title : Crystal structure of STING CTD complex with c-di-GMP  
Authors : Ouyang, S.; Ru, H.; Shaw, N.; Jiang, Y.; Niu, F.; Zhu, Y.; Qiu, W.; Li, Y.;  
Liu, Z.-J.  
Deposited on : 2012-03-29  
Resolution : 2.15 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	
1	B	265	

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3063 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transmembrane protein 173.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	178	1460	923	258	273	6	0	1	0
1	B	177	1456	919	254	277	6	6	4	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	115	MET	-	expression tag	UNP Q86WV6
A	116	HIS	-	expression tag	UNP Q86WV6
A	117	HIS	-	expression tag	UNP Q86WV6
A	118	HIS	-	expression tag	UNP Q86WV6
A	119	HIS	-	expression tag	UNP Q86WV6
A	120	HIS	-	expression tag	UNP Q86WV6
A	121	HIS	-	expression tag	UNP Q86WV6
A	122	SER	-	expression tag	UNP Q86WV6
A	123	SER	-	expression tag	UNP Q86WV6
A	124	GLY	-	expression tag	UNP Q86WV6
A	125	VAL	-	expression tag	UNP Q86WV6
A	126	ASP	-	expression tag	UNP Q86WV6
A	127	LEU	-	expression tag	UNP Q86WV6
A	128	GLY	-	expression tag	UNP Q86WV6
A	129	THR	-	expression tag	UNP Q86WV6
A	130	GLU	-	expression tag	UNP Q86WV6
A	131	ASN	-	expression tag	UNP Q86WV6
A	132	LEU	-	expression tag	UNP Q86WV6
A	133	TYR	-	expression tag	UNP Q86WV6
A	134	PHE	-	expression tag	UNP Q86WV6
A	135	GLN	-	expression tag	UNP Q86WV6
A	136	SER	-	expression tag	UNP Q86WV6
A	137	ASN	-	expression tag	UNP Q86WV6
A	138	ALA	-	expression tag	UNP Q86WV6
B	115	MET	-	expression tag	UNP Q86WV6

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	116	HIS	-	expression tag	UNP Q86WV6
B	117	HIS	-	expression tag	UNP Q86WV6
B	118	HIS	-	expression tag	UNP Q86WV6
B	119	HIS	-	expression tag	UNP Q86WV6
B	120	HIS	-	expression tag	UNP Q86WV6
B	121	HIS	-	expression tag	UNP Q86WV6
B	122	SER	-	expression tag	UNP Q86WV6
B	123	SER	-	expression tag	UNP Q86WV6
B	124	GLY	-	expression tag	UNP Q86WV6
B	125	VAL	-	expression tag	UNP Q86WV6
B	126	ASP	-	expression tag	UNP Q86WV6
B	127	LEU	-	expression tag	UNP Q86WV6
B	128	GLY	-	expression tag	UNP Q86WV6
B	129	THR	-	expression tag	UNP Q86WV6
B	130	GLU	-	expression tag	UNP Q86WV6
B	131	ASN	-	expression tag	UNP Q86WV6
B	132	LEU	-	expression tag	UNP Q86WV6
B	133	TYR	-	expression tag	UNP Q86WV6
B	134	PHE	-	expression tag	UNP Q86WV6
B	135	GLN	-	expression tag	UNP Q86WV6
B	136	SER	-	expression tag	UNP Q86WV6
B	137	ASN	-	expression tag	UNP Q86WV6
B	138	ALA	-	expression tag	UNP Q86WV6

- Molecule 2 is 9,9'-[(2R,3R,3aS,5S,7aR,9R,10R,10aS,12S,14aR)-3,5,10,12-tetrahydroxy-5,12-dioxidoctahydro-2H,7H-difuro[3,2-d:3',2'-j][1,3,7,9,2,8]tetraoxadiphosphacyclododecine-2,9-diyl]bis(2-amino-1,9-dihydro-6H-purin-6-one) (CCD ID: C2E) (formula: C<sub>20</sub>H<sub>24</sub>N<sub>10</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	46	20	10	14	2	0	0

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	1	1	1	0	0
3	B	1	1	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	50	50	50	0	0
4	B	49	49	49	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.38Å 72.84Å 62.44Å 90.00° 97.03° 90.00°	Depositor
Resolution (Å)	41.01 – 2.15 41.01 – 2.15	Depositor EDS
% Data completeness (in resolution range)	96.8 (41.01-2.15) 85.2 (41.01-2.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 2.14Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, $R_{free}$	0.209 , 0.257 0.215 , 0.259	Depositor DCC
$R_{free}$ test set	1455 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.1	Xtrriage
Anisotropy	0.389	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 42.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.276 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3063	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: C2E, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/1490	0.85	3/2017 (0.1%)
1	B	0.47	0/1494	0.84	4/2022 (0.2%)
All	All	0.48	0/2984	0.84	7/4039 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	172	LEU	CA-C-N	-7.67	112.10	119.85
1	B	172	LEU	C-N-CA	-7.67	112.10	119.85
1	A	306	GLN	N-CA-C	7.61	119.36	111.14
1	A	302	ALA	CA-C-N	6.23	125.64	118.85
1	A	302	ALA	C-N-CA	6.23	125.64	118.85
1	B	302	ALA	CA-C-N	5.29	126.45	119.84
1	B	302	ALA	C-N-CA	5.29	126.45	119.84

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1460	0	1439	18	0
1	B	1456	0	1435	18	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	46	0	22	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	50	0	0	3	0
4	B	49	0	0	3	0
All	All	3063	0	2896	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:TYR:OH	1:A:336:GLU:OE1	2.03	0.76
1:B:199:TYR:OH	1:B:336:GLU:OE1	2.04	0.75
1:A:197:ARG:HH21	1:A:307:ASN:HB3	1.60	0.66
1:A:305:SER:OG	1:A:307:ASN:ND2	2.28	0.65
1:A:165:ILE:HA	1:A:169[A]:ARG:HD2	1.80	0.64
1:B:178:ARG:HH11	1:B:227:GLN:HA	1.64	0.62
1:B:273:GLN:O	4:B:521:HOH:O	2.17	0.60
1:B:301:ASP:N	1:B:301:ASP:OD1	2.35	0.60
1:A:155:VAL:HG23	1:B:158:GLY:HA2	1.84	0.59
1:B:306:GLN:O	1:B:309:CYS:N	2.36	0.59
1:A:174:GLU:HG2	4:A:547:HOH:O	2.07	0.54
1:B:190:LEU:O	1:B:192:GLY:N	2.43	0.52
1:A:247:LEU:HD21	1:A:329:VAL:HB	1.93	0.51
1:A:306:GLN:N	1:A:307:ASN:HA	2.25	0.51
1:A:211:ASN:HB3	1:A:214:MET:HG3	1.94	0.50
1:A:246:GLU:OE2	1:A:253:ARG:NH1	2.40	0.50
1:B:305:SER:HB3	1:B:307:ASN:H	1.77	0.50
1:A:176:GLN:OE1	1:A:306:GLN:NE2	2.43	0.49
1:B:171:ILE:O	1:B:175:LEU:HB2	2.12	0.49
1:A:308:ASN:OD1	1:A:308:ASN:N	2.41	0.49
1:B:199:TYR:HE2	1:B:310:ARG:HD3	1.78	0.48
1:B:284:ARG:HD2	4:B:528:HOH:O	2.13	0.48
1:B:306:GLN:O	1:B:308:ASN:N	2.47	0.48
1:A:209:PRO:HD2	1:A:261:TYR:CE2	2.50	0.47
1:B:178:ARG:HD2	1:B:227:GLN:HA	1.97	0.46
1:B:174:GLU:O	1:B:178:ARG:HG3	2.16	0.46
1:A:338:LYS:O	1:A:342:THR:HB	2.17	0.45
1:B:210:ASP:OD1	4:B:527:HOH:O	2.21	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:197:ARG:HG2	1:B:308:ASN:HA	1.99	0.44
1:A:280:SER:OG	1:A:282:GLU:HG2	2.17	0.44
1:A:284:ARG:HD2	4:A:511:HOH:O	2.18	0.44
1:A:223:ASP:OD1	4:A:543:HOH:O	2.20	0.43
1:A:199:TYR:HD2	1:A:312:ILE:HD11	1.83	0.43
1:B:211:ASN:HB3	1:B:214:MET:HE3	2.02	0.42
1:B:199:TYR:CD2	1:B:312:ILE:HD11	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/265 (65%)	167 (96%)	6 (4%)	0	100	100
1	B	175/265 (66%)	164 (94%)	8 (5%)	3 (2%)	7	2
All	All	348/530 (66%)	331 (95%)	14 (4%)	3 (1%)	14	8

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	191	ARG
1	B	304	GLU
1	B	307	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/231 (69%)	146 (92%)	13 (8%)	10	6
1	B	160/231 (69%)	147 (92%)	13 (8%)	11	6
All	All	319/462 (69%)	293 (92%)	26 (8%)	11	6

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	159	LEU
1	A	171	ILE
1	A	212	LEU
1	A	223	ASP
1	A	227	GLN
1	A	239	VAL
1	A	253	ARG
1	A	272	SER
1	A	282	GLU
1	A	301	ASP
1	A	307	ASN
1	A	308	ASN
1	A	342	THR
1	B	162	SER
1	B	174	GLU
1	B	196	GLN
1	B	211	ASN
1	B	272[A]	SER
1	B	272[B]	SER
1	B	301	ASP
1	B	304	GLU
1	B	305	SER
1	B	322	SER
1	B	325	LEU
1	B	330	LEU
1	B	333	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	185	HIS
1	A	211	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	307	ASN
1	B	157	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	C2E	A	401	-	52,52,52	2.58	12 (23%)	78,82,82	1.83	20 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C2E	A	401	-	-	1/30/62/62	0/6/7/7

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	C2E	O6-C6	9.60	1.41	1.23
2	A	401	C2E	O61-C61	8.51	1.39	1.23
2	A	401	C2E	C2'-C3'	-6.46	1.38	1.53
2	A	401	C2E	C2-N2	4.91	1.45	1.34
2	A	401	C2E	C5A-C4A	-4.30	1.38	1.51
2	A	401	C2E	C21-N21	3.34	1.42	1.34
2	A	401	C2E	C3'-C4'	-2.73	1.45	1.52
2	A	401	C2E	O4'-C4'	-2.62	1.39	1.45
2	A	401	C2E	C2A-C3A	-2.57	1.47	1.53
2	A	401	C2E	O2A-C2A	-2.46	1.36	1.43
2	A	401	C2E	C6-N1	-2.36	1.34	1.38
2	A	401	C2E	O4A-C4A	2.15	1.49	1.45

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	C2E	C5-C4-N3	-5.45	119.72	128.39
2	A	401	C2E	C51-C41-N31	-4.74	120.85	128.39
2	A	401	C2E	C2-N3-C4	4.16	119.47	112.30
2	A	401	C2E	C21-N31-C41	4.07	119.31	112.30
2	A	401	C2E	N9-C4-N3	3.87	133.70	125.95
2	A	401	C2E	C61-C51-N71	3.15	136.01	130.29
2	A	401	C2E	C41-C51-N71	-3.10	105.76	110.67
2	A	401	C2E	N9-C8-N7	-3.00	107.83	113.40
2	A	401	C2E	C2-N1-C6	-2.90	119.85	125.11
2	A	401	C2E	C81-N71-C51	2.79	109.23	104.26
2	A	401	C2E	N91-C81-N71	-2.71	108.37	113.40
2	A	401	C2E	O6-C6-C5	-2.53	119.86	126.53
2	A	401	C2E	C8-N7-C5	2.52	108.74	104.26
2	A	401	C2E	N91-C41-N31	2.49	130.94	125.95
2	A	401	C2E	C5-C6-N1	2.45	119.49	113.25
2	A	401	C2E	C21-N11-C61	-2.41	120.74	125.11
2	A	401	C2E	C4-C5-N7	-2.29	107.05	110.67
2	A	401	C2E	C51-C61-N11	2.17	118.78	113.25
2	A	401	C2E	C6-C5-N7	2.13	134.17	130.29
2	A	401	C2E	C8-N9-C4	2.01	109.80	106.03

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	C2E	C3A-O3A-P1-O2P

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	178/265 (67%)	-0.93	0 <a href="#">100</a>   <a href="#">100</a>	27, 55, 78, 98	2 (1%)
1	B	177/265 (66%)	-0.94	1 (0%) <a href="#">85</a>   <a href="#">88</a>	27, 53, 82, 102	4 (2%)
All	All	355/530 (66%)	-0.93	1 (0%) <a href="#">90</a>   <a href="#">92</a>	27, 55, 81, 102	6 (1%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	318	ALA	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	C2E	A	401	46/46	0.99	0.03	37,44,54,68	0
3	CA	A	402	1/1	1.00	0.02	57,57,57,57	0
3	CA	B	401	1/1	1.00	0.01	53,53,53,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.