



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:49 PM UTC

PDB ID : 4EF5 / pdb_00004ef5
Title : Crystal structure of STING CTD
Authors : Ouyang, S.; Ru, H.; Shaw, N.; Jiang, Y.; Niu, F.; Zhu, Y.; Qiu, W.; Li, Y.;
Liu, Z.-J.
Deposited on : 2012-03-29
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1476 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transmembrane protein 173.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	177	Total	C	N	O	S	0	3	0
			1455	920	256	273	6			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	115	MET	-	expression tag	UNP Q86WV6
A	116	HIS	-	expression tag	UNP Q86WV6
A	117	HIS	-	expression tag	UNP Q86WV6
A	118	HIS	-	expression tag	UNP Q86WV6
A	119	HIS	-	expression tag	UNP Q86WV6
A	120	HIS	-	expression tag	UNP Q86WV6
A	121	HIS	-	expression tag	UNP Q86WV6
A	122	SER	-	expression tag	UNP Q86WV6
A	123	SER	-	expression tag	UNP Q86WV6
A	124	GLY	-	expression tag	UNP Q86WV6
A	125	VAL	-	expression tag	UNP Q86WV6
A	126	ASP	-	expression tag	UNP Q86WV6
A	127	LEU	-	expression tag	UNP Q86WV6
A	128	GLY	-	expression tag	UNP Q86WV6
A	129	THR	-	expression tag	UNP Q86WV6
A	130	GLU	-	expression tag	UNP Q86WV6
A	131	ASN	-	expression tag	UNP Q86WV6
A	132	LEU	-	expression tag	UNP Q86WV6
A	133	TYR	-	expression tag	UNP Q86WV6
A	134	PHE	-	expression tag	UNP Q86WV6
A	135	GLN	-	expression tag	UNP Q86WV6
A	136	SER	-	expression tag	UNP Q86WV6
A	137	ASN	-	expression tag	UNP Q86WV6
A	138	ALA	-	expression tag	UNP Q86WV6

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	21	Total	O	0	0
			21	21		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transmembrane protein 173



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	81.04Å 90.36Å 73.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.18 – 2.45 45.18 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.18-2.45) 99.9 (45.18-2.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.20 (at 2.45Å)	Xtriage
Refinement program	PHENIX 1.7.3 _928	Depositor
R, R_{free}	0.227 , 0.288 0.230 , 0.288	Depositor DCC
R_{free} test set	489 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	59.8	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1476	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/1491	0.77	0/2018

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1455	0	1437	19	1
2	A	21	0	0	7	1
All	All	1476	0	1437	19	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:TYR:O	2:A:402:HOH:O	1.91	0.86
1:A:314:TYR:OH	2:A:414:HOH:O	2.03	0.74
1:A:205:ASP:OD1	2:A:418:HOH:O	2.07	0.72
1:A:246:GLU:OE2	1:A:253:ARG:NH1	2.30	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:ASP:OD2	2:A:419:HOH:O	2.14	0.64
1:A:153:PHE:O	2:A:421:HOH:O	2.16	0.59
1:A:199:TYR:OH	1:A:336:GLU:OE1	2.22	0.57
1:A:324:SER:HB3	1:A:327:GLN:HB2	1.91	0.52
1:A:303:PRO:C	1:A:305:SER:H	2.18	0.50
1:A:308:ASN:OD1	1:A:308:ASN:N	2.43	0.49
1:A:284:ARG:HD2	2:A:410:HOH:O	2.16	0.45
1:A:203:PRO:HB2	1:A:205:ASP:HB2	1.99	0.45
1:A:276:GLN:OE1	1:A:276:GLN:N	2.36	0.44
1:A:216:ASP:HB3	1:A:219:ILE:HG13	2.00	0.43
1:A:220:ARG:NH1	1:A:221:PHE:O	2.53	0.42
1:A:206:CYS:SG	1:A:284:ARG:HD3	2.60	0.41
1:A:303:PRO:C	1:A:305:SER:N	2.78	0.41
1:A:334:ARG:HD2	2:A:416:HOH:O	2.20	0.41
1:A:303:PRO:O	1:A:305:SER:N	2.54	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:341:VAL:O	2:A:419:HOH:O[6_555]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	174/265 (66%)	161 (92%)	12 (7%)	1 (1%)	21 27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/231 (69%)	146 (92%)	13 (8%)	10	13

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	170	LEU
1	A	171	ILE
1	A	175	LEU
1	A	212	LEU
1	A	224	LYS
1	A	272	SER
1	A	282	GLU
1	A	306	GLN
1	A	307	ASN
1	A	308	ASN
1	A	309	CYS
1	A	325	LEU
1	A	330	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	306	GLN
1	A	327	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	177/265 (66%)	0.82	21 (11%) 9 7	29, 60, 96, 113	3 (1%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	190	LEU	6.9
1	A	240	TYR	4.9
1	A	318	ALA	4.6
1	A	214	MET	4.2
1	A	185	HIS	3.7
1	A	303	PRO	3.3
1	A	305	SER	3.3
1	A	153	PHE	3.2
1	A	307	ASN	3.1
1	A	189	LEU	2.8
1	A	186	TYR	2.7
1	A	306	GLN	2.7
1	A	152	ASN	2.6
1	A	215	ALA	2.6
1	A	323	PHE	2.5
1	A	322	SER	2.5
1	A	321	SER	2.4
1	A	171	ILE	2.3
1	A	343	VAL	2.2
1	A	197[A]	ARG	2.1
1	A	310	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.