



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:40 PM UTC

PDB ID : 7EFR / pdb_00007efr
Title : Structure of SARS-CoV-2 spike receptor-binding domain in complex with high affinity ACE2 mutant (T27W,N330Y)
Authors : Lu, G.W.; Ye, F.; Lin, X.
Deposited on : 2021-03-23
Resolution : 2.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	599	2% 83% 16% .
2	B	217	5% 77% 12% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	1003	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6609 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Processed angiotensin-converting enzyme 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	596	Total	C	N	O	S	0	0	0
			4859	3115	801	914	29			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	ALA	-	expression tag	UNP Q9BYF1
A	18	ASP	-	expression tag	UNP Q9BYF1
A	27	TRP	THR	engineered mutation	UNP Q9BYF1
A	330	TYR	ASN	engineered mutation	UNP Q9BYF1

- Molecule 2 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	194	Total	C	N	O	S	0	0	0
			1536	985	256	287	8			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	109	Total	O	0	0
			109	109		
4	B	35	Total	O	0	0
			35	35		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Processed angiotensin-converting enzyme 2



- Molecule 2: Spike glycoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	139.58Å 139.58Å 156.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.28 – 2.49 26.28 – 2.49	Depositor EDS
% Data completeness (in resolution range)	99.7 (26.28-2.49) 99.5 (26.28-2.49)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 2.50Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.190 , 0.229 0.193 , 0.229	Depositor DCC
R_{free} test set	2700 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	50.1	Xtriage
Anisotropy	0.379	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6609	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/4997	0.55	3/6791 (0.0%)
2	B	0.39	0/1580	0.58	0/2151
All	All	0.36	0/6577	0.56	3/8942 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	133	CYS	CA-CB-SG	-7.20	97.85	114.40
1	A	493	HIS	CA-C-N	5.91	132.83	121.54
1	A	493	HIS	C-N-CA	5.91	132.83	121.54

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	480	CYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4859	0	4627	80	0
2	B	1536	0	1452	19	0
3	A	56	0	52	11	0
3	B	14	0	13	0	0
4	A	109	0	0	8	0
4	B	35	0	0	1	0
All	All	6609	0	6144	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:GLN:OE1	3:A:1003:NAG:C6	1.69	1.37
1:A:81:GLN:CD	3:A:1003:NAG:H61	1.67	1.18
1:A:81:GLN:OE1	3:A:1003:NAG:H61	0.80	0.98
1:A:74:LYS:CE	3:A:1003:NAG:H83	1.97	0.93
1:A:74:LYS:HE2	3:A:1003:NAG:H83	1.53	0.88
1:A:455:MET:HE3	1:A:480:MET:HB2	1.58	0.84
2:B:472:ILE:HG12	2:B:482:GLY:HA2	1.61	0.82
1:A:406:GLU:HG3	1:A:518:ARG:NH2	1.96	0.81
1:A:246:ALA:HA	1:A:249:MET:HE3	1.67	0.77
1:A:74:LYS:HE3	3:A:1003:NAG:H83	1.67	0.76
1:A:515:TYR:HD1	1:A:518:ARG:HH11	1.32	0.75
1:A:291:ILE:HD11	1:A:415:PRO:HG3	1.67	0.75
1:A:402:GLU:C	1:A:518:ARG:HG3	2.13	0.73
1:A:494:ASP:HB3	1:A:496:THR:H	1.57	0.70
1:A:288:LYS:NZ	1:A:431:ASP:OD2	2.24	0.69
2:B:360:ASN:H	2:B:523:THR:HG23	1.57	0.69
1:A:432:ASN:ND2	4:A:1101:HOH:O	2.24	0.69
1:A:81:GLN:OE1	3:A:1003:NAG:C5	2.41	0.67
1:A:229:THR:HG23	1:A:516:TYR:OH	1.98	0.64
1:A:52:THR:O	1:A:340:GLN:NE2	2.31	0.63
1:A:455:MET:CE	1:A:480:MET:HB2	2.27	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:420:SER:HB2	3:A:1004:NAG:H62	1.82	0.61
1:A:406:GLU:HG3	1:A:518:ARG:HH21	1.68	0.59
2:B:379:CYS:HB3	2:B:382:VAL:HG23	1.85	0.59
1:A:419:LYS:HD2	1:A:428:PHE:HB3	1.85	0.58
1:A:455:MET:HE3	1:A:480:MET:CB	2.32	0.58
1:A:85:LEU:HA	1:A:88:ILE:HD12	1.86	0.58
1:A:74:LYS:HE2	3:A:1003:NAG:C8	2.32	0.57
1:A:253:PRO:O	1:A:254:SER:HB3	2.04	0.57
1:A:402:GLU:O	1:A:518:ARG:HG3	2.04	0.57
1:A:20:THR:HG23	1:A:23:GLU:H	1.70	0.56
1:A:187:LYS:HD2	1:A:199:TYR:CZ	2.40	0.56
1:A:461:TRP:O	1:A:465:LYS:HG2	2.05	0.56
1:A:322:ASN:ND2	4:A:1102:HOH:O	2.25	0.55
1:A:245:ARG:HB2	1:A:262:LEU:HD21	1.88	0.54
1:A:442:GLN:NE2	4:A:1105:HOH:O	2.38	0.54
2:B:472:ILE:CG1	2:B:482:GLY:HA2	2.36	0.54
1:A:114:LYS:NZ	1:A:114:LYS:HB3	2.22	0.54
1:A:403:ALA:HA	1:A:518:ARG:CG	2.38	0.54
1:A:233:ILE:HD13	1:A:450:LEU:HD13	1.89	0.53
2:B:359:SER:HA	2:B:524:VAL:CG2	2.38	0.53
1:A:25:ALA:HB1	1:A:97:LEU:HD11	1.91	0.53
1:A:474:MET:HE1	1:A:499:ASP:HB2	1.91	0.52
1:A:477:TRP:CE3	1:A:500:PRO:HG3	2.44	0.52
1:A:137:ASN:HB3	1:A:140:GLU:HG2	1.92	0.51
1:A:181:GLU:OE1	1:A:470:LYS:HE3	2.11	0.50
2:B:393:THR:O	2:B:523:THR:HG22	2.12	0.50
1:A:98:GLN:HA	1:A:101:GLN:HG2	1.94	0.50
1:A:134:ASN:HB3	1:A:137:ASN:H	1.77	0.49
1:A:134:ASN:HB2	1:A:140:GLU:CD	2.37	0.49
1:A:536:GLU:N	1:A:536:GLU:OE1	2.45	0.49
1:A:411:SER:HB3	1:A:543:ASP:HA	1.94	0.49
1:A:134:ASN:N	1:A:140:GLU:OE1	2.45	0.49
2:B:368:LEU:O	2:B:371:SER:HB3	2.12	0.49
1:A:492:PRO:HD3	1:A:613:TYR:CG	2.48	0.49
1:A:137:ASN:HB3	1:A:140:GLU:CG	2.43	0.48
1:A:228:HIS:HB2	4:A:1106:HOH:O	2.13	0.48
2:B:481:ASN:O	2:B:483:VAL:HG23	2.13	0.47
1:A:26:LYS:HE2	3:A:1002:NAG:HN2	1.78	0.47
3:A:1002:NAG:O7	3:A:1002:NAG:O3	2.29	0.47
2:B:393:THR:HG21	2:B:518:LEU:H	1.79	0.47
1:A:238:GLU:HG2	1:A:604:VAL:HG12	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:432:ASN:ND2	4:A:1109:HOH:O	2.47	0.46
2:B:384:PRO:HA	2:B:387:LEU:HD12	1.98	0.46
1:A:173:GLY:HA3	4:A:1144:HOH:O	2.15	0.46
1:A:212:VAL:HG21	1:A:565:PRO:HG3	1.97	0.45
2:B:359:SER:HA	2:B:524:VAL:HG22	1.97	0.45
2:B:450:ASN:ND2	4:B:701:HOH:O	2.44	0.45
1:A:291:ILE:HD12	1:A:291:ILE:HA	1.70	0.45
1:A:321:PRO:HD2	1:A:380:GLN:OE1	2.17	0.45
1:A:293:VAL:O	1:A:297:MET:HG2	2.17	0.45
1:A:31:LYS:NZ	1:A:35:GLU:OE1	2.45	0.45
1:A:293:VAL:HG23	1:A:297:MET:HE3	1.99	0.44
1:A:73:LEU:HD12	1:A:73:LEU:HA	1.84	0.44
1:A:388:GLN:HG3	1:A:392:LEU:HB2	2.00	0.44
1:A:310:GLU:HG3	4:A:1155:HOH:O	2.18	0.43
1:A:187:LYS:NZ	1:A:509:ASP:OD2	2.51	0.42
1:A:309:LYS:O	1:A:313:LYS:HG3	2.19	0.42
1:A:403:ALA:HA	1:A:518:ARG:HG3	2.00	0.42
1:A:119:ILE:HG21	1:A:183:TYR:HB2	2.01	0.42
1:A:245:ARG:HD2	4:A:1163:HOH:O	2.18	0.42
2:B:354:ASN:O	2:B:398:ASP:HA	2.19	0.42
1:A:403:ALA:HA	1:A:518:ARG:HG2	2.01	0.42
1:A:131:LYS:HB2	1:A:131:LYS:HE2	1.65	0.42
2:B:391:CYS:HB3	2:B:522:ALA:HB1	2.02	0.42
2:B:481:ASN:OD1	2:B:481:ASN:N	2.52	0.41
1:A:181:GLU:OE1	1:A:470:LYS:CE	2.68	0.41
1:A:168:TRP:CD1	1:A:502:SER:HB2	2.56	0.41
1:A:187:LYS:HD2	1:A:199:TYR:CE1	2.56	0.41
1:A:229:THR:OG1	1:A:581:VAL:HB	2.20	0.41
1:A:237:TYR:CZ	1:A:451:PRO:HG2	2.56	0.41
1:A:318:VAL:O	1:A:551:GLY:HA3	2.21	0.41
1:A:492:PRO:HD3	1:A:613:TYR:CD2	2.56	0.41
2:B:425:LEU:HD21	2:B:512:VAL:HG11	2.02	0.41
1:A:535:HIS:CG	1:A:536:GLU:N	2.89	0.40
2:B:481:ASN:O	2:B:482:GLY:C	2.63	0.40
1:A:165:TRP:CH2	1:A:490:PRO:HD2	2.56	0.40
2:B:392:PHE:HB3	2:B:516:GLU:O	2.21	0.40
1:A:381:TYR:CD2	1:A:404:VAL:HG21	2.56	0.40
2:B:481:ASN:C	2:B:483:VAL:N	2.79	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	594/599 (99%)	582 (98%)	11 (2%)	1 (0%)	43	63
2	B	192/217 (88%)	179 (93%)	13 (7%)	0	100	100
All	All	786/816 (96%)	761 (97%)	24 (3%)	1 (0%)	48	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	494	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	523/528 (99%)	522 (100%)	1 (0%)	87	95
2	B	167/190 (88%)	167 (100%)	0	100	100
All	All	690/718 (96%)	689 (100%)	1 (0%)	88	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	607	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	221	GLN
1	A	394	ASN
1	A	508	ASN
1	A	540	HIS
1	A	552	GLN
2	B	440	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	601	2	14,14,15	0.99	2 (14%)	17,19,21	0.83	1 (5%)
3	NAG	A	1004	1	14,14,15	0.97	1 (7%)	17,19,21	0.51	0
3	NAG	A	1002	1	14,14,15	0.58	0	17,19,21	0.51	0
3	NAG	A	1001	1	14,14,15	0.65	0	17,19,21	1.00	1 (5%)
3	NAG	A	1003	1	14,14,15	0.29	0	17,19,21	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	601	2	-	0/6/23/26	0/1/1/1
3	NAG	A	1004	1	-	1/6/23/26	0/1/1/1
3	NAG	A	1002	1	-	3/6/23/26	0/1/1/1
3	NAG	A	1001	1	-	2/6/23/26	0/1/1/1
3	NAG	A	1003	1	-	6/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	NAG	O5-C1	2.91	1.48	1.43
3	A	1004	NAG	O5-C1	-2.90	1.38	1.43
3	B	601	NAG	C1-C2	2.06	1.55	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	NAG	C1-O5-C5	3.07	116.30	112.19
3	B	601	NAG	C1-O5-C5	2.85	116.00	112.19

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1003	NAG	C8-C7-N2-C2
3	A	1003	NAG	O7-C7-N2-C2
3	A	1001	NAG	O5-C5-C6-O6
3	A	1003	NAG	C4-C5-C6-O6
3	A	1002	NAG	O5-C5-C6-O6
3	A	1003	NAG	O5-C5-C6-O6
3	A	1001	NAG	C4-C5-C6-O6
3	A	1004	NAG	O5-C5-C6-O6
3	A	1003	NAG	C1-C2-N2-C7
3	A	1002	NAG	C4-C5-C6-O6
3	A	1002	NAG	C3-C2-N2-C7
3	A	1003	NAG	C3-C2-N2-C7

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1004	NAG	1	0
3	A	1002	NAG	2	0
3	A	1003	NAG	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	596/599 (99%)	-0.30	11 (1%) 67 64	36, 48, 76, 128	0
2	B	194/217 (89%)	0.05	11 (5%) 29 26	40, 53, 92, 100	0
All	All	790/816 (96%)	-0.21	22 (2%) 55 50	36, 49, 83, 128	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	138	PRO	6.9
2	B	480	CYS	4.3
1	A	135	PRO	4.2
1	A	339	VAL	4.1
2	B	382	VAL	4.0
2	B	481	ASN	3.8
1	A	133	CYS	3.7
2	B	478	THR	3.6
2	B	482	GLY	3.6
1	A	140	GLU	3.5
1	A	136	ASP	3.4
1	A	103	ASN	3.2
2	B	527	PRO	3.1
1	A	614	ALA	3.0
2	B	479	PRO	2.9
2	B	335	LEU	2.8
1	A	536	GLU	2.6
2	B	334	ASN	2.6
1	A	139	GLN	2.4
2	B	405	ASP	2.4
1	A	134	ASN	2.3
2	B	488	CYS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	1003	14/15	0.36	0.31	62,72,85,93	0
3	NAG	A	1004	14/15	0.67	0.19	88,95,109,110	0
3	NAG	B	601	14/15	0.77	0.14	63,83,90,94	0
3	NAG	A	1002	14/15	0.83	0.13	77,89,107,108	0
3	NAG	A	1001	14/15	0.86	0.14	50,85,94,95	0

6.5 Other polymers [i](#)

There are no such residues in this entry.