



Full wwPDB EM Validation Report ⓘ

Mar 9, 2026 – 05:19 AM UTC

PDB ID : 7EG6 / pdb_00007eg6
EMDB ID : EMD-31106
Title : Snf5 Finger Helix bound to the nucleosome
Authors : Chen, Z.C.; Chen, K.J.; He, Z.Y.; Ye, Y.P.
Deposited on : 2021-03-24
Resolution : 3.10 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14724 (2.60 - 3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	M	905	 98%
2	A	135	 71% 27%
2	E	135	 69% 30%
3	B	102	 74% 7% 20%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	102	
4	C	129	
4	G	129	
5	D	122	
5	H	122	
6	I	235	
7	J	235	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 12118 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SWI/SNF chromatin-remodeling complex subunit SNF5.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	M	18	Total	C	N	O	0	0
			159	90	40	29		

- Molecule 2 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	A	98	Total	C	N	O	S	0	0
			800	505	153	139	3		
2	E	95	Total	C	N	O	S	0	0
			778	491	148	136	3		

- Molecule 3 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	B	82	Total	C	N	O	S	0	0
			653	413	127	112	1		
3	F	86	Total	C	N	O	S	0	0
			672	424	130	117	1		

- Molecule 4 is a protein called Histone H2A type 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
4	C	107	Total	C	N	O	0	0
			811	510	158	143		
4	G	107	Total	C	N	O	0	0
			815	513	159	143		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	99	ARG	GLY	conflict	UNP P06897
C	123	SER	ALA	conflict	UNP P06897
G	99	ARG	GLY	conflict	UNP P06897

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	123	SER	ALA	conflict	UNP P06897

- Molecule 5 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	D	93	Total	C	N	O	S	0	0
			718	451	128	137	2		
5	H	93	Total	C	N	O	S	0	0
			726	457	130	137	2		

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	29	THR	SER	conflict	UNP P02281
H	29	THR	SER	conflict	UNP P02281

- Molecule 6 is a DNA chain called DNA (235-MER).


Mol	Chain	Residues	Atoms					AltConf	Trace
6	I	146	Total	C	N	O	P	0	0
			2978	1414	542	876	146		

- Molecule 7 is a DNA chain called DNA (235-MER).


Mol	Chain	Residues	Atoms					AltConf	Trace
7	J	146	Total	C	N	O	P	0	0
			3008	1424	562	876	146		




- Molecule 2: Histone H3.2




- Molecule 2: Histone H3.2




- Molecule 3: Histone H4




- Molecule 3: Histone H4



- Molecule 4: Histone H2A type 1



- Molecule 4: Histone H2A type 1



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	525389	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.060	Depositor
Minimum map value	-0.020	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.0077	Depositor
Map size ($\text{\AA}$)	429.68002, 429.68002, 429.68002	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.0742, 1.0742, 1.0742	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	M	0.33	0/158	0.71	0/204
2	A	0.30	0/812	0.47	0/1091
2	E	0.31	0/788	0.49	1/1057 (0.1%)
3	B	0.32	0/660	0.53	0/885
3	F	0.30	0/680	0.48	0/912
4	C	0.29	0/821	0.45	0/1112
4	G	0.30	0/825	0.46	0/1116
5	D	0.28	0/729	0.40	0/985
5	H	0.27	0/737	0.40	0/993
6	I	0.33	0/3337	0.47	0/5144
7	J	0.34	0/3377	0.44	0/5214
All	All	0.32	0/12924	0.46	1/18713 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	117	VAL	N-CA-C	-5.05	108.20	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	M	159	0	166	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	800	0	829	2	0
2	E	778	0	813	1	0
3	B	653	0	695	5	0
3	F	672	0	698	4	0
4	C	811	0	849	2	0
4	G	815	0	860	8	0
5	D	718	0	725	3	0
5	H	726	0	747	9	0
6	I	2978	0	1639	6	0
7	J	3008	0	1639	5	0
All	All	12118	0	9660	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:H:33:SER:OG	5:H:34:TYR:N	2.39	0.54
2:E:63:ARG:HH21	6:I:60:DA:H5''	1.74	0.51
4:G:29:ARG:NH1	5:H:33:SER:O	2.43	0.51
2:A:61:LEU:HD12	3:B:37:LEU:HD23	1.92	0.50
2:A:76:GLN:HE22	3:B:21:VAL:HA	1.76	0.49
6:I:10:DC:H42	7:J:138:DG:H1	1.59	0.49
3:B:44:LYS:HD2	4:G:115:LEU:HB3	1.95	0.49
6:I:144:DC:H2'	6:I:145:DT:H72	1.94	0.48
3:F:59:LYS:NZ	3:F:63:GLU:OE2	2.47	0.47
3:F:92:ARG:HH21	5:H:98:LEU:HA	1.80	0.47
6:I:125:DC:H2''	6:I:126:DG:H5''	1.97	0.46
6:I:48:DT:H2''	6:I:49:DA:C8	2.52	0.45
3:F:22:LEU:HB3	3:F:25:ASN:HD21	1.80	0.45
5:H:83:ARG:NH2	6:I:41:DA:OP2	2.49	0.45
4:G:16:THR:O	4:G:19:SER:OG	2.35	0.44
3:B:47:SER:OG	3:B:48:GLY:N	2.52	0.43
3:F:75:HIS:HB2	5:H:93:THR:HG21	2.01	0.43
1:M:669:ARG:NH1	4:G:92:GLU:OE1	2.52	0.43
4:C:67:GLY:HA3	5:D:46:HIS:CD2	2.54	0.42
5:D:116:THR:O	5:D:120:SER:HB3	2.19	0.42
1:M:665:LYS:NZ	4:G:91:GLU:H	2.18	0.42
4:G:39:TYR:OH	5:H:68:GLU:OE2	2.37	0.42
7:J:68:DG:H2''	7:J:69:DG:C8	2.55	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:H:54:LYS:HE3	5:H:54:LYS:HB3	1.90	0.42
7:J:9:DT:H2"	7:J:10:DA:C8	2.55	0.41
4:G:67:GLY:HA3	5:H:46:HIS:CD2	2.56	0.41
3:B:32:PRO:HA	3:B:35:ARG:HG2	2.03	0.41
7:J:144:DC:H2"	7:J:145:DC:C6	2.55	0.41
4:G:63:LEU:HD13	5:H:42:LEU:HB2	2.03	0.40
4:C:54:VAL:HG21	5:D:95:VAL:HG21	2.04	0.40
7:J:17:DT:H2"	7:J:18:DG:C8	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	M	16/905 (2%)	16 (100%)	0	0	100	100
2	A	96/135 (71%)	96 (100%)	0	0	100	100
2	E	93/135 (69%)	93 (100%)	0	0	100	100
3	B	80/102 (78%)	78 (98%)	2 (2%)	0	100	100
3	F	84/102 (82%)	83 (99%)	1 (1%)	0	100	100
4	C	105/129 (81%)	103 (98%)	2 (2%)	0	100	100
4	G	105/129 (81%)	104 (99%)	1 (1%)	0	100	100
5	D	91/122 (75%)	90 (99%)	1 (1%)	0	100	100
5	H	91/122 (75%)	91 (100%)	0	0	100	100
All	All	761/1881 (40%)	754 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	M	17/823 (2%)	17 (100%)	0	100	100
2	A	84/110 (76%)	84 (100%)	0	100	100
2	E	82/110 (74%)	82 (100%)	0	100	100
3	B	67/78 (86%)	67 (100%)	0	100	100
3	F	67/78 (86%)	67 (100%)	0	100	100
4	C	81/101 (80%)	81 (100%)	0	100	100
4	G	82/101 (81%)	82 (100%)	0	100	100
5	D	77/102 (76%)	77 (100%)	0	100	100
5	H	79/102 (78%)	79 (100%)	0	100	100
All	All	636/1605 (40%)	636 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	39	HIS
2	A	55	GLN
2	A	68	GLN
2	A	76	GLN
2	A	113	HIS
3	B	27	GLN
3	B	75	HIS
5	D	46	HIS
5	D	81	ASN
5	D	92	GLN
2	E	125	GLN
3	F	75	HIS
4	G	24	GLN
4	G	73	ASN
4	G	89	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-31106. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200



Y Index: 200




Z Index: 200

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 171



Y Index: 161




Z Index: 211


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0077. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 144 nm^3 ; this corresponds to an approximate mass of 130 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹


8 Fourier-Shell correlation ⓘ

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-31106 and PDB model 7EG6. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0077 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0077).

9.4 Atom inclusion ⓘ



At the recommended contour level, 96% of all backbone atoms, 95% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0077) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9470	0.5130
A	0.9570	0.5560
B	0.9620	0.5540
C	0.9540	0.5420
D	0.9800	0.5390
E	0.9710	0.5600
F	0.9200	0.5390
G	0.9430	0.5540
H	0.9650	0.5360
I	0.9530	0.4860
J	0.9470	0.4790
M	0.4150	0.3570

1.0

0.0

<0.0