



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:51 PM UTC

PDB ID : 3EHI / pdb_00003ehi
Title : Crystal Structure of Human Thymidylate Synthase M190K with Loop 181-197 stabilized in the inactive conformation
Authors : Lovelace, L.L.; Gibson, L.M.; Johnson, S.R.; Berger, S.H.; Lebioda, L.
Deposited on : 2008-09-12
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	313	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	X	318	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2392 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Thymidylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	X	273	2220	1420	384	403	13	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
X	190	LYS	MET	engineered mutation	UNP P04818

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	X	1	5	4	1	0	0
2	X	1	5	4	1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	X	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	X	158	Total	O	0	0
			158	158		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thymidylate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	96.05Å 96.05Å 80.06Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.59 – 2.00 41.59 – 2.00	Depositor EDS
% Data completeness (in resolution range)	88.4 (41.59-2.00) 88.7 (41.59-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.198 , 0.238 0.195 , 0.235	Depositor DCC
R_{free} test set	1321 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	49.3	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2392	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SO4, CME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	X	1.17	4/2255 (0.2%)	1.12	8/3046 (0.3%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	164	VAL	CA-CB	6.96	1.63	1.54
1	X	181	ALA	N-CA	6.86	1.59	1.46
1	X	223	VAL	C-O	-5.13	1.20	1.24
1	X	80	PHE	CA-C	5.04	1.59	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	187	LEU	CA-C-N	-6.75	111.40	119.84
1	X	187	LEU	C-N-CA	-6.75	111.40	119.84
1	X	262	ILE	N-CA-C	6.04	116.21	110.42
1	X	181	ALA	N-CA-C	5.51	126.43	111.00
1	X	121	LEU	N-CA-C	5.41	122.32	110.80
1	X	190	LYS	N-CA-C	-5.15	105.10	111.33
1	X	269	LEU	N-CA-C	5.06	118.55	112.38
1	X	226	ASN	N-CA-C	5.04	117.67	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	2220	0	2188	57	0
2	X	10	0	0	0	0
3	X	4	0	6	5	0
4	X	158	0	0	8	0
All	All	2392	0	2194	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:229:SER:HB2	4:X:361:HOH:O	1.13	1.31
1:X:229:SER:CB	4:X:361:HOH:O	1.69	1.10
1:X:50:ARG:HH12	1:X:118:LEU:HD12	1.09	1.09
1:X:252:LEU:HD22	3:X:318:EDO:H12	1.40	1.03
1:X:280:ARG:HH21	1:X:280:ARG:HG3	1.33	0.93
1:X:239:HIS:HE1	1:X:285:VAL:H	1.15	0.88
1:X:42:ARG:HD3	4:X:437:HOH:O	1.75	0.85
1:X:50:ARG:NH1	1:X:118:LEU:HD12	1.92	0.85
1:X:147:ARG:NH1	1:X:151:SER:HB3	2.00	0.76
1:X:50:ARG:HH12	1:X:118:LEU:CD1	1.97	0.75
1:X:160:GLN:HB3	1:X:179:MET:HG3	1.67	0.74
1:X:260:ASN:ND2	1:X:310:GLU:HB2	2.03	0.74
1:X:280:ARG:HH21	1:X:280:ARG:CG	2.00	0.74
1:X:278:LYS:HE2	4:X:389:HOH:O	1.88	0.72
1:X:239:HIS:CE1	1:X:285:VAL:H	2.03	0.68
1:X:214:GLN:NE2	4:X:365:HOH:O	2.27	0.67
1:X:180:CME:O	1:X:197:ALA:O	2.16	0.63
1:X:212:LEU:HD23	1:X:250:HIS:CD2	2.36	0.60
1:X:280:ARG:HG3	1:X:299:GLU:OE1	2.02	0.60
1:X:171:ASN:ND2	4:X:457:HOH:O	2.31	0.56
1:X:140:ARG:NH2	1:X:289:ASP:OD1	2.39	0.56
1:X:239:HIS:HE1	1:X:285:VAL:N	1.94	0.55
1:X:160:GLN:CB	1:X:179:MET:HG3	2.38	0.54
1:X:280:ARG:HG3	1:X:280:ARG:NH2	2.12	0.53
1:X:260:ASN:HD22	1:X:310:GLU:HB2	1.74	0.52
1:X:236:MET:HE2	1:X:296:PHE:CZ	2.43	0.52
1:X:121:LEU:HD12	1:X:121:LEU:H	1.75	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:180:CME:O	1:X:181:ALA:O	2.29	0.51
1:X:97:ASN:HD21	1:X:99:LYS:HG3	1.75	0.51
1:X:180:CME:O	1:X:181:ALA:C	2.54	0.50
1:X:184:PRO:HA	1:X:187:LEU:HD12	1.93	0.50
1:X:306:THR:HG21	1:X:308:LYS:NZ	2.26	0.50
1:X:187:LEU:HD22	1:X:189:LEU:HD12	1.95	0.48
1:X:187:LEU:O	1:X:188:PRO:C	2.56	0.48
1:X:280:ARG:NH2	1:X:299:GLU:OE1	2.46	0.48
1:X:189:LEU:HA	1:X:193:PRO:HD3	1.95	0.47
1:X:179:MET:HB3	1:X:179:MET:HE3	1.58	0.47
1:X:186:ASP:C	1:X:188:PRO:HD2	2.41	0.45
1:X:179:MET:HB2	1:X:179:MET:HE2	1.65	0.45
1:X:244:LYS:HB2	1:X:244:LYS:HE3	1.70	0.45
1:X:280:ARG:CG	1:X:280:ARG:NH2	2.69	0.45
1:X:165:ILE:O	1:X:169:LYS:HG3	2.17	0.44
1:X:134:VAL:HG11	1:X:184:PRO:HB3	2.00	0.44
1:X:250:HIS:ND1	3:X:318:EDO:H11	2.33	0.44
1:X:116:ASP:O	1:X:118:LEU:N	2.51	0.43
1:X:252:LEU:CD2	3:X:318:EDO:H12	2.30	0.43
1:X:229:SER:OG	4:X:361:HOH:O	2.08	0.43
1:X:133:PRO:HG3	4:X:400:HOH:O	2.18	0.43
1:X:123:PHE:CG	1:X:124:SER:N	2.87	0.42
1:X:98:ALA:HB2	1:X:131:LEU:HD21	2.02	0.42
1:X:294:GLU:H	1:X:294:GLU:CD	2.28	0.42
1:X:252:LEU:HD22	3:X:318:EDO:C1	2.30	0.41
1:X:250:HIS:CG	3:X:318:EDO:H11	2.54	0.41
1:X:304:HIS:HB3	1:X:305:PRO:CD	2.51	0.41
1:X:190:LYS:C	1:X:192:LEU:H	2.28	0.41
1:X:73:LEU:HD21	1:X:81:TRP:HB2	2.03	0.41
1:X:34:LEU:HD11	1:X:76:THR:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	X	267/313 (85%)	248 (93%)	13 (5%)	6 (2%)	5	2

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	117	PHE
1	X	181	ALA
1	X	119	ASP
1	X	121	LEU
1	X	187	LEU
1	X	191	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	X	238/269 (88%)	221 (93%)	17 (7%)	13	10

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	82	LYS
1	X	126	ARG
1	X	140	ARG
1	X	164	VAL
1	X	198	LEU
1	X	221	LEU
1	X	244	LYS
1	X	252	LEU
1	X	259	LEU
1	X	272	GLU
1	X	274	ARG
1	X	280	ARG
1	X	284	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	287	LYS
1	X	292	LYS
1	X	299	GLU
1	X	308	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	97	ASN
1	X	138	GLN
1	X	171	ASN
1	X	205	ASN
1	X	226	ASN
1	X	239	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CME	X	199	1	8,9,10	1.08	1 (12%)	6,9,11	1.91	2 (33%)
1	CME	X	180	1	8,9,10	0.78	0	6,9,11	1.86	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	X	199	1	-	0/5/8/10	-
1	CME	X	180	1	-	1/5/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	199	CME	CB-SG	-2.25	1.74	1.81

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	180	CME	CB-CA-C	-4.09	99.71	110.80
1	X	199	CME	CB-CA-C	-3.58	101.09	110.80
1	X	199	CME	CE-SD-SG	2.39	113.96	103.46

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	X	180	CME	CZ-CE-SD-SG

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	X	180	CME	3	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	X	317	-	4,4,4	0.56	0	6,6,6	0.60	0
3	EDO	X	318	-	3,3,3	0.72	0	2,2,2	0.19	0
2	SO4	X	316	-	4,4,4	0.21	0	6,6,6	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	X	318	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	X	318	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	X	318	EDO	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	271/313 (86%)	0.66	21 (7%) 19 18	38, 51, 87, 110	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	120	SER	5.9
1	X	121	LEU	5.4
1	X	117	PHE	4.7
1	X	118	LEU	4.5
1	X	189	LEU	4.4
1	X	192	LEU	3.4
1	X	123	PHE	3.3
1	X	101	LEU	3.2
1	X	119	ASP	2.8
1	X	187	LEU	2.8
1	X	188	PRO	2.7
1	X	131	LEU	2.6
1	X	122	GLY	2.6
1	X	125	THR	2.5
1	X	142	PHE	2.4
1	X	103	SER	2.4
1	X	310	GLU	2.2
1	X	181	ALA	2.2
1	X	126	ARG	2.2
1	X	191	ALA	2.2
1	X	26	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	X	199	10/11	0.91	0.13	40,44,56,59	0
1	CME	X	180	10/11	0.93	0.13	50,55,69,69	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	X	318	4/4	0.88	0.36	48,52,56,58	0
2	SO4	X	316	5/5	0.93	0.09	67,68,68,68	0
2	SO4	X	317	5/5	0.94	0.08	57,61,65,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.