



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:33 PM UTC

PDB ID : 1EI5 / pdb_00001ei5
Title : CRYSTAL STRUCTURE OF A D-AMINOPEPTIDASE FROM
OCHROBACTRUM ANTHROPI
Authors : Bompard-Gilles, C.; Remaut, H.; Villeret, V.; Prange, T.; Fanuel, L.; Joris, J.;
Frere, J.-M.; Van Beeumen, J.
Deposited on : 2000-02-24
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	520	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4329 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-AMINOPEPTIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	518	Total	C	N	O	S	0	0	0
			3961	2493	697	748	23			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	368	Total	O	0	0
			368	368		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: D-AMINOPEPTIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	82.86Å 82.86Å 204.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	12.50 – 1.90 12.50 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (12.50-1.90) 96.3 (12.50-1.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.20 (at 1.90Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.163 , 0.192 0.151 , 0.179	Depositor DCC
R_{free} test set	2776 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtriage
Anisotropy	0.262	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4329	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	2/4046 (0.0%)	1.53	44/5484 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	519	GLN	C-O	-7.63	1.17	1.24
1	A	520	PRO	N-CD	6.42	1.56	1.47

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	482	ALA	CA-C-O	-14.07	106.17	120.66
1	A	335	SER	N-CA-C	11.19	127.18	109.50
1	A	520	PRO	CA-N-CD	-11.10	96.46	112.00
1	A	355	THR	N-CA-CB	-10.92	94.48	110.53
1	A	335	SER	CA-C-O	10.19	133.03	121.11
1	A	261	THR	N-CA-CB	-9.99	94.31	111.69
1	A	169	THR	N-CA-CB	-9.27	96.42	110.33
1	A	308	THR	N-CA-CB	-9.26	94.56	111.13
1	A	196	THR	N-CA-CB	-9.17	94.73	110.32
1	A	203	THR	CA-CB-OG1	8.42	122.23	109.60
1	A	335	SER	CA-C-N	8.32	137.42	121.54
1	A	335	SER	C-N-CA	8.32	137.42	121.54
1	A	172	THR	N-CA-CB	-7.86	98.35	110.29
1	A	68	THR	CA-CB-OG1	7.47	120.80	109.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	495	ASP	CB-CA-C	-7.44	95.61	110.42
1	A	375	THR	CA-CB-OG1	7.29	120.53	109.60
1	A	483	PRO	N-CA-CB	7.21	110.53	102.60
1	A	393	THR	CA-CB-OG1	7.10	120.26	109.60
1	A	217	THR	N-CA-CB	7.09	122.60	110.68
1	A	196	THR	CA-CB-OG1	6.51	119.36	109.60
1	A	257	SER	N-CA-C	6.41	120.41	112.59
1	A	308	THR	OG1-CB-CG2	6.37	122.03	109.30
1	A	158	PHE	CA-CB-CG	-6.37	107.43	113.80
1	A	312	PHE	CA-CB-CG	6.35	120.15	113.80
1	A	298	ARG	NE-CZ-NH2	6.32	124.89	119.20
1	A	261	THR	OG1-CB-CG2	6.23	121.76	109.30
1	A	468	GLY	CA-C-O	-6.11	115.83	121.63
1	A	254	ARG	NE-CZ-NH2	6.03	124.63	119.20
1	A	517	ARG	NE-CZ-NH1	-5.83	115.67	121.50
1	A	308	THR	CB-CA-C	5.79	121.13	109.38
1	A	168	HIS	CA-C-O	5.63	126.39	120.42
1	A	519	GLN	CA-C-O	5.63	126.62	120.94
1	A	346	TRP	CA-CB-CG	5.56	124.17	113.60
1	A	188	LYS	N-CA-C	5.50	118.04	111.71
1	A	355	THR	CA-CB-CG2	5.44	119.75	110.50
1	A	123	ASP	CA-CB-CG	5.44	118.04	112.60
1	A	308	THR	CA-CB-OG1	5.43	117.75	109.60
1	A	359	LEU	CA-C-O	-5.32	114.58	120.38
1	A	254	ARG	NE-CZ-NH1	-5.28	116.22	121.50
1	A	189	ARG	CD-NE-CZ	-5.23	117.08	124.40
1	A	396	ARG	CD-NE-CZ	5.19	131.67	124.40
1	A	365	GLY	CA-C-N	5.15	131.38	121.54
1	A	365	GLY	C-N-CA	5.15	131.38	121.54
1	A	203	THR	N-CA-CB	5.02	118.77	110.69

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	482	ALA	Peptide,Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3961	0	3801	64	0
2	A	368	0	0	11	0
All	All	4329	0	3801	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:429:ARG:HB3	1:A:519:GLN:HB2	1.47	0.95
1:A:162:ALA:HB1	1:A:223:MET:HE2	1.57	0.87
1:A:129:LEU:H	1:A:132:GLN:HE21	1.22	0.83
1:A:352:ASP:OD1	1:A:355:THR:HB	1.82	0.78
1:A:172:THR:HG22	1:A:175:ASP:H	1.50	0.76
1:A:68:THR:HG22	2:A:611:HOH:O	1.85	0.76
1:A:378:GLU:OE2	1:A:393:THR:HG23	1.85	0.75
1:A:355:THR:HG23	1:A:357:LEU:HG	1.70	0.74
1:A:68:THR:HG21	1:A:224:GLY:HA2	1.72	0.71
1:A:429:ARG:CB	1:A:519:GLN:HB2	2.20	0.69
1:A:519:GLN:NE2	1:A:520:PRO:HD2	2.10	0.66
1:A:169:THR:HG21	1:A:176:ILE:HD11	1.78	0.65
1:A:45:ASP:OD2	1:A:203:THR:HG21	1.97	0.64
1:A:129:LEU:H	1:A:132:GLN:NE2	1.95	0.64
1:A:495:ASP:HB3	1:A:497:LYS:H	1.63	0.63
1:A:287:HIS:HD2	1:A:288:GLY:O	1.83	0.62
1:A:141:LYS:HE2	2:A:654:HOH:O	1.99	0.62
1:A:107:GLN:HE21	1:A:148:GLY:HA2	1.66	0.61
1:A:261:THR:HG23	1:A:262:PHE:O	2.00	0.61
1:A:393:THR:HG21	1:A:406:ARG:HE	1.65	0.61
1:A:393:THR:CG2	1:A:406:ARG:HE	2.13	0.60
1:A:428:GLY:HA2	1:A:520:PRO:HB2	1.83	0.60
1:A:68:THR:HG21	1:A:224:GLY:O	2.04	0.58
1:A:165:ILE:O	1:A:169:THR:HB	2.04	0.58
1:A:195:ASP:HB3	1:A:198:LEU:HD22	1.85	0.58
1:A:500:ILE:HB	1:A:517:ARG:NH2	2.18	0.57
1:A:172:THR:CG2	1:A:175:ASP:H	2.17	0.56
1:A:183:ALA:HB3	1:A:184:PRO:HD3	1.90	0.54
1:A:418:VAL:HG12	1:A:468:GLY:HA3	1.89	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:519:GLN:O	1:A:520:PRO:C	2.52	0.53
1:A:375:THR:HG21	2:A:645:HOH:O	2.08	0.53
1:A:169:THR:HG23	1:A:171:ARG:HB2	1.91	0.52
1:A:10:GLU:OE2	1:A:37:LEU:HD22	2.11	0.50
1:A:355:THR:HG23	1:A:357:LEU:CG	2.39	0.50
1:A:308:THR:HG21	1:A:322:ALA:O	2.11	0.49
1:A:169:THR:CG2	1:A:171:ARG:H	2.25	0.49
1:A:203:THR:HG22	2:A:602:HOH:O	2.12	0.49
1:A:300:HIS:HD2	2:A:543:HOH:O	1.96	0.48
1:A:246:ARG:O	1:A:254:ARG:NH2	2.47	0.48
1:A:517:ARG:NE	1:A:520:PRO:O	2.46	0.47
1:A:196:THR:HB	1:A:220:ILE:O	2.13	0.47
1:A:393:THR:HB	1:A:406:ARG:HG2	1.97	0.46
1:A:300:HIS:HE1	2:A:716:HOH:O	1.98	0.46
1:A:169:THR:HG22	1:A:171:ARG:H	1.81	0.46
1:A:222:TRP:CD2	1:A:226:ALA:HB3	2.51	0.45
1:A:424:HIS:HD2	2:A:603:HOH:O	1.98	0.45
1:A:217:THR:HG22	2:A:805:HOH:O	2.16	0.45
1:A:403:GLU:CD	1:A:412:ARG:HH11	2.24	0.45
1:A:355:THR:CG2	1:A:357:LEU:H	2.30	0.45
1:A:375:THR:CG2	2:A:645:HOH:O	2.64	0.45
1:A:257:SER:HB2	1:A:285:THR:HG21	1.99	0.44
1:A:428:GLY:CA	1:A:520:PRO:HB2	2.45	0.44
1:A:68:THR:HG21	1:A:224:GLY:CA	2.45	0.44
1:A:189:ARG:HH11	1:A:189:ARG:HD3	1.51	0.43
1:A:270:TYR:OH	1:A:287:HIS:HE1	2.02	0.43
1:A:364:ALA:CB	1:A:379:MET:HE1	2.48	0.43
1:A:519:GLN:HB3	1:A:520:PRO:HD2	2.01	0.43
1:A:519:GLN:HB3	1:A:520:PRO:CD	2.49	0.42
1:A:172:THR:HG23	1:A:174:VAL:HB	2.02	0.41
1:A:18:GLN:HG2	2:A:772:HOH:O	2.20	0.41
1:A:392:VAL:HG12	1:A:393:THR:HG22	2.01	0.41
1:A:292:ARG:HG3	2:A:593:HOH:O	2.20	0.41
1:A:65:LYS:HZ3	1:A:155:ASN:ND2	2.19	0.40
1:A:45:ASP:OD2	1:A:203:THR:CG2	2.65	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	516/520 (99%)	503 (98%)	10 (2%)	3 (1%)	21	13

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	366	HIS
1	A	483	PRO
1	A	336	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	401/422 (95%)	379 (94%)	22 (6%)	19	11

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	LEU
1	A	68	THR
1	A	81	LEU
1	A	140	LEU
1	A	164	LEU
1	A	169	THR
1	A	172	THR
1	A	196	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	198	LEU
1	A	203	THR
1	A	217	THR
1	A	261	THR
1	A	263	LYS
1	A	276	LEU
1	A	306	LEU
1	A	308	THR
1	A	340	VAL
1	A	355	THR
1	A	375	THR
1	A	393	THR
1	A	423	LYS
1	A	503	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	107	GLN
1	A	132	GLN
1	A	155	ASN
1	A	275	ASN
1	A	287	HIS
1	A	297	GLN
1	A	300	HIS
1	A	424	HIS
1	A	519	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	518/520 (99%)	-0.50	10 (1%) 66 70	12, 19, 36, 60	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	520	PRO	8.8
1	A	519	GLN	7.3
1	A	495	ASP	3.4
1	A	335	SER	3.3
1	A	337	VAL	3.2
1	A	334	SER	3.1
1	A	496	ASP	2.6
1	A	3	LYS	2.4
1	A	518	VAL	2.4
1	A	366	HIS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.