



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 11:54 AM UTC

PDB ID : 5EI5 / pdb_00005ei5
Title : Crystal structure of MSF-aged Torpedo californica Acetylcholinesterase in complex with alkylene-linked bis-tacrine dimer (7 carbon linker)
Authors : Pesaresi, A.; Lamba, D.
Deposited on : 2015-10-29
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	
2	B	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	03S	A	603	-	X	X	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 4724 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholinesterase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	534	Total	C	N	O	S	0	0	0
			4263	2731	724	786	22			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is methanesulfonic acid (CCD ID: 03S) (formula: $\text{CH}_4\text{O}_3\text{S}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	1	2	1		

- Molecule 5 is N,N'-DI-1,2,3,4-TETRAHYDROACRIDIN-9-YLHEPTANE-1,7-DIAMINE (CCD ID: AA7) (formula: $\text{C}_{33}\text{H}_{40}\text{N}_4$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	C	N	0
			37	33	4	


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	364	Total	O	0	0
			364	364		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetylcholinesterase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	111.28Å 111.28Å 136.74Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.23 – 2.10 48.23 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.2 (48.23-2.10) 99.2 (48.23-2.10)	Depositor EDS
R_{merge}	0.22	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.13 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.173 , 0.206 (Not available) , 0.243	Depositor DCC
R_{free} test set	2898 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	0.108	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4724	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, 03S, AA7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.35	12/4387 (0.3%)	1.17	13/5955 (0.2%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	58	TRP	C-N	-12.66	1.15	1.33
1	A	59	ASN	C-N	-9.78	1.20	1.33
1	A	355	GLY	C-O	-6.09	1.16	1.23
1	A	380	ASP	C-N	-5.85	1.24	1.33
1	A	63	TYR	C-O	-5.61	1.17	1.24
1	A	518	VAL	C-O	-5.51	1.17	1.24
1	A	212	SER	CA-C	5.46	1.59	1.52
1	A	172	ASP	C-O	5.42	1.30	1.24
1	A	112	MET	N-CA	5.37	1.52	1.46
1	A	243	ARG	C-O	-5.23	1.18	1.24
1	A	392	ASP	CA-C	5.11	1.59	1.52
1	A	22	VAL	C-O	-5.09	1.18	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4	SER	N-CA-C	6.82	119.30	109.14
1	A	129	VAL	CB-CA-C	-6.61	96.77	111.77
1	A	492	TRP	CA-C-N	-6.46	113.98	120.31
1	A	492	TRP	C-N-CA	-6.46	113.98	120.31
1	A	291	SER	N-CA-C	6.37	118.02	111.14
1	A	381	ASP	N-CA-C	-6.33	105.55	113.28
1	A	173	GLN	N-CA-C	-6.13	104.68	111.36
1	A	381	ASP	N-CA-CB	5.39	118.92	110.46
1	A	341	LYS	N-CA-C	5.33	117.17	111.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	282	LEU	CA-C-N	-5.13	114.45	119.64
1	A	282	LEU	C-N-CA	-5.13	114.45	119.64
1	A	517	ARG	CA-CB-CG	-5.13	103.83	114.10
1	A	380	ASP	O-C-N	-5.03	115.90	122.59

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4263	0	4103	33	0
2	B	28	0	25	1	0
3	A	28	0	26	0	0
4	A	4	0	3	2	0
5	A	37	0	40	17	0
6	A	364	0	0	2	0
All	All	4724	0	4197	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:TYR:HB2	5:A:604:AA7:H5C2	1.26	1.09
1:A:70:TYR:CB	5:A:604:AA7:H5C2	2.06	0.83
5:A:604:AA7:C28	5:A:604:AA7:H191	2.13	0.78
1:A:279:TRP:CE3	5:A:604:AA7:H4C1	2.24	0.72
1:A:279:TRP:CZ3	5:A:604:AA7:H4C1	2.27	0.69
1:A:119:GLY:HA2	4:A:603:03S:H213	1.75	0.68
1:A:374:GLN:HE21	1:A:520:MET:HE1	1.61	0.66
5:A:604:AA7:H191	5:A:604:AA7:H281	1.77	0.64
5:A:604:AA7:H14	5:A:604:AA7:H251	1.79	0.64
5:A:604:AA7:H191	5:A:604:AA7:H282	1.80	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330:PHE:CZ	5:A:604:AA7:H201	2.37	0.60
1:A:223:ILE:HA	1:A:320:LEU:O	2.06	0.55
5:A:604:AA7:C28	5:A:604:AA7:C19	2.85	0.54
1:A:468:ARG:HD3	6:A:711:HOH:O	2.07	0.53
1:A:374:GLN:NE2	1:A:520:MET:HE1	2.22	0.53
1:A:374:GLN:HE21	1:A:520:MET:CE	2.21	0.52
5:A:604:AA7:H192	6:A:746:HOH:O	2.08	0.52
1:A:23:LEU:O	1:A:24:SER:HB2	2.10	0.51
1:A:146:LEU:C	1:A:146:LEU:HD12	2.36	0.51
1:A:233:TRP:CZ3	4:A:603:03S:H113	2.45	0.51
1:A:321:LEU:HD11	1:A:408:VAL:CG2	2.40	0.51
1:A:451:PRO:HA	1:A:458:TYR:CD1	2.45	0.51
1:A:279:TRP:HB2	5:A:604:AA7:N7	2.27	0.49
1:A:330:PHE:HZ	5:A:604:AA7:H201	1.74	0.49
5:A:604:AA7:H14	5:A:604:AA7:C25	2.42	0.49
1:A:484:GLU:HB3	1:A:487:SER:HB3	1.94	0.49
1:A:531:LEU:C	1:A:531:LEU:HD23	2.38	0.48
2:B:1:NAG:H61	2:B:2:NAG:C1	2.44	0.47
1:A:316:LYS:HE3	1:A:414:PHE:HB3	1.95	0.46
1:A:320:LEU:C	1:A:320:LEU:HD23	2.40	0.46
1:A:455:GLU:CD	1:A:455:GLU:H	2.23	0.46
1:A:419:TYR:CZ	1:A:494:LEU:HD13	2.51	0.46
1:A:23:LEU:O	1:A:24:SER:CB	2.64	0.45
1:A:278:GLU:OE1	5:A:604:AA7:H5C1	2.16	0.45
1:A:484:GLU:OE1	1:A:485:PRO:HD2	2.18	0.44
1:A:330:PHE:CE1	5:A:604:AA7:H281	2.54	0.43
1:A:495:PHE:C	1:A:495:PHE:CD1	2.96	0.43
1:A:456:LEU:O	1:A:457:ASN:HB2	2.18	0.42
1:A:330:PHE:CD1	5:A:604:AA7:H301	2.55	0.41
1:A:143:LEU:HD13	1:A:143:LEU:C	2.46	0.41
1:A:321:LEU:HD11	1:A:408:VAL:HG23	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	532/534 (100%)	508 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	467/467 (100%)	450 (96%)	17 (4%)	31	34

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASP
1	A	15	VAL
1	A	46	ARG
1	A	129	VAL
1	A	161	SER
1	A	221	ARG
1	A	228	SER
1	A	279	TRP
1	A	294	PRO
1	A	295	VAL
1	A	299	GLU
1	A	357	LYS
1	A	366	LEU
1	A	391	LEU
1	A	461	GLU
1	A	490	SER
1	A	505	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	65	ASN
1	A	251	ASN
1	A	272	GLN
1	A	280	ASN
1	A	374	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	602	1	14,14,15	0.78	0	17,19,21	2.05	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	602	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	NAG	C1-O5-C5	4.04	117.59	112.19
3	A	602	NAG	C1-C2-N2	-3.45	104.99	110.43
3	A	602	NAG	O5-C5-C6	3.26	114.00	107.66
3	A	602	NAG	O5-C1-C2	-3.12	106.47	111.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	NAG	C3-C4-C5	-3.06	104.69	110.23
3	A	602	NAG	C4-C3-C2	2.12	114.12	111.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	0.56	0	17,19,21	1.44	3 (17%)
2	NAG	B	2	2	14,14,15	0.29	0	17,19,21	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	O5-C5-C6	3.04	113.59	107.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	O6-C6-C5	2.36	119.38	111.33
2	B	1	NAG	C1-O5-C5	-2.23	109.19	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	NAG	O5-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	NAG	1	0
2	B	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	601	1	14,14,15	0.29	0	17,19,21	0.56	0
3	NAG	A	602	1	14,14,15	0.78	0	17,19,21	2.05	6 (35%)
4	03S	A	603	1	1,3,4	3.83	1 (100%)	2,3,6	4.04	2 (100%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	AA7	A	604	-	42,42,42	2.83	10 (23%)	51,57,57	1.78	14 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	601	1	-	1/6/23/26	0/1/1/1
3	NAG	A	602	1	-	0/6/23/26	0/1/1/1
5	AA7	A	604	-	-	7/12/26/26	0/6/6/6

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	604	AA7	C31-C26	-8.73	1.37	1.50
5	A	604	AA7	C28-C27	-8.39	1.38	1.51
5	A	604	AA7	C6-C1	-7.42	1.39	1.50
5	A	604	AA7	C3-C2	-7.31	1.40	1.51
5	A	604	AA7	C33-N32	-4.18	1.31	1.37
4	A	603	03S	C13-S10	-3.83	1.60	1.78
5	A	604	AA7	C8-N7	-3.11	1.32	1.37
5	A	604	AA7	C35-C34	-2.80	1.38	1.43
5	A	604	AA7	C34-C33	-2.72	1.38	1.42
5	A	604	AA7	C14-C9	-2.36	1.37	1.42
5	A	604	AA7	C10-C9	-2.07	1.39	1.43

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	604	AA7	C26-N32-C33	5.22	123.86	117.67
4	A	603	03S	O12-S10-C13	4.93	114.23	105.48
5	A	604	AA7	C1-N7-C8	4.09	122.51	117.67
3	A	602	NAG	C1-O5-C5	4.04	117.59	112.19
5	A	604	AA7	C27-C26-N32	-4.00	120.83	123.66
3	A	602	NAG	C1-C2-N2	-3.45	104.99	110.43
5	A	604	AA7	C34-C33-N32	-3.31	119.30	122.80
3	A	602	NAG	O5-C5-C6	3.26	114.00	107.66
3	A	602	NAG	O5-C1-C2	-3.12	106.47	111.29
3	A	602	NAG	C3-C4-C5	-3.06	104.69	110.23
4	A	603	03S	O11-S10-C13	2.91	110.64	105.48
5	A	604	AA7	C35-C27-C26	-2.81	116.96	118.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	604	AA7	C2-C1-N7	-2.72	121.74	123.66
5	A	604	AA7	C29-C28-C27	2.68	118.29	112.84
5	A	604	AA7	C30-C31-C26	2.65	117.60	113.48
5	A	604	AA7	C5-C6-C1	2.58	117.50	113.48
5	A	604	AA7	C17-C8-C9	2.36	121.68	119.13
5	A	604	AA7	C9-C8-N7	-2.35	120.31	122.80
3	A	602	NAG	C4-C3-C2	2.12	114.12	111.02
5	A	604	AA7	C10-C2-C1	-2.12	117.38	118.68
5	A	604	AA7	C15-C14-C9	-2.05	118.14	120.91
5	A	604	AA7	C31-C26-C27	2.02	123.46	121.40

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	604	AA7	C23-C24-C25-N11
3	A	601	NAG	O5-C5-C6-O6
5	A	604	AA7	N36-C19-C20-C21
5	A	604	AA7	C19-C20-C21-C22
5	A	604	AA7	C21-C22-C23-C24
5	A	604	AA7	C20-C21-C22-C23
5	A	604	AA7	C27-C35-N36-C19
5	A	604	AA7	C9-C10-N11-C25


There are no ring outliers.

2 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	603	03S	2	0
5	A	604	AA7	17	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	59:ASN	C	60:ALA	N	1.20
1	A	58:TRP	C	59:ASN	N	1.15

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	534/534 (100%)	-0.10	9 (1%) 69 71	14, 25, 45, 75	2 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	HIS	26.1
1	A	2	ASP	15.7
1	A	486	HIS	4.5
1	A	485	PRO	4.5
1	A	535	THR	3.7
1	A	279	TRP	2.6
1	A	488	GLN	2.4
1	A	454	LYS	2.2
1	A	268	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	602	14/15	0.84	0.14	57,60,67,76	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	B	2	14/15	0.71	0.21	54,57,68,76	0
2	NAG	B	1	14/15	0.90	0.11	31,35,41,44	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	601	14/15	0.55	0.27	58,68,74,87	0
5	AA7	A	604	37/37	0.78	0.14	20,20,20,20	0
3	NAG	A	602	14/15	0.84	0.14	57,60,67,76	0
4	03S	A	603	4/5	0.98	0.06	25,30,31,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.