



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:16 PM UTC

PDB ID : 5EIB / pdb_00005eib
Title : Crystal structure of CPAP PN2-3 C-terminal loop-helix in complex with DARPin-tubulin
Authors : Li, H.; Zheng, X.
Deposited on : 2015-10-29
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	451	
2	D	445	
3	E	169	
4	F	23	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8684 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	429	Total	C	N	O	S	0	0	0
			3364	2132	572	639	21			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	430	Total	C	N	O	S	0	1	0
			3375	2116	579	655	25			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	203	SER	CYS	see sequence details	UNP Q6B856


- Molecule 3 is a protein called Designed ankyrin repeat protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	156	Total	C	N	O	S	0	0	0
			1159	730	199	227	3			

- Molecule 4 is a protein called Peptide from Centromere protein J.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	F	15	Total	C	N	O	0	0	0
			123	80	24	19			

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	D	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	281	Total	O	0	0
			281	281		
7	D	183	Total	O	0	0
			183	183		
7	E	125	Total	O	0	0
			125	125		
7	F	8	Total	O	0	0
			8	8		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain



- Molecule 2: Tubulin beta-2B chain



- Molecule 3: Designed ankyrin repeat protein



- Molecule 4: Peptide from Centromere protein J

Chain F:

57%9%35%

K372	L376	R377	R378	T386	ASN	ALA	LYS	SER	LYS	PHE	GLN	LYS
------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.86Å 91.12Å 83.31Å 90.00° 96.97° 90.00°	Depositor
Resolution (Å)	41.35 – 2.10 41.35 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.9 (41.35-2.10) 99.1 (41.35-2.10)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.78 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.178 , 0.222 0.178 , 0.222	Depositor DCC
R_{free} test set	3131 reflections (4.55%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.310	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8684	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.61	2/3441 (0.1%)	0.86	6/4672 (0.1%)
2	D	0.70	5/3452 (0.1%)	0.96	18/4677 (0.4%)
3	E	0.53	0/1175	0.79	0/1597
4	F	0.40	0/125	0.77	0/164
All	All	0.64	7/8193 (0.1%)	0.89	24/11110 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	9

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	110	ILE	C-O	-7.87	1.15	1.24
1	C	109	THR	C-O	7.32	1.33	1.24
2	D	107	HIS	C-O	7.12	1.31	1.23
2	D	111	GLY	CA-C	-6.70	1.44	1.52
2	D	112	ALA	C-O	-6.29	1.16	1.24
2	D	111	GLY	C-O	-5.47	1.17	1.23
2	D	104	ALA	C-O	-5.04	1.18	1.24

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	107	HIS	N-CA-C	-10.57	94.65	111.04
2	D	110	GLU	N-CA-C	-10.03	101.17	113.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	262	TYR	CA-C-N	8.80	129.67	119.47
1	C	262	TYR	C-N-CA	8.80	129.67	119.47
2	D	106	GLY	CA-C-O	-8.65	111.66	121.00
2	D	114	LEU	N-CA-C	7.94	121.03	111.82
2	D	108	TYR	N-CA-C	-7.91	101.75	111.40
2	D	109	THR	N-CA-C	-7.88	99.50	111.56
2	D	111	GLY	N-CA-C	-7.27	103.46	112.77
2	D	283	TYR	N-CA-C	-7.14	98.18	108.60
1	C	113	GLU	N-CA-C	6.55	119.24	111.71
2	D	109	THR	CA-C-N	-6.37	110.69	122.09
2	D	109	THR	C-N-CA	-6.37	110.69	122.09
2	D	288	VAL	O-C-N	6.32	124.55	120.07
2	D	106	GLY	N-CA-C	-6.00	105.54	112.50
1	C	36	MET	CA-C-N	5.85	127.15	119.84
1	C	36	MET	C-N-CA	5.85	127.15	119.84
2	D	108	TYR	CA-C-N	-5.67	111.44	121.81
2	D	108	TYR	C-N-CA	-5.67	111.44	121.81
2	D	170	SER	CA-C-N	-5.28	116.27	122.93
2	D	170	SER	C-N-CA	-5.28	116.27	122.93
2	D	98	GLY	N-CA-C	5.17	118.08	112.29
1	C	178	SER	N-CA-C	5.03	117.60	110.50
2	D	112	ALA	N-CA-C	-5.01	106.00	111.82

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	106	GLY	Mainchain
2	D	108	TYR	Mainchain,Peptide
2	D	279	GLY	Peptide
2	D	280	SER	Peptide
2	D	281	GLN	Peptide
2	D	282	GLN	Peptide
2	D	283	TYR	Peptide
2	D	98	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3364	0	3274	44	0
2	D	3375	0	3247	54	0
3	E	1159	0	1158	18	0
4	F	123	0	130	2	0
5	C	32	0	12	0	0
5	D	32	0	12	1	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	C	281	0	0	10	0
7	D	183	0	0	11	0
7	E	125	0	0	10	0
7	F	8	0	0	0	0
All	All	8684	0	7833	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (113) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:139:VAL:CG2	7:E:284:HOH:O	1.78	1.29
1:C:297:GLU:OE2	1:C:339:ARG:NH2	1.74	1.21
3:E:139:VAL:HG21	7:E:284:HOH:O	1.34	1.19
2:D:308:ARG:NH2	7:D:601:HOH:O	1.82	1.10
2:D:88:ARG:NH1	7:D:603:HOH:O	1.94	0.99
2:D:372:LYS:NZ	7:D:605:HOH:O	1.95	0.97
3:E:29:GLU:OE1	7:E:201:HOH:O	1.84	0.95
2:D:278:ARG:NH1	7:D:606:HOH:O	1.98	0.94
2:D:167[B]:ASN:ND2	2:D:252:LEU:HD22	1.85	0.92
2:D:437:ASP:OD2	7:D:604:HOH:O	1.94	0.84
1:C:263:PRO:O	7:C:601:HOH:O	1.94	0.84
1:C:340:SER:O	7:C:602:HOH:O	1.94	0.83
1:C:229:ARG:HD3	7:C:604:HOH:O	1.79	0.80
1:C:220:GLU:HG3	2:D:326:LYS:HD3	1.62	0.79
2:D:15:GLN:HG3	7:D:657:HOH:O	1.83	0.79
1:C:2:ARG:HH22	1:C:50:ASN:HB3	1.51	0.76
2:D:167[B]:ASN:HD22	2:D:252:LEU:HD22	1.49	0.75
1:C:270:ALA:HB3	1:C:302:MET:HE2	1.71	0.70
3:E:23:ARG:NH2	7:E:202:HOH:O	2.12	0.69
2:D:39:ASP:O	7:D:607:HOH:O	2.10	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:71:GLU:HB2	1:C:98:ASP:HB3	1.76	0.67
2:D:308:ARG:HG2	2:D:308:ARG:HH11	1.60	0.67
1:C:336:LYS:C	7:C:618:HOH:O	2.40	0.63
1:C:334:THR:HG23	1:C:338:LYS:HE3	1.81	0.63
3:E:134:LYS:NZ	7:E:204:HOH:O	2.31	0.62
2:D:70:LEU:HB3	2:D:98:GLY:HA2	1.83	0.61
2:D:215:ARG:NH2	3:E:160:ASP:OD1	2.30	0.61
2:D:278:ARG:NH1	2:D:278:ARG:HG2	2.15	0.60
1:C:334:THR:CG2	1:C:338:LYS:HE3	2.31	0.60
2:D:400:ARG:HD2	3:E:112:TRP:CE2	2.38	0.59
3:E:126:LEU:O	3:E:130:GLU:HG3	2.02	0.58
1:C:108:TYR:HA	1:C:112:LYS:HE2	1.85	0.58
1:C:161:TYR:HB3	1:C:164:LYS:HG2	1.85	0.58
2:D:123:ARG:O	2:D:127:GLU:HG3	2.04	0.57
1:C:265:ILE:HG23	1:C:432:TYR:CE1	2.39	0.57
1:C:156:ARG:HD3	7:C:605:HOH:O	2.04	0.56
1:C:2:ARG:N	1:C:131:GLY:O	2.38	0.56
2:D:147:SER:HG	2:D:190:SER:HG	1.50	0.56
2:D:248:LEU:HD23	2:D:354:ALA:HB2	1.86	0.56
2:D:106:GLY:O	2:D:111:GLY:N	2.36	0.56
1:C:390:ARG:HD3	7:C:773:HOH:O	2.05	0.55
2:D:278:ARG:HG2	2:D:278:ARG:HH11	1.72	0.54
1:C:339:ARG:HB3	1:C:341:ILE:CD1	2.38	0.54
1:C:36:MET:HE1	1:C:49:PHE:CZ	2.43	0.54
2:D:396:THR:O	2:D:400:ARG:HG3	2.09	0.53
2:D:66:ILE:HD13	2:D:122:VAL:HG22	1.91	0.53
2:D:283:TYR:CB	2:D:284:ARG:HA	2.38	0.52
1:C:2:ARG:NH2	1:C:50:ASN:HB3	2.22	0.52
2:D:396:THR:OG1	7:D:602:HOH:O	1.92	0.52
2:D:400:ARG:HD2	3:E:112:TRP:NE1	2.25	0.51
2:D:323:MET:HE2	2:D:373:MET:HG2	1.93	0.51
1:C:223:THR:HG23	1:C:226:ASN:H	1.75	0.51
3:E:139:VAL:HG23	7:E:242:HOH:O	2.11	0.51
2:D:140:SER:OG	5:D:501:GTP:H5'	2.10	0.51
2:D:406:HIS:CE1	2:D:407:TRP:CD1	3.00	0.50
1:C:11:GLN:O	1:C:15:GLN:HG3	2.13	0.48
1:C:256:GLN:NE2	7:C:623:HOH:O	2.45	0.48
1:C:108:TYR:O	1:C:112:LYS:HG2	2.13	0.48
1:C:266:HIS:HD2	7:C:601:HOH:O	1.97	0.48
2:D:171:VAL:HA	2:D:204:ILE:O	2.14	0.47
2:D:147:SER:OG	2:D:190:SER:OG	2.27	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:138:ASP:OD2	7:E:203:HOH:O	2.19	0.47
2:D:70:LEU:CB	2:D:98:GLY:HA2	2.44	0.47
1:C:28:HIS:CE1	1:C:49:PHE:HB3	2.49	0.47
1:C:293:ASN:HB3	1:C:339:ARG:HH12	1.79	0.47
2:D:283:TYR:CB	2:D:284:ARG:CA	2.93	0.47
2:D:278:ARG:HH11	2:D:278:ARG:CG	2.28	0.46
7:D:634:HOH:O	4:F:378:ARG:HD2	2.16	0.46
2:D:284:ARG:NH2	2:D:290:GLU:OE2	2.48	0.46
2:D:158:ARG:HH22	2:D:199:ASP:CG	2.22	0.46
2:D:406:HIS:CE1	2:D:407:TRP:HD1	2.33	0.46
2:D:167[B]:ASN:OD1	2:D:200:GLU:HG3	2.16	0.46
2:D:196:GLU:HB2	4:F:376:LEU:HD21	1.98	0.45
1:C:326:LYS:NZ	7:C:609:HOH:O	2.30	0.44
1:C:36:MET:HE3	1:C:61:HIS:NE2	2.33	0.44
2:D:75:MET:HG2	2:D:94:PHE:CD1	2.53	0.44
3:E:78:ILE:HD12	7:E:287:HOH:O	2.18	0.44
3:E:139:VAL:CG2	7:E:242:HOH:O	2.65	0.43
2:D:2:ARG:N	2:D:133:GLN:HG3	2.33	0.43
1:C:36:MET:HA	1:C:37:PRO:HD2	1.67	0.43
1:C:223:THR:HG22	1:C:226:ASN:OD1	2.18	0.43
1:C:413:MET:HE3	1:C:417:GLU:HB3	1.99	0.43
1:C:188:ILE:HG12	1:C:425:MET:HG3	2.01	0.43
2:D:432:TYR:OH	7:D:608:HOH:O	2.21	0.42
2:D:251:ASP:OD2	7:D:609:HOH:O	2.21	0.42
1:C:332:ILE:O	1:C:335:ILE:HG13	2.20	0.42
3:E:65:VAL:HG23	7:E:241:HOH:O	2.18	0.42
3:E:91:GLY:HA2	3:E:128:ILE:HD12	2.02	0.42
1:C:30:ILE:HG12	1:C:36:MET:HE2	2.02	0.42
1:C:223:THR:HG23	1:C:225:THR:N	2.34	0.42
2:D:109:THR:HG22	2:D:110:GLU:OE1	2.20	0.42
2:D:167[B]:ASN:HA	2:D:200:GLU:O	2.19	0.42
3:E:59:HIS:O	3:E:63:VAL:HG23	2.19	0.42
1:C:133:GLN:OE1	1:C:251:ASP:HB2	2.19	0.42
2:D:58:GLY:HA2	2:D:59:ASN:HA	1.65	0.41
2:D:180:THR:HG22	2:D:182:VAL:HG22	2.00	0.41
2:D:311:ARG:HH12	2:D:345:GLU:CD	2.28	0.41
1:C:36:MET:HE3	1:C:61:HIS:CD2	2.55	0.41
2:D:138:THR:HG22	2:D:169:PHE:HB2	2.03	0.41
2:D:320:ARG:HA	2:D:356:CYS:O	2.20	0.41
1:C:88:HIS:HB2	1:C:91:GLN:HG2	2.03	0.41
1:C:210:TYR:CZ	1:C:222:PRO:HD2	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:37:PRO:O	1:C:38:SER:HB2	2.20	0.41
2:D:272:PHE:CZ	2:D:275:LEU:HD23	2.56	0.41
1:C:156:ARG:NH1	7:C:605:HOH:O	2.14	0.41
2:D:283:TYR:HA	2:D:284:ARG:O	2.21	0.41
3:E:121:ALA:HB1	3:E:161:LEU:HD21	2.02	0.41
1:C:31:GLN:NE2	1:C:37:PRO:HD3	2.37	0.40
1:C:171:ILE:HD13	1:C:204:VAL:HB	2.03	0.40
3:E:109:VAL:HG12	3:E:115:THR:HG22	2.04	0.40
2:D:165:ILE:HG21	2:D:252:LEU:HB3	2.04	0.40
2:D:287:THR:HB	2:D:289:PRO:HD2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	425/451 (94%)	415 (98%)	10 (2%)	0	100	100
2	D	429/445 (96%)	422 (98%)	7 (2%)	0	100	100
3	E	154/169 (91%)	152 (99%)	2 (1%)	0	100	100
4	F	13/23 (56%)	13 (100%)	0	0	100	100
All	All	1021/1088 (94%)	1002 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	363/379 (96%)	355 (98%)	8 (2%)	45	53
2	D	370/383 (97%)	360 (97%)	10 (3%)	39	45
3	E	120/132 (91%)	118 (98%)	2 (2%)	53	62
4	F	12/19 (63%)	12 (100%)	0	100	100
All	All	865/913 (95%)	845 (98%)	20 (2%)	44	51

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	2	ARG
1	C	158	SER
1	C	178	SER
1	C	188	ILE
1	C	311	LYS
1	C	339	ARG
1	C	430	LYS
1	C	433	GLU
2	D	60	LYS
2	D	109	THR
2	D	110	GLU
2	D	117	SER
2	D	171	VAL
2	D	278	ARG
2	D	280	SER
2	D	286	LEU
2	D	325	MET
2	D	344	VAL
3	E	65	VAL
3	E	78	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	107	HIS
1	C	256	GLN
1	C	266	HIS
1	C	372	GLN
2	D	8	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	14	ASN
2	D	136	GLN
2	D	406	HIS
3	E	36	ASN
3	E	59	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GTP	D	501	6	33,34,34	1.04	3 (9%)	50,54,54	1.74	13 (26%)
5	GTP	C	501	6	33,34,34	2.45	16 (48%)	50,54,54	1.63	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	D	501	6	-	2/22/38/38	0/3/3/3
5	GTP	C	501	6	-	8/22/38/38	0/3/3/3

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	501	GTP	C5-N7	-4.94	1.29	1.39
5	C	501	GTP	PB-O1B	-4.55	1.35	1.50
5	C	501	GTP	PG-O3G	-4.15	1.39	1.54
5	C	501	GTP	C6-N1	-4.02	1.31	1.38
5	C	501	GTP	C4-N9	-3.92	1.28	1.38
5	C	501	GTP	PG-O1G	-3.72	1.38	1.50
5	C	501	GTP	PA-O1A	-3.61	1.38	1.50
5	C	501	GTP	C8-N9	-3.10	1.30	1.37
5	C	501	GTP	PB-O2B	-3.05	1.41	1.55
5	C	501	GTP	PA-O2A	-3.01	1.41	1.55
5	C	501	GTP	PG-O2G	-2.98	1.43	1.54
5	C	501	GTP	C5-C4	2.78	1.46	1.38
5	C	501	GTP	O4'-C4'	-2.57	1.39	1.45
5	C	501	GTP	PB-O3A	-2.46	1.56	1.59
5	D	501	GTP	PB-O3A	2.44	1.62	1.59
5	C	501	GTP	PB-O3B	-2.37	1.56	1.59
5	D	501	GTP	C5-N7	-2.22	1.34	1.39
5	C	501	GTP	O6-C6	-2.18	1.19	1.23
5	D	501	GTP	PA-O3A	2.12	1.61	1.59

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	501	GTP	C5-C4-N3	-5.24	120.04	128.39
5	C	501	GTP	C2-N3-C4	4.37	119.83	112.30
5	D	501	GTP	C5-C4-N3	-4.37	121.43	128.39
5	D	501	GTP	C2-N3-C4	4.33	119.76	112.30
5	C	501	GTP	N9-C4-N3	3.73	133.41	125.95
5	D	501	GTP	N9-C8-N7	-3.22	107.43	113.40
5	C	501	GTP	C6-C5-N7	3.16	136.03	130.29
5	D	501	GTP	C8-N7-C5	3.05	109.69	104.26
5	D	501	GTP	O2A-PA-O3A	3.04	115.49	107.27
5	D	501	GTP	N9-C4-N3	2.77	131.49	125.95
5	C	501	GTP	O3A-PB-O1B	-2.75	102.42	110.70
5	D	501	GTP	C2-N1-C6	-2.73	120.15	125.11
5	D	501	GTP	O5'-PA-O1A	-2.63	98.52	108.94
5	C	501	GTP	C4-C5-N7	-2.62	106.52	110.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	501	GTP	O3'-C3'-C4'	-2.55	103.76	111.08
5	C	501	GTP	O2A-PA-O3A	2.49	114.01	107.27
5	D	501	GTP	C5-C6-N1	2.44	119.45	113.25
5	D	501	GTP	O3B-PB-O1B	2.39	117.89	110.70
5	C	501	GTP	O2'-C2'-C3'	2.31	119.21	111.82
5	D	501	GTP	O6-C6-C5	-2.19	120.75	126.53
5	D	501	GTP	O2A-PA-O5'	2.03	116.77	107.57

There are no chirality outliers.

All (10) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	C	501	GTP	PB-O3B-PG-O2G
5	C	501	GTP	C5'-O5'-PA-O3A
5	C	501	GTP	C5'-O5'-PA-O1A
5	C	501	GTP	C5'-O5'-PA-O2A
5	D	501	GTP	PB-O3B-PG-O2G
5	C	501	GTP	PB-O3B-PG-O3G
5	D	501	GTP	PB-O3B-PG-O3G
5	C	501	GTP	C4'-C5'-O5'-PA
5	C	501	GTP	PB-O3A-PA-O1A
5	C	501	GTP	PB-O3A-PA-O2A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	501	GTP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	429/451 (95%)	-0.10	9 (2%) 63 66	14, 26, 48, 68	0
2	D	430/445 (96%)	0.10	18 (4%) 40 42	16, 30, 52, 70	1 (0%)
3	E	156/169 (92%)	-0.22	1 (0%) 85 87	14, 23, 42, 60	0
4	F	15/23 (65%)	0.41	0 100 100	28, 40, 51, 51	0
All	All	1030/1088 (94%)	-0.03	28 (2%) 56 59	14, 27, 49, 70	1 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	437	VAL	4.2
2	D	283	TYR	3.9
2	D	106	GLY	3.8
2	D	110	GLU	3.3
2	D	407	TRP	3.1
2	D	279	GLY	3.0
1	C	262	TYR	3.0
2	D	75	MET	2.9
1	C	2	ARG	2.9
1	C	47	ASP	2.8
2	D	111	GLY	2.8
2	D	441	ASP	2.7
1	C	50	ASN	2.7
2	D	58	GLY	2.6
2	D	282	GLN	2.5
1	C	365	GLY	2.5
2	D	109	THR	2.4
2	D	107	HIS	2.3
3	E	168	LEU	2.2
2	D	108	TYR	2.2
2	D	2	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	335	ILE	2.2
2	D	179	ASP	2.2
1	C	436	GLY	2.1
2	D	98	GLY	2.1
2	D	56	ALA	2.1
2	D	57	THR	2.0
1	C	359	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	MG	C	502	1/1	0.68	0.12	25,25,25,25	0
6	MG	D	502	1/1	0.96	0.07	32,32,32,32	0
5	GTP	D	501	32/32	0.97	0.06	18,23,31,39	0
5	GTP	C	501	32/32	0.98	0.05	10,16,21,36	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GTP D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GTP C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.