



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 24, 2026 – 01:03 PM UTC

PDB ID : 6EIF / pdb_00006eif
Title : DYRK1A in complex with XMD7-117
Authors : Rothweiler, U.
Deposited on : 2017-09-19
Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	 92% 7%
1	B	368	 87% 5% 8%
1	C	368	 83% 14%
1	D	368	 86% 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11105 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dual specificity tyrosine-phosphorylation-regulated kinase 1A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	344	Total	C	N	O	P	S	0	0	0
			2800	1802	479	501	1	17			
1	B	338	Total	C	N	O	P	S	0	0	0
			2739	1763	468	491	1	16			
1	C	317	Total	C	N	O	P	S	0	0	0
			2539	1632	427	462	1	17			
1	D	331	Total	C	N	O	P	S	0	0	0
			2693	1741	450	484	1	17			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	GLY	-	expression tag	UNP Q13627
A	124	ALA	-	expression tag	UNP Q13627
A	125	SER	-	expression tag	UNP Q13627
B	123	GLY	-	expression tag	UNP Q13627
B	124	ALA	-	expression tag	UNP Q13627
B	125	SER	-	expression tag	UNP Q13627
C	123	GLY	-	expression tag	UNP Q13627
C	124	ALA	-	expression tag	UNP Q13627
C	125	SER	-	expression tag	UNP Q13627
D	123	GLY	-	expression tag	UNP Q13627
D	124	ALA	-	expression tag	UNP Q13627
D	125	SER	-	expression tag	UNP Q13627

- Molecule 2 is 4-(3-pyridin-3-yl-1 {H}-pyrrolo[2,3-b]pyridin-5-yl)benzenesulfonamide (CCD ID: B5T) (formula: C₁₈H₁₄N₄O₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			25	18	4	2	1		
2	B	1	Total	C	N	O	S	0	0
			25	18	4	2	1		
2	C	1	Total	C	N	O	S	0	0
			25	18	4	2	1		
2	D	1	Total	C	N	O	S	0	0
			25	18	4	2	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	95	Total	O	0	0
			95	95		
3	B	31	Total	O	0	0
			31	31		
3	C	47	Total	O	0	0
			47	47		
3	D	61	Total	O	0	0
			61	61		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 1A

Chain A: 




- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 1A

Chain B: 




- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 1A

Chain C: 



- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 1A

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	88.44Å 88.54Å 231.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.57 – 2.22 48.57 – 2.22	Depositor EDS
% Data completeness (in resolution range)	99.1 (48.57-2.22) 96.6 (48.57-2.22)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.218 , 0.250 0.210 , 0.208	Depositor DCC
R_{free} test set	2101 reflections (2.33%)	wwPDB-VP
Wilson B-factor (Å ²)	33.8	Xtriage
Anisotropy	0.194	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 16.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.37$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	0.377 for k,h,-l	Xtriage
Reported twinning fraction	0.517 for H, K, L 0.483 for K, H, -L	Depositor
Outliers	1 of 89258 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11105	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: B5T, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/2848	0.78	0/3842
1	B	0.56	0/2785	0.75	0/3759
1	C	0.55	0/2577	0.76	0/3483
1	D	0.57	0/2739	0.79	0/3699
All	All	0.56	0/10949	0.77	0/14783

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2800	0	2787	5	0
1	B	2739	0	2710	10	0
1	C	2539	0	2444	4	0
1	D	2693	0	2672	9	0
2	A	25	0	0	1	0
2	B	25	0	0	0	0
2	C	25	0	0	0	0
2	D	25	0	0	0	0
3	A	95	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	31	0	0	0	0
3	C	47	0	0	0	0
3	D	61	0	0	0	0
All	All	11105	0	10613	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204:VAL:HG22	1:B:236:LEU:HD21	1.57	0.87
1:D:257:VAL:HG11	1:D:261:LEU:HD23	1.65	0.78
1:A:222:VAL:HG21	1:A:306:VAL:HG12	1.72	0.71
1:A:249:LEU:HD11	1:A:354:MET:HE1	1.74	0.69
1:B:224:LEU:HD11	1:B:236:LEU:HD22	1.79	0.64
1:C:455:LEU:HD11	1:C:459:MET:HE3	1.83	0.59
1:A:222:VAL:HG21	1:A:306:VAL:CG1	2.35	0.57
1:D:142:ASP:HB3	1:D:148:ILE:HD11	1.90	0.52
1:D:294:LEU:HG	1:D:306:VAL:HG21	1.92	0.50
1:B:210:MET:HE3	1:B:283:ILE:HD13	1.93	0.50
1:B:249:LEU:HD22	1:B:357:GLY:HA2	1.95	0.49
1:B:269:MET:HE1	1:B:305:ILE:HG13	1.93	0.49
1:A:165:ILE:HD12	2:A:501:B5T:CAI	2.42	0.49
1:B:224:LEU:CD1	1:B:236:LEU:HD22	2.44	0.48
1:A:222:VAL:HG22	1:A:241:LEU:HD11	1.94	0.48
1:C:289:LYS:HA	1:C:350:ILE:HD11	1.96	0.48
1:C:222:VAL:HG21	1:C:306:VAL:CG1	2.46	0.46
1:B:273:LEU:HD12	1:B:347:LEU:HD22	1.99	0.45
1:D:222:VAL:HG21	1:D:306:VAL:HG13	1.98	0.45
1:B:341:ALA:HB1	1:B:467:ARG:HD2	2.00	0.44
1:D:222:VAL:CG1	1:D:306:VAL:HG22	2.48	0.44
1:D:429:VAL:HG23	1:D:449:TYR:HB3	1.98	0.44
1:D:424:HIS:CE1	1:D:429:VAL:HG11	2.54	0.43
1:D:210:MET:HE2	1:D:283:ILE:HD13	2.01	0.43
1:D:252:THR:HG21	1:D:256:GLY:O	2.19	0.42
1:B:227:HIS:HB3	1:B:236:LEU:HD23	2.01	0.41
1:B:210:MET:HE1	1:B:308:PHE:CE2	2.56	0.41
1:C:352:VAL:O	1:C:356:THR:HG22	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/368 (92%)	328 (97%)	11 (3%)	0	100	100
1	B	331/368 (90%)	317 (96%)	14 (4%)	0	100	100
1	C	306/368 (83%)	295 (96%)	10 (3%)	1 (0%)	36	41
1	D	324/368 (88%)	308 (95%)	16 (5%)	0	100	100
All	All	1300/1472 (88%)	1248 (96%)	51 (4%)	1 (0%)	48	56

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	287	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/324 (93%)	299 (100%)	1 (0%)	86	93
1	B	292/324 (90%)	290 (99%)	2 (1%)	76	86
1	C	266/324 (82%)	264 (99%)	2 (1%)	73	84
1	D	290/324 (90%)	288 (99%)	2 (1%)	76	86
All	All	1148/1296 (89%)	1141 (99%)	7 (1%)	78	88

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	286	CYS
1	B	134	LYS
1	B	320	GLN
1	C	255	ARG
1	C	329	SER
1	D	209	LEU
1	D	215	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	213	HIS
1	A	371	ASN
1	A	404	ASN
1	B	192	ASN
1	B	223	HIS
1	B	232	ASN
1	B	260	ASN
1	B	313	GLN
1	B	323	GLN
1	C	213	HIS
1	C	297	ASN
1	C	313	GLN
1	C	371	ASN
1	C	404	ASN
1	C	424	HIS
1	D	227	HIS
1	D	297	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PTR	B	321	1	15,16,17	0.81	0	17,22,24	0.80	1 (5%)
1	PTR	D	321	1	15,16,17	0.81	0	17,22,24	0.84	0
1	PTR	C	321	1	15,16,17	0.83	0	17,22,24	0.76	0
1	PTR	A	321	1	15,16,17	0.86	0	17,22,24	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	B	321	1	-	5/10/11/13	0/1/1/1
1	PTR	D	321	1	-	0/10/11/13	0/1/1/1
1	PTR	C	321	1	-	1/10/11/13	0/1/1/1
1	PTR	A	321	1	-	1/10/11/13	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	321	PTR	OH-P-O1P	-2.12	102.41	109.48

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	321	PTR	O-C-CA-CB
1	B	321	PTR	O-C-CA-CB
1	B	321	PTR	CZ-OH-P-O1P
1	C	321	PTR	O-C-CA-CB
1	B	321	PTR	CZ-OH-P-O3P
1	B	321	PTR	N-CA-CB-CG
1	B	321	PTR	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	B5T	B	501	-	28,28,28	3.39	12 (42%)	41,41,41	2.12	10 (24%)
2	B5T	D	501	-	28,28,28	3.32	13 (46%)	41,41,41	2.06	8 (19%)
2	B5T	A	501	-	28,28,28	3.39	13 (46%)	41,41,41	2.14	11 (26%)
2	B5T	C	501	-	28,28,28	3.34	13 (46%)	41,41,41	2.18	9 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B5T	B	501	-	-	4/14/14/14	0/4/4/4
2	B5T	D	501	-	-	0/14/14/14	0/4/4/4
2	B5T	A	501	-	-	4/14/14/14	0/4/4/4
2	B5T	C	501	-	-	0/14/14/14	0/4/4/4

All (51) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	B5T	CAT-CAQ	-10.23	1.34	1.49
2	B	501	B5T	CAT-CAQ	-10.18	1.34	1.49
2	C	501	B5T	CAT-CAQ	-10.12	1.34	1.49
2	D	501	B5T	CAT-CAQ	-9.85	1.35	1.49
2	A	501	B5T	CAR-CAN	-6.18	1.34	1.41
2	B	501	B5T	CAR-CAN	-6.05	1.34	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	B5T	CAR-CAQ	-5.93	1.34	1.47
2	A	501	B5T	CAR-CAQ	-5.91	1.34	1.47
2	D	501	B5T	CAR-CAN	-5.90	1.35	1.41
2	C	501	B5T	CAR-CAQ	-5.86	1.34	1.47
2	C	501	B5T	CAR-CAN	-5.77	1.35	1.41
2	D	501	B5T	CAR-CAQ	-5.75	1.34	1.47
2	A	501	B5T	OAD-SAB	5.54	1.53	1.43
2	D	501	B5T	OAC-SAB	5.52	1.53	1.43
2	B	501	B5T	OAD-SAB	5.50	1.53	1.43
2	B	501	B5T	OAC-SAB	5.48	1.53	1.43
2	C	501	B5T	OAC-SAB	5.46	1.53	1.43
2	D	501	B5T	OAD-SAB	5.46	1.53	1.43
2	A	501	B5T	OAC-SAB	5.41	1.53	1.43
2	C	501	B5T	OAD-SAB	5.40	1.53	1.43
2	A	501	B5T	CAP-CAQ	-4.37	1.33	1.37
2	B	501	B5T	CAP-CAQ	-4.34	1.33	1.37
2	C	501	B5T	CAP-CAQ	-4.28	1.33	1.37
2	D	501	B5T	CAP-CAQ	-4.09	1.33	1.37
2	C	501	B5T	CAK-CAH	-3.61	1.40	1.49
2	D	501	B5T	CAK-CAH	-3.57	1.40	1.49
2	B	501	B5T	CAK-CAH	-3.53	1.40	1.49
2	B	501	B5T	CAU-NAV	3.36	1.41	1.34
2	A	501	B5T	CAU-NAV	3.33	1.41	1.34
2	D	501	B5T	CAU-NAV	3.29	1.41	1.34
2	C	501	B5T	CAU-NAV	3.23	1.41	1.34
2	A	501	B5T	CAK-CAH	-3.22	1.41	1.49
2	A	501	B5T	CAL-NAM	3.14	1.40	1.34
2	B	501	B5T	CAL-NAM	3.13	1.40	1.34
2	D	501	B5T	CAL-NAM	3.09	1.40	1.34
2	C	501	B5T	CAL-NAM	3.02	1.40	1.34
2	B	501	B5T	CAS-CAR	-3.00	1.35	1.39
2	A	501	B5T	CAS-CAR	-2.99	1.35	1.39
2	D	501	B5T	CAS-CAR	-2.84	1.35	1.39
2	C	501	B5T	CAS-CAR	-2.76	1.35	1.39
2	C	501	B5T	CAP-NAO	-2.63	1.32	1.37
2	C	501	B5T	CAW-NAV	2.61	1.41	1.33
2	B	501	B5T	CAW-NAV	2.60	1.41	1.33
2	D	501	B5T	CAP-NAO	-2.58	1.33	1.37
2	D	501	B5T	CAW-NAV	2.55	1.41	1.33
2	A	501	B5T	CAP-NAO	-2.55	1.33	1.37
2	A	501	B5T	CAW-NAV	2.55	1.41	1.33
2	B	501	B5T	CAP-NAO	-2.39	1.33	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	B5T	CAN-NAO	-2.18	1.33	1.36
2	D	501	B5T	CAN-NAO	-2.18	1.33	1.36
2	A	501	B5T	CAN-NAO	-2.04	1.33	1.36

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	B5T	OAD-SAB-OAC	-6.57	108.69	118.80
2	D	501	B5T	OAD-SAB-OAC	-6.48	108.83	118.80
2	B	501	B5T	OAD-SAB-OAC	-6.19	109.28	118.80
2	A	501	B5T	OAD-SAB-OAC	-6.16	109.33	118.80
2	C	501	B5T	CAR-CAQ-CAP	5.86	109.65	105.98
2	B	501	B5T	CAR-CAQ-CAP	5.74	109.58	105.98
2	A	501	B5T	CAR-CAQ-CAP	5.72	109.57	105.98
2	D	501	B5T	CAR-CAQ-CAP	5.15	109.21	105.98
2	B	501	B5T	CAQ-CAP-NAO	-4.76	107.39	110.46
2	A	501	B5T	CAQ-CAP-NAO	-4.76	107.39	110.46
2	C	501	B5T	CAQ-CAP-NAO	-4.67	107.45	110.46
2	A	501	B5T	CAR-CAN-NAM	-4.50	120.86	126.31
2	A	501	B5T	CAS-CAR-CAN	4.44	120.92	117.64
2	B	501	B5T	CAR-CAN-NAM	-4.34	121.05	126.31
2	C	501	B5T	CAS-CAR-CAN	4.28	120.80	117.64
2	D	501	B5T	CAQ-CAP-NAO	-4.27	107.71	110.46
2	C	501	B5T	CAR-CAN-NAM	-4.18	121.24	126.31
2	B	501	B5T	CAS-CAR-CAN	4.18	120.73	117.64
2	D	501	B5T	CAR-CAN-NAM	-4.14	121.30	126.31
2	D	501	B5T	CAS-CAR-CAN	3.75	120.41	117.64
2	B	501	B5T	CAK-CAL-NAM	-3.31	119.02	124.28
2	D	501	B5T	CAK-CAL-NAM	-3.22	119.16	124.28
2	A	501	B5T	CAK-CAL-NAM	-3.10	119.35	124.28
2	C	501	B5T	CAK-CAL-NAM	-2.95	119.59	124.28
2	D	501	B5T	OAD-SAB-CAE	2.67	110.36	107.35
2	C	501	B5T	OAD-SAB-CAE	2.64	110.34	107.35
2	B	501	B5T	OAC-SAB-CAE	2.49	110.17	107.35
2	A	501	B5T	NAO-CAN-NAM	2.33	130.46	126.69
2	B	501	B5T	NAO-CAN-NAM	2.25	130.34	126.69
2	C	501	B5T	CAT-CAU-NAV	-2.19	120.24	123.50
2	A	501	B5T	CAT-CAU-NAV	-2.19	120.25	123.50
2	A	501	B5T	OAC-SAB-CAE	2.16	109.80	107.35
2	A	501	B5T	CAL-NAM-CAN	2.16	120.43	116.19
2	B	501	B5T	CAL-NAM-CAN	2.12	120.34	116.19
2	B	501	B5T	CAT-CAU-NAV	-2.11	120.36	123.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	501	B5T	CAL-NAM-CAN	2.04	120.19	116.19
2	A	501	B5T	OAD-SAB-CAE	2.02	109.64	107.35
2	C	501	B5T	NAO-CAN-NAM	2.00	129.93	126.69

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	B5T	CAF-CAE-SAB-OAD
2	A	501	B5T	CAJ-CAE-SAB-OAD
2	B	501	B5T	CAF-CAE-SAB-OAD
2	B	501	B5T	CAJ-CAE-SAB-OAD
2	A	501	B5T	CAF-CAE-SAB-NAA
2	A	501	B5T	CAJ-CAE-SAB-NAA
2	B	501	B5T	CAF-CAE-SAB-NAA
2	B	501	B5T	CAJ-CAE-SAB-NAA


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	B5T	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand B5T B 501




Ligand B5T D 501



Ligand B5T A 501



Ligand B5T C 501



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	343/368 (93%)	-1.22	0 100 100	25, 36, 54, 69	0
1	B	337/368 (91%)	-0.88	0 100 100	34, 56, 82, 94	0
1	C	316/368 (85%)	-0.96	0 100 100	30, 52, 78, 91	0
1	D	330/368 (89%)	-1.16	0 100 100	29, 41, 61, 70	0
All	All	1326/1472 (90%)	-1.06	0 100 100	25, 45, 76, 94	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PTR	B	321	16/17	0.99	0.05	50,52,53,54	0
1	PTR	C	321	16/17	0.99	0.05	64,66,67,67	0
1	PTR	D	321	16/17	0.99	0.03	33,34,34,34	0
1	PTR	A	321	16/17	1.00	0.02	29,30,31,31	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	B5T	A	501	25/25	0.99	0.05	54,55,64,65	0
2	B5T	B	501	25/25	0.99	0.05	72,73,79,80	0
2	B5T	C	501	25/25	0.99	0.04	47,48,50,51	0
2	B5T	D	501	25/25	0.99	0.04	50,51,53,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around B5T A 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around B5T B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around B5T C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around B5T D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.