



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:58 PM UTC

PDB ID : 7EI3 / pdb_00007ei3
Title : Crystal structure of MasL, a thiolase from Massilia sp. YMA4
Authors : Lin, C.C.; Huang, K.F.; Yang, Y.L.
Deposited on : 2021-03-30
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	407	90% 5% .
1	B	407	93% . .
1	C	407	90% 5% . .
1	D	407	91% . . .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13180 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetyl-CoA C-acyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	0	0
			2872	1793	531	540	8			
1	B	393	Total	C	N	O	S	0	0	0
			2880	1799	532	541	8			
1	C	393	Total	C	N	O	S	0	0	0
			2880	1799	532	541	8			
1	D	394	Total	C	N	O	S	0	0	0
			2889	1805	534	542	8			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP A0A5C7BKK5
A	2	VAL	-	expression tag	UNP A0A5C7BKK5
A	183	THR	VAL	engineered mutation	UNP A0A5C7BKK5
A	395	LYS	-	expression tag	UNP A0A5C7BKK5
A	396	LEU	-	expression tag	UNP A0A5C7BKK5
A	397	ALA	-	expression tag	UNP A0A5C7BKK5
A	398	ALA	-	expression tag	UNP A0A5C7BKK5
A	399	ALA	-	expression tag	UNP A0A5C7BKK5
A	400	LEU	-	expression tag	UNP A0A5C7BKK5
A	401	GLU	-	expression tag	UNP A0A5C7BKK5
A	402	HIS	-	expression tag	UNP A0A5C7BKK5
A	403	HIS	-	expression tag	UNP A0A5C7BKK5
A	404	HIS	-	expression tag	UNP A0A5C7BKK5
A	405	HIS	-	expression tag	UNP A0A5C7BKK5
A	406	HIS	-	expression tag	UNP A0A5C7BKK5
A	407	HIS	-	expression tag	UNP A0A5C7BKK5
B	1	MET	-	initiating methionine	UNP A0A5C7BKK5
B	2	VAL	-	expression tag	UNP A0A5C7BKK5
B	183	THR	VAL	engineered mutation	UNP A0A5C7BKK5
B	395	LYS	-	expression tag	UNP A0A5C7BKK5
B	396	LEU	-	expression tag	UNP A0A5C7BKK5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	397	ALA	-	expression tag	UNP A0A5C7BKK5
B	398	ALA	-	expression tag	UNP A0A5C7BKK5
B	399	ALA	-	expression tag	UNP A0A5C7BKK5
B	400	LEU	-	expression tag	UNP A0A5C7BKK5
B	401	GLU	-	expression tag	UNP A0A5C7BKK5
B	402	HIS	-	expression tag	UNP A0A5C7BKK5
B	403	HIS	-	expression tag	UNP A0A5C7BKK5
B	404	HIS	-	expression tag	UNP A0A5C7BKK5
B	405	HIS	-	expression tag	UNP A0A5C7BKK5
B	406	HIS	-	expression tag	UNP A0A5C7BKK5
B	407	HIS	-	expression tag	UNP A0A5C7BKK5
C	1	MET	-	initiating methionine	UNP A0A5C7BKK5
C	2	VAL	-	expression tag	UNP A0A5C7BKK5
C	183	THR	VAL	engineered mutation	UNP A0A5C7BKK5
C	395	LYS	-	expression tag	UNP A0A5C7BKK5
C	396	LEU	-	expression tag	UNP A0A5C7BKK5
C	397	ALA	-	expression tag	UNP A0A5C7BKK5
C	398	ALA	-	expression tag	UNP A0A5C7BKK5
C	399	ALA	-	expression tag	UNP A0A5C7BKK5
C	400	LEU	-	expression tag	UNP A0A5C7BKK5
C	401	GLU	-	expression tag	UNP A0A5C7BKK5
C	402	HIS	-	expression tag	UNP A0A5C7BKK5
C	403	HIS	-	expression tag	UNP A0A5C7BKK5
C	404	HIS	-	expression tag	UNP A0A5C7BKK5
C	405	HIS	-	expression tag	UNP A0A5C7BKK5
C	406	HIS	-	expression tag	UNP A0A5C7BKK5
C	407	HIS	-	expression tag	UNP A0A5C7BKK5
D	1	MET	-	initiating methionine	UNP A0A5C7BKK5
D	2	VAL	-	expression tag	UNP A0A5C7BKK5
D	183	THR	VAL	engineered mutation	UNP A0A5C7BKK5
D	395	LYS	-	expression tag	UNP A0A5C7BKK5
D	396	LEU	-	expression tag	UNP A0A5C7BKK5
D	397	ALA	-	expression tag	UNP A0A5C7BKK5
D	398	ALA	-	expression tag	UNP A0A5C7BKK5
D	399	ALA	-	expression tag	UNP A0A5C7BKK5
D	400	LEU	-	expression tag	UNP A0A5C7BKK5
D	401	GLU	-	expression tag	UNP A0A5C7BKK5
D	402	HIS	-	expression tag	UNP A0A5C7BKK5
D	403	HIS	-	expression tag	UNP A0A5C7BKK5
D	404	HIS	-	expression tag	UNP A0A5C7BKK5
D	405	HIS	-	expression tag	UNP A0A5C7BKK5
D	406	HIS	-	expression tag	UNP A0A5C7BKK5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	407	HIS	-	expression tag	UNP A0A5C7BKK5

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	434	Total 434	O 434	0	0
2	B	406	Total 406	O 406	0	0
2	C	428	Total 428	O 428	0	0
2	D	391	Total 391	O 391	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetyl-CoA C-acyltransferase

Chain A: 




- Molecule 1: Acetyl-CoA C-acyltransferase

Chain B: 



- Molecule 1: Acetyl-CoA C-acyltransferase

Chain C: 



- Molecule 1: Acetyl-CoA C-acyltransferase

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	71.79Å 77.00Å 98.08Å 79.77° 79.00° 62.98°	Depositor
Resolution (Å)	29.90 – 1.78 29.90 – 1.78	Depositor EDS
% Data completeness (in resolution range)	92.0 (29.90-1.78) 92.0 (29.90-1.78)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.23 (at 1.78Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.128 , 0.180 0.145 , 0.193	Depositor DCC
R_{free} test set	7924 reflections (4.53%)	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13180	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	0/2919	1.19	1/3968 (0.0%)
1	B	0.98	0/2927	1.20	3/3979 (0.1%)
1	C	0.99	0/2927	1.19	1/3979 (0.0%)
1	D	0.98	0/2936	1.22	6/3990 (0.2%)
All	All	0.98	0/11709	1.20	11/15916 (0.1%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	131	ARG	CG-CD-NE	-9.77	90.51	112.00
1	D	131	ARG	NE-CZ-NH1	-8.37	113.13	121.50
1	D	131	ARG	NE-CZ-NH2	7.10	125.59	119.20
1	B	369	ARG	CG-CD-NE	-6.61	97.45	112.00
1	D	145	LEU	N-CA-CB	-6.11	101.14	110.12
1	D	4	ASP	N-CA-CB	5.33	117.84	110.17
1	B	15	PRO	N-CA-C	-5.21	103.59	111.41
1	A	15	PRO	N-CA-C	-5.20	103.61	111.41
1	D	15	PRO	N-CA-C	-5.17	103.65	111.41
1	C	15	PRO	N-CA-C	-5.03	103.87	111.41
1	B	105	ARG	CG-CD-NE	-5.01	100.97	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2872	0	2891	14	0
1	B	2880	0	2902	5	0
1	C	2880	0	2902	16	0
1	D	2889	0	2915	18	0
2	A	434	0	0	4	0
2	B	406	0	0	3	0
2	C	428	0	0	8	0
2	D	391	0	0	8	0
All	All	13180	0	11610	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:4:ASP:HB2	2:D:760:HOH:O	1.40	1.16
1:D:156:ASP:HB2	2:D:768:HOH:O	1.62	0.96
1:B:175:GLU:HG2	2:B:843:HOH:O	1.66	0.96
1:C:207:THR:HG22	1:C:212:GLU:HG2	1.55	0.88
1:A:156:ASP:HB3	2:A:764:HOH:O	1.80	0.82
1:C:207:THR:HG22	1:C:212:GLU:CG	2.21	0.70
1:C:4:ASP:N	2:C:501:HOH:O	2.31	0.62
1:A:136:MET:HE1	2:C:623:HOH:O	2.04	0.56
1:B:175:GLU:CB	2:B:843:HOH:O	2.53	0.56
1:A:175:GLU:HG3	2:A:730:HOH:O	2.05	0.56
1:D:310:GLN:HG2	2:D:675:HOH:O	2.08	0.52
1:B:140:GLN:HG2	1:B:142:LEU:HD23	1.91	0.52
1:C:350:HIS:HD2	1:C:355:THR:OG1	1.93	0.51
1:C:143:ASP:CG	1:D:131:ARG:HH12	2.18	0.51
1:A:280:ILE:H	1:A:304:GLN:NE2	2.08	0.51
1:D:140:GLN:HG2	1:D:142:LEU:HD23	1.93	0.50
1:C:350:HIS:HE1	2:C:819:HOH:O	1.92	0.50
1:D:362:LYS:HE3	2:D:600:HOH:O	2.10	0.50
1:C:140:GLN:HG2	1:C:142:LEU:HD23	1.94	0.49
1:C:143:ASP:OD2	1:D:131:ARG:NH1	2.40	0.49
1:D:373:ARG:NH1	2:D:503:HOH:O	2.46	0.49
2:C:675:HOH:O	1:D:105:ARG:HD2	2.13	0.48
1:C:362:LYS:HD3	2:C:534:HOH:O	2.14	0.48
1:D:48:GLU:HG3	2:D:732:HOH:O	2.14	0.48
1:A:223:PRO:HG3	2:A:566:HOH:O	2.13	0.47
1:C:276:VAL:CG1	1:C:392:GLU:HB3	2.46	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:GLN:HG2	1:A:142:LEU:HD23	1.98	0.46
1:C:362:LYS:CD	2:C:534:HOH:O	2.65	0.44
1:A:136:MET:CE	2:C:623:HOH:O	2.65	0.44
1:D:39:LEU:C	1:D:39:LEU:HD23	2.44	0.43
1:A:127:MET:HE2	1:A:127:MET:HB3	1.94	0.43
1:C:41:GLU:HG3	2:C:696:HOH:O	2.19	0.43
1:B:39:LEU:C	1:B:39:LEU:HD23	2.44	0.43
1:B:175:GLU:HB3	2:B:843:HOH:O	2.17	0.42
1:D:179:ARG:HG2	1:D:227:GLU:CD	2.43	0.42
1:D:41:GLU:HG3	2:D:522:HOH:O	2.20	0.42
1:D:395:LYS:O	1:D:396:LEU:HB3	2.19	0.42
1:A:34:HIS:CD2	1:A:204:VAL:HG12	2.55	0.42
1:A:48:GLU:H	1:A:48:GLU:CD	2.28	0.42
1:C:149:GLY:O	1:D:65:MET:HE2	2.18	0.42
1:C:39:LEU:C	1:C:39:LEU:HD23	2.45	0.42
1:D:396:LEU:C	2:D:504:HOH:O	2.63	0.42
1:C:395:LYS:HB3	1:C:395:LYS:HE3	1.77	0.41
1:D:3:LYS:HD2	1:D:105:ARG:NH2	2.34	0.41
1:D:48:GLU:H	1:D:48:GLU:CD	2.28	0.41
1:A:39:LEU:HD23	1:A:39:LEU:C	2.46	0.41
1:A:208:THR:O	1:A:209:ARG:C	2.62	0.41
1:A:156:ASP:CB	2:A:764:HOH:O	2.54	0.41
1:A:204:VAL:HG22	1:A:215:PHE:HB3	2.02	0.41
1:C:276:VAL:HG13	1:C:392:GLU:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	390/407 (96%)	380 (97%)	7 (2%)	3 (1%)	16 6

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	391/407 (96%)	381 (97%)	7 (2%)	3 (1%)	16	6
1	C	391/407 (96%)	380 (97%)	7 (2%)	4 (1%)	12	4
1	D	392/407 (96%)	382 (97%)	7 (2%)	3 (1%)	16	6
All	All	1564/1628 (96%)	1523 (97%)	28 (2%)	13 (1%)	16	6

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	209	ARG
1	C	209	ARG
1	A	352	ILE
1	B	352	ILE
1	C	352	ILE
1	D	352	ILE
1	A	89	VAL
1	B	89	VAL
1	B	209	ARG
1	C	89	VAL
1	D	89	VAL
1	D	209	ARG
1	C	66	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/298 (96%)	277 (97%)	9 (3%)	35	14
1	B	287/298 (96%)	282 (98%)	5 (2%)	53	36
1	C	287/298 (96%)	279 (97%)	8 (3%)	38	17
1	D	288/298 (97%)	281 (98%)	7 (2%)	43	24
All	All	1148/1192 (96%)	1119 (98%)	29 (2%)	42	22

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	GLU
1	A	52	GLU
1	A	59	LEU
1	A	89	VAL
1	A	136	MET
1	A	145	LEU
1	A	209	ARG
1	A	210	LYS
1	A	395	LYS
1	B	52	GLU
1	B	89	VAL
1	B	209	ARG
1	B	239	LYS
1	B	396	LEU
1	C	52	GLU
1	C	89	VAL
1	C	140	GLN
1	C	210	LYS
1	C	213	THR
1	C	373	ARG
1	C	395	LYS
1	C	396	LEU
1	D	48	GLU
1	D	52	GLU
1	D	89	VAL
1	D	127	MET
1	D	209	ARG
1	D	373	ARG
1	D	396	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	304	GLN
1	B	103	GLN
1	B	306	GLN
1	B	308	GLN
1	B	318	ASN
1	C	110	ASN
1	C	308	GLN
1	C	318	ASN
1	C	350	HIS
1	D	110	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	306	GLN
1	D	318	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/407 (96%)	-0.47	1 (0%) 90 93	13, 19, 37, 105	0
1	B	393/407 (96%)	-0.43	2 (0%) 87 91	13, 20, 38, 108	0
1	C	393/407 (96%)	-0.48	1 (0%) 90 93	13, 18, 34, 106	0
1	D	394/407 (96%)	-0.48	1 (0%) 90 93	13, 19, 36, 96	0
All	All	1572/1628 (96%)	-0.46	5 (0%) 90 93	13, 19, 37, 108	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	396	LEU	3.9
1	B	396	LEU	3.7
1	D	396	LEU	2.7
1	B	209	ARG	2.3
1	A	210	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.