



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 24, 2026 – 05:13 AM UTC

PDB ID : 7EIN / pdb_00007ein
Title : SARS-CoV-2 main proteinase complex with microbial metabolite leupeptin
Authors : Fu, L.F.; Feng, Y.; Qi, J.X.; Gao, G.F.
Deposited on : 2021-03-31
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	19% 86% 12% .
1	B	306	8% 88% 10% .
2	C	4	75% 25%
2	D	4	100%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5180 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	300	Total	C	N	O	S	0	2	0
			2319	1467	393	437	22			
1	B	300	Total	C	N	O	S	0	3	0
			2313	1465	389	436	23			

- Molecule 2 is a protein called leupeptin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	4	Total	C	N	O	0	0	0
			30	20	6	4			
2	D	4	Total	C	N	O	0	0	0
			30	20	6	4			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	230	Total	O	0	0
			230	230		
3	B	253	Total	O	0	0
			253	253		
3	C	2	Total	O	0	0
			2	2		
3	D	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3C-like proteinase



- Molecule 1: 3C-like proteinase



- Molecule 2: leupeptin



- Molecule 2: leupeptin



There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.04Å 53.12Å 113.64Å 90.00° 101.05° 90.00°	Depositor
Resolution (Å)	47.96 – 1.70 47.96 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.2 (47.96-1.70) 99.2 (47.96-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.78 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158, PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.235 , 0.260 0.238 , 0.262	Depositor DCC
R_{free} test set	2786 reflections (4.68%)	wwPDB-VP
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.132	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5180	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 93.68 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.6273e-09. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, OAR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.74	11/2373 (0.5%)	0.52	1/3227 (0.0%)
1	B	0.56	7/2373 (0.3%)	0.45	0/3228
2	C	0.34	0/15	0.79	0/19
2	D	0.33	0/15	0.88	0/19
All	All	0.65	18/4776 (0.4%)	0.48	1/6493 (0.0%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	33	ASP	C-O	-10.47	1.14	1.24
1	A	33	ASP	C-O	-10.31	1.14	1.24
1	B	35	VAL	C-O	-8.76	1.15	1.24
1	A	36	VAL	C-O	-8.72	1.14	1.24
1	A	32	LEU	C-O	-8.67	1.14	1.24
1	B	82	MET	C-O	-8.59	1.13	1.24
1	B	32	LEU	C-O	-8.10	1.14	1.24
1	A	83	GLN	C-O	-7.73	1.15	1.24
1	A	84	ASN	C-O	-7.60	1.14	1.24
1	A	152	ILE	C-O	-7.23	1.16	1.24
1	A	35	VAL	C-O	-6.89	1.17	1.24
1	A	153	ASP	C-O	-6.79	1.16	1.24
1	B	31	TRP	C-O	-6.65	1.15	1.23
1	A	26	THR	C-O	-5.89	1.16	1.24
1	B	34	ASP	C-O	-5.56	1.16	1.23
1	A	157	VAL	C-O	-5.50	1.18	1.24
1	A	31	TRP	C-O	-5.46	1.17	1.24
1	B	127	GLN	C-O	-5.27	1.17	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	152	ILE	CB-CA-C	-5.53	102.44	110.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2319	0	2263	18	0
1	B	2313	0	2258	15	0
2	C	30	0	36	0	0
2	D	30	0	36	0	0
3	A	230	0	0	7	0
3	B	253	0	0	2	0
3	C	2	0	0	0	0
3	D	3	0	0	0	0
All	All	5180	0	4593	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:ARG:HA	1:A:87:LEU:HG	1.79	0.65
1:A:7:ALA:HA	1:A:127:GLN:HG2	1.81	0.63
1:A:86:VAL:HG13	1:A:179:GLY:HA2	1.82	0.62
1:B:233:VAL:HG11	1:B:269:LYS:HG3	1.85	0.59
1:A:241:PRO:HD2	3:A:596:HOH:O	2.05	0.56
1:A:110:GLN:HG3	3:A:535:HOH:O	2.07	0.55
1:B:40:ARG:HA	1:B:87:LEU:HG	1.89	0.55
1:A:285:ALA:HB3	1:B:285:ALA:HB3	1.89	0.54
1:A:235:MET:HA	1:A:235:MET:HE2	1.91	0.53
1:B:235:MET:HA	1:B:235:MET:HE2	1.92	0.51
1:B:86:VAL:HG13	1:B:179:GLY:HA2	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:PRO:HG3	3:A:610:HOH:O	2.11	0.51
1:A:6:MET:O	1:A:127:GLN:HG3	2.11	0.50
1:A:217:ARG:HD2	3:A:449:HOH:O	2.12	0.50
1:B:279:ARG:NH2	3:B:412:HOH:O	2.46	0.48
1:A:76:ARG:HD3	3:A:530:HOH:O	2.15	0.46
1:A:298:ARG:NH2	3:A:410:HOH:O	2.42	0.46
1:A:7:ALA:HA	1:A:127:GLN:CG	2.44	0.46
1:B:78:ILE:HG13	1:B:90:LYS:HD3	1.98	0.45
1:B:113:SER:O	1:B:149:GLY:HA2	2.17	0.44
1:B:221:ASN:ND2	1:B:270:GLU:HG3	2.33	0.42
1:A:213:ILE:HG21	1:A:300:CYS:HB3	2.01	0.42
1:A:141:LEU:HD23	1:B:1:SER:OG	2.18	0.42
1:A:109:GLY:HA2	1:A:200:ILE:HD13	2.00	0.42
1:A:279:ARG:NH2	3:A:422:HOH:O	2.52	0.42
1:B:115:LEU:HD11	1:B:122:PRO:HB3	2.01	0.41
1:B:76:ARG:HD3	3:B:562:HOH:O	2.20	0.41
1:B:67:LEU:HD23	1:B:67:LEU:HA	1.93	0.41
1:B:163:HIS:CE1	1:B:172:HIS:HB3	2.56	0.41
1:A:163:HIS:CE1	1:A:172:HIS:HB3	2.55	0.41
1:B:95:ASN:HB3	1:B:98:THR:OG1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	300/306 (98%)	296 (99%)	3 (1%)	1 (0%)	36	22
1	B	301/306 (98%)	296 (98%)	4 (1%)	1 (0%)	36	22
2	C	2/4 (50%)	2 (100%)	0	0	100	100
2	D	2/4 (50%)	2 (100%)	0	0	100	100
All	All	605/620 (98%)	596 (98%)	7 (1%)	2 (0%)	36	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	TYR
1	B	154	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/263 (98%)	255 (99%)	3 (1%)	63	51
1	B	258/263 (98%)	258 (100%)	0	100	100
2	C	2/2 (100%)	2 (100%)	0	100	100
2	D	2/2 (100%)	2 (100%)	0	100	100
All	All	520/530 (98%)	517 (99%)	3 (1%)	78	72

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	SER
1	A	22	CYS
1	A	270	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	ASN
1	A	180	ASN
1	B	72	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	OAR	D	4	1,2	10,10,10	0.22	0	9,11,11	0.40	0
2	OAR	C	4	1,2	10,10,10	2.33	2 (20%)	9,11,11	2.20	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OAR	D	4	1,2	-	2/9/9/9	-
2	OAR	C	4	1,2	-	3/9/9/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	4	OAR	C-CA	6.56	1.63	1.52
2	C	4	OAR	CB-CA	3.26	1.57	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4	OAR	CB-CA-C	-6.48	103.39	112.25

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	4	OAR	CA-CB-CG-CD
2	D	4	OAR	NE-CD-CG-CB

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	C	4	OAR	CG-CD-NE-CZ
2	C	4	OAR	O-C-CA-N
2	D	4	OAR	O-C-CA-N

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	300/306 (98%)	1.16	59 (19%) 3 2	8, 21, 42, 54	2 (0%)
1	B	300/306 (98%)	0.54	25 (8%) 17 18	3, 14, 33, 43	3 (1%)
2	C	2/4 (50%)	1.97	0 100 100	24, 24, 24, 27	0
2	D	2/4 (50%)	0.85	0 100 100	14, 14, 14, 17	0
All	All	604/620 (97%)	0.85	84 (13%) 6 6	3, 17, 37, 54	5 (0%)

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	300	CYS	7.6
1	A	223	PHE	4.8
1	A	73	VAL	4.7
1	A	168	PRO	4.6
1	A	154	TYR	4.4
1	A	22	CYS	4.2
1	A	275	GLY	4.0
1	A	294	PHE	3.9
1	A	169	THR	3.9
1	B	24	THR	3.9
1	A	254[A]	SER	3.8
1	A	298	ARG	3.8
1	A	274	ASN	3.8
1	A	276	MET	3.7
1	B	3	PHE	3.6
1	A	119	ASN	3.6
1	A	270	GLU	3.6
1	A	153	ASP	3.6
1	A	277	ASN	3.5
1	A	194	ALA	3.4
1	A	118	TYR	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	235	MET	3.4
1	B	232	LEU	3.4
1	A	218	TRP	3.4
1	B	154	TYR	3.3
1	B	1	SER	3.3
1	B	300	CYS	3.2
1	A	1	SER	3.2
1	A	232	LEU	3.1
1	A	3	PHE	3.1
1	A	67	LEU	3.0
1	A	143	GLY	3.0
1	A	197	ASP	3.0
1	A	74	GLN	2.9
1	A	237	TYR	2.9
1	B	294	PHE	2.9
1	A	142	ASN	2.9
1	A	141	LEU	2.9
1	A	198	THR	2.8
1	A	60	ARG	2.8
1	A	167	LEU	2.8
1	A	229	ASP	2.7
1	A	273	GLN	2.7
1	B	235	MET	2.7
1	A	222	ARG	2.7
1	B	197	ASP	2.7
1	B	229	ASP	2.7
1	B	277	ASN	2.6
1	B	169	THR	2.6
1	A	44	CYS	2.6
1	A	127	GLN	2.6
1	A	128	CYS	2.5
1	B	73	VAL	2.5
1	A	71	GLY	2.5
1	B	74	GLN	2.5
1	B	236	LYS	2.5
1	B	168	PRO	2.5
1	A	47	GLU	2.5
1	A	227	LEU	2.4
1	A	271	LEU	2.4
1	B	47	GLU	2.4
1	B	23	GLY	2.4
1	A	272	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	233	VAL	2.4
1	A	236	LYS	2.4
1	A	299	GLN	2.3
1	A	215	GLY	2.3
1	A	216	ASP	2.3
1	A	234	ALA	2.3
1	B	153	ASP	2.3
1	B	22[A]	CYS	2.2
1	A	279	ARG	2.2
1	B	217	ARG	2.2
1	B	223	PHE	2.2
1	B	233	VAL	2.2
1	B	274	ASN	2.2
1	A	92	ASP	2.2
1	A	165	MET	2.2
1	A	2	GLY	2.2
1	A	72	ASN	2.1
1	A	256	GLN	2.1
1	A	59	ILE	2.1
1	A	221	ASN	2.1
1	B	198	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	OAR	C	4	11/11	0.82	0.15	24,27,37,38	0
2	OAR	D	4	11/11	0.87	0.12	14,16,29,29	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.