



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 26, 2026 – 01:27 PM EDT

PDB ID : 8EIL / pdb\_00008eil  
Title : C-Terminal Domain of BrxL from Acinetobacter BREX type I phage restriction system  
Authors : Doyle, L.A.; Stoddard, B.L.; Kaiser, B.  
Deposited on : 2022-09-15  
Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	80% 17% . .
1	B	186	83% 13% . .
1	C	186	80% 15% . .
1	D	186	81% 13% . 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SIN	D	701	-	X	-	-

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 5208 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protease Lon-related BREX system protein BrxL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	181	1290	833	208	243	6	0	0	0
1	B	180	1300	838	209	246	7	0	0	0
1	C	178	1243	799	206	233	5	0	0	0
1	D	177	1233	792	205	231	5	0	0	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	494	GLY	-	expression tag	UNP A0A3R9EDI8
A	495	SER	-	expression tag	UNP A0A3R9EDI8
A	496	HIS	-	expression tag	UNP A0A3R9EDI8
A	497	MET	-	expression tag	UNP A0A3R9EDI8
B	494	GLY	-	expression tag	UNP A0A3R9EDI8
B	495	SER	-	expression tag	UNP A0A3R9EDI8
B	496	HIS	-	expression tag	UNP A0A3R9EDI8
B	497	MET	-	expression tag	UNP A0A3R9EDI8
C	494	GLY	-	expression tag	UNP A0A3R9EDI8
C	495	SER	-	expression tag	UNP A0A3R9EDI8
C	496	HIS	-	expression tag	UNP A0A3R9EDI8
C	497	MET	-	expression tag	UNP A0A3R9EDI8
D	494	GLY	-	expression tag	UNP A0A3R9EDI8
D	495	SER	-	expression tag	UNP A0A3R9EDI8
D	496	HIS	-	expression tag	UNP A0A3R9EDI8
D	497	MET	-	expression tag	UNP A0A3R9EDI8

- Molecule 2 is MALONIC ACID (CCD ID: MLA) (formula: C<sub>3</sub>H<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	C O	0	0
			7	3 4		
2	B	1	Total	C O	0	0
			7	3 4		

- Molecule 3 is SUCCINIC ACID (CCD ID: SIN) (formula:  $C_4H_6O_4$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	C O	0	0
			8	4 4		

- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	40	Total 40	O 40	0	0
4	B	35	Total 35	O 35	0	0
4	C	24	Total 24	O 24	0	0
4	D	21	Total 21	O 21	0	0



## 4 Data and refinement statistics i

Property	Value	Source
Space group	F 2 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	213.34Å 213.34Å 213.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.94 – 2.25 48.94 – 2.25	Depositor EDS
% Data completeness (in resolution range)	92.6 (48.94-2.25) 92.1 (48.94-2.25)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.13 (at 2.24Å)	Xtrriage
Refinement program	PHENIX 1.18_3855	Depositor
R, $R_{free}$	0.216 , 0.230 0.216 , 0.231	Depositor DCC
$R_{free}$ test set	2024 reflections (5.33%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.3	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 63.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.499 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5208	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIN, MLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.16	0/1317	0.34	0/1794
1	B	0.16	0/1327	0.37	0/1807
1	C	0.16	0/1270	0.39	0/1731
1	D	0.15	0/1260	0.33	0/1719
All	All	0.16	0/5174	0.36	0/7051

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1290	0	1226	16	0
1	B	1300	0	1252	13	0
1	C	1243	0	1144	14	0
1	D	1233	0	1119	15	0
2	A	7	0	2	0	0
2	B	7	0	2	0	0
3	D	8	0	4	0	0
4	A	40	0	0	0	0
4	B	35	0	0	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	24	0	0	0	0
4	D	21	0	0	0	0
All	All	5208	0	4749	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:520:LEU:HD22	1:D:593:PHE:HA	1.67	0.74
1:C:520:LEU:HD22	1:C:593:PHE:HA	1.70	0.73
1:D:516:TYR:HD1	1:D:575:GLU:HA	1.59	0.67
1:C:516:TYR:OH	1:C:579:THR:OG1	2.14	0.65
1:D:582:LEU:HD22	1:D:584:HIS:H	1.62	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/186 (95%)	176 (99%)	1 (1%)	0	100	100
1	B	176/186 (95%)	173 (98%)	3 (2%)	0	100	100
1	C	172/186 (92%)	167 (97%)	4 (2%)	1 (1%)	21	21
1	D	171/186 (92%)	168 (98%)	3 (2%)	0	100	100
All	All	696/744 (94%)	684 (98%)	11 (2%)	1 (0%)	48	56

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	561	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/150 (84%)	119 (94%)	7 (6%)	19	20
1	B	132/150 (88%)	126 (96%)	6 (4%)	24	29
1	C	116/150 (77%)	107 (92%)	9 (8%)	11	10
1	D	113/150 (75%)	105 (93%)	8 (7%)	13	12
All	All	487/600 (81%)	457 (94%)	30 (6%)	16	16

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	532	SER
1	D	617	SER
1	C	590	LEU
1	D	667	ASP
1	D	582	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	543	GLN
1	D	554	ASN
1	B	661	GLN
1	C	543	GLN
1	C	570	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MLA	A	701	-	6,6,6	1.37	0	7,7,7	1.30	0
3	SIN	D	701	-	7,7,7	1.04	0	8,8,8	1.66	3 (37%)
2	MLA	B	701	-	6,6,6	1.42	0	7,7,7	1.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLA	A	701	-	-	2/4/4/4	-
3	SIN	D	701	-	-	5/5/5/5	-
2	MLA	B	701	-	-	2/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	701	SIN	C2-C3-C4	-2.04	108.25	113.67
3	D	701	SIN	O2-C1-C2	2.03	120.42	114.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	701	SIN	O4-C4-C3	2.02	120.39	114.00

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	701	SIN	C1-C2-C3-C4
2	B	701	MLA	O1A-C1-C2-C3
2	B	701	MLA	O1B-C1-C2-C3
2	A	701	MLA	O1B-C1-C2-C3
2	A	701	MLA	O1A-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	181/186 (97%)	-1.21	0 100 100	32, 53, 82, 98	0
1	B	180/186 (96%)	-1.20	0 100 100	34, 53, 84, 95	0
1	C	178/186 (95%)	-1.03	0 100 100	34, 71, 101, 115	0
1	D	177/186 (95%)	-1.05	0 100 100	35, 71, 98, 121	0
All	All	716/744 (96%)	-1.12	0 100 100	32, 61, 95, 121	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MLA	A	701	7/7	0.98	0.03	57,62,73,74	0
2	MLA	B	701	7/7	0.98	0.04	59,61,71,72	0
3	SIN	D	701	8/8	0.99	0.03	55,67,68,71	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.