



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:31 PM UTC

PDB ID : 3EJP / pdb_00003ejp
Title : Golgi alpha-Mannosidase II in complex with 5-substituted swainsonine analog:
(5R)-5-[2'-oxo-2'-(phenyl)ethyl]-swainsonine
Authors : Kuntz, D.A.; Rose, D.R.
Deposited on : 2008-09-18
Resolution : 1.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2531 (1.34-1.30)
Clashscore	190562	2585 (1.34-1.30)
Ramachandran outliers	187476	2528 (1.34-1.30)
Sidechain outliers	187428	2528 (1.34-1.30)
RSRZ outliers	180081	2528 (1.34-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1045	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9759 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alpha-mannosidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1016	Total	C	N	O	S	0	42	0
			8415	5345	1480	1548	42			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ARG	-	expression tag	UNP Q24451
A	2	SER	-	expression tag	UNP Q24451
A	3	SER	-	expression tag	UNP Q24451
A	4	HIS	-	expression tag	UNP Q24451
A	5	HIS	-	expression tag	UNP Q24451
A	6	HIS	-	expression tag	UNP Q24451
A	7	HIS	-	expression tag	UNP Q24451
A	8	HIS	-	expression tag	UNP Q24451
A	9	HIS	-	expression tag	UNP Q24451
A	10	GLY	-	expression tag	UNP Q24451
A	11	GLU	-	expression tag	UNP Q24451
A	12	PHE	-	expression tag	UNP Q24451
A	907	LYS	GLU	SEE REMARK 999	UNP Q24451

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is (4R)-2-METHYLPENTANE-2,4-DIOL (CCD ID: MRD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is 1-phenyl-2-[(1S,2R,5R,8R,8aR)-1,2,8-trihydroxyoctahydroindolizin-5-yl]ethanone (CCD ID: HN2) (formula: C₁₆H₂₁NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			21	16	1	4		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1295	Total	O	0	5
			1300	1300		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.03Å 109.88Å 138.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.22 – 1.32 29.22 – 1.32	Depositor EDS
% Data completeness (in resolution range)	92.1 (29.22-1.32) 92.1 (29.22-1.32)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 1.32Å)	Xtriage
Refinement program	REFMAC, CNS	Depositor
R, R_{free}	0.157 , 0.182 0.155 , 0.153	Depositor DCC
R_{free} test set	5091 reflections (2.24%)	wwPDB-VP
Wilson B-factor (Å ²)	11.4	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9759	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MRD, HN2, NAG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	1/8781 (0.0%)	0.75	1/11914 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	772	GLY	N-CA	5.08	1.49	1.44

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	194	ASN	N-CA-C	5.22	118.65	111.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8415	0	8211	55	0
2	A	14	0	13	1	0
3	A	1	0	0	0	0
4	A	8	0	14	2	0
5	A	21	0	19	0	0
6	A	1300	0	0	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	9759	0	8257	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:610:GLN:HG3	6:A:1827:HOH:O	1.41	1.21
1:A:498[B]:GLN:HG2	1:A:526:LEU:HD12	1.16	1.12
1:A:136[A]:ARG:HD3	6:A:1338:HOH:O	1.72	0.89
1:A:599[A]:HIS:CD2	6:A:1827:HOH:O	2.34	0.80
1:A:75[A]:TYR:HD1	6:A:2154:HOH:O	1.65	0.79
1:A:96:ILE:CG2	6:A:2101:HOH:O	2.32	0.77
1:A:96:ILE:HG22	6:A:2101:HOH:O	1.84	0.76
1:A:380:GLN:HG2	6:A:1643:HOH:O	1.84	0.76
1:A:541:THR:HG23	6:A:1239:HOH:O	1.87	0.74
1:A:136[B]:ARG:HD2	6:A:1880:HOH:O	1.88	0.71
1:A:82[A]:LYS:HE3	1:A:375[A]:GLN:NE2	2.06	0.70
1:A:82[A]:LYS:HD2	1:A:375[A]:GLN:HE22	1.58	0.69
1:A:280:LYS:NZ	1:A:301:VAL:HG21	2.07	0.68
1:A:599[A]:HIS:HD2	1:A:610:GLN:CG	2.06	0.67
1:A:651[B]:LEU:HD22	1:A:659:LEU:HD11	1.77	0.67
1:A:1011[B]:ARG:NH1	6:A:1681:HOH:O	2.27	0.66
1:A:640:LYS:HD2	1:A:646:TYR:CE2	2.31	0.65
1:A:599[A]:HIS:CD2	1:A:610:GLN:HG3	2.32	0.64
1:A:678:LEU:HD12	1:A:769[A]:MET:HE1	1.78	0.64
1:A:280:LYS:HZ2	1:A:301:VAL:HG21	1.61	0.63
4:A:1048:MRD:H1C3	6:A:1112:HOH:O	1.98	0.63
1:A:979[A]:LEU:HD21	1:A:999:VAL:HG11	1.81	0.62
1:A:904:ARG:HG2	1:A:985:MET:SD	2.40	0.61
1:A:742:GLN:HG3	6:A:1438:HOH:O	2.02	0.60
4:A:1048:MRD:H1C1	6:A:2192:HOH:O	2.00	0.59
1:A:47:GLU:OE2	1:A:51:ARG:HD3	2.03	0.58
1:A:975:VAL:HG21	1:A:1003:LEU:CD1	2.34	0.58
1:A:138:TYR:CE1	1:A:193:MET:HE1	2.40	0.57
1:A:599[A]:HIS:HD2	1:A:610:GLN:HG3	1.68	0.56
1:A:498[B]:GLN:HG2	1:A:526:LEU:CD1	2.11	0.55
1:A:599[A]:HIS:CD2	1:A:610:GLN:CG	2.90	0.54
1:A:678:LEU:CD1	1:A:769[A]:MET:HE1	2.38	0.54
2:A:1046:NAG:H83	2:A:1046:NAG:H3	1.90	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48:LEU:HD11	1:A:236:GLU:HG2	1.90	0.53
1:A:653:ARG:HD3	1:A:654:LYS:O	2.09	0.52
1:A:895:VAL:HG12	1:A:897:GLU:HG3	1.92	0.52
1:A:62:TRP:CD2	1:A:65:GLY:HA3	2.45	0.52
1:A:288:LYS:NZ	6:A:2093:HOH:O	2.43	0.52
1:A:47:GLU:OE2	1:A:51:ARG:CD	2.57	0.51
1:A:82[A]:LYS:CE	1:A:375[A]:GLN:NE2	2.73	0.51
1:A:174:HIS:CE1	1:A:176:ARG:HD3	2.46	0.49
1:A:653:ARG:HD2	1:A:656:PRO:HA	1.94	0.49
1:A:686:LEU:HD22	1:A:697:ILE:HG12	1.95	0.48
1:A:707:PRO:HG2	1:A:797[A]:HIS:CE1	2.48	0.48
1:A:280:LYS:HZ2	1:A:301:VAL:CG2	2.26	0.48
1:A:251:TRP:C	1:A:251:TRP:CD1	2.91	0.48
1:A:313:ALA:O	1:A:316[B]:ASP:HB2	2.14	0.47
1:A:82[A]:LYS:CD	1:A:375[A]:GLN:HE22	2.26	0.47
1:A:651[B]:LEU:HD11	1:A:744:VAL:CG1	2.45	0.47
1:A:491[A]:GLN:NE2	6:A:1552:HOH:O	2.30	0.44
1:A:96:ILE:HG23	1:A:479:LYS:HE2	1.99	0.44
1:A:96:ILE:HG23	1:A:479:LYS:CE	2.49	0.43
1:A:48:LEU:HG	1:A:52[A]:MET:SD	2.59	0.42
1:A:204:ASP:H	1:A:205:PRO:HD3	1.85	0.42
1:A:599[A]:HIS:HD2	1:A:610:GLN:HG2	1.84	0.41
1:A:117[B]:HIS:HE1	1:A:351:ASP:OD1	2.04	0.41
1:A:96:ILE:HG22	1:A:96:ILE:O	2.19	0.40
1:A:225:LEU:HD21	1:A:264[A]:MET:SD	2.61	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1056/1045 (101%)	1030 (98%)	24 (2%)	2 (0%)	43	17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	95	TRP
1	A	204	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	944/929 (102%)	938 (99%)	6 (1%)	78 54

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	275	CYS
1	A	585	ASN
1	A	651[A]	LEU
1	A	651[B]	LEU
1	A	653	ARG
1	A	898	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	ASN
1	A	347	ASN
1	A	371	ASN
1	A	923	GLN
1	A	1005	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1046	1	14,14,15	0.64	0	17,19,21	2.96	5 (29%)
4	MRD	A	1048	-	7,7,7	0.35	0	9,10,10	0.75	0
5	HN2	A	1049	-	23,23,23	0.86	0	27,33,33	0.98	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1046	1	-	6/6/23/26	0/1/1/1
4	MRD	A	1048	-	-	2/5/5/5	-
5	HN2	A	1049	-	-	0/8/37/37	0/3/3/3

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1046	NAG	C1-O5-C5	8.10	123.05	112.19
2	A	1046	NAG	C2-N2-C7	6.40	131.47	122.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1046	NAG	O5-C1-C2	3.59	116.85	111.29
2	A	1046	NAG	C8-C7-N2	3.21	121.44	116.12
2	A	1046	NAG	O5-C5-C4	2.53	116.98	110.83
5	A	1049	HN2	C6-C5-N4	-2.21	105.31	110.65

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1048	MRD	C2-C3-C4-O4
4	A	1048	MRD	C2-C3-C4-C5
2	A	1046	NAG	C8-C7-N2-C2
2	A	1046	NAG	O7-C7-N2-C2
2	A	1046	NAG	C4-C5-C6-O6
2	A	1046	NAG	O5-C5-C6-O6
2	A	1046	NAG	C3-C2-N2-C7
2	A	1046	NAG	C1-C2-N2-C7

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1046	NAG	1	0
4	A	1048	MRD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	1016/1045 (97%)	0.25	63 (6%)	26 34	6, 12, 23, 44	42 (4%)

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	993	HIS	7.8
1	A	1045	SER	6.6
1	A	990	PRO	5.8
1	A	614	THR	5.4
1	A	991	GLU	5.1
1	A	703	SER	4.8
1	A	583	LEU	4.7
1	A	613	THR	4.4
1	A	655	ASN	4.2
1	A	1028	ALA	4.1
1	A	534	SER	4.0
1	A	702	ASP	3.6
1	A	992	GLU	3.5
1	A	662	GLY	3.4
1	A	1024	ASP	3.3
1	A	994	THR	3.3
1	A	654	LYS	3.3
1	A	676	ILE	3.3
1	A	682	ASN	3.3
1	A	638	ASP	3.2
1	A	989	THR	3.1
1	A	661	LEU	3.1
1	A	535	GLY	3.1
1	A	988	GLY	3.1
1	A	78	HIS	3.0
1	A	144	ASN	3.0
1	A	290	MET	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	521	PHE	3.0
1	A	77	ALA	3.0
1	A	537	GLU	2.9
1	A	538	ASP	2.9
1	A	368	ALA	2.9
1	A	602	THR	2.8
1	A	392	ALA	2.8
1	A	396	GLU	2.7
1	A	347	ASN	2.7
1	A	683	GLY	2.7
1	A	704	PRO	2.7
1	A	603	LEU	2.7
1	A	1030	GLU	2.7
1	A	700	THR	2.6
1	A	720	HIS	2.6
1	A	701	GLN	2.6
1	A	539	SER	2.6
1	A	721	GLY	2.5
1	A	784	SER	2.4
1	A	1022	HIS	2.4
1	A	1029	PRO	2.4
1	A	659	LEU	2.4
1	A	653	ARG	2.4
1	A	940	ILE	2.3
1	A	1027	VAL	2.3
1	A	663	GLN	2.3
1	A	987	CYS	2.2
1	A	451	MET	2.2
1	A	541	THR	2.2
1	A	291	GLY	2.2
1	A	739	GLU	2.2
1	A	57[A]	ILE	2.1
1	A	313	ALA	2.1
1	A	616	TYR	2.1
1	A	706	VAL	2.1
1	A	536	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	A	1046	14/15	0.61	0.17	34,40,45,46	0
4	MRD	A	1048	8/8	0.92	0.09	14,18,20,22	0
5	HN2	A	1049	21/21	0.98	0.04	7,9,15,16	0
3	ZN	A	1047	1/1	1.00	0.01	8,8,8,8	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.