



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:32 PM UTC

PDB ID : 5EJR / pdb_00005ejr
Title : Structure of Dictyostelium Discoideum Myosin VII MyTH4-FERM MF2 domain
Authors : Planelles-Herrero, V.J.; Sirkia, H.; Sourigues, Y.; Clause, J.; Titus, M.A.; Houdusse, A.
Deposited on : 2015-11-02
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	507	8% 90% 8% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4388 atoms, of which 2 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin-I heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	503	4087	2607	679	777	24	0	5	0

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	H		
2	A	1	4	2	2	0	0	0
2	A	1	4	2	2	0	0	0
2	A	1	4	2	2	0	0	0
2	A	1	4	2	2	0	0	0
2	A	1	5	2	1	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C H O 5 2 1 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	271	Total O 271 271	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Myosin-I heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	40.02Å 58.95Å 64.36Å 90.07° 96.78° 108.48°	Depositor
Resolution (Å)	41.22 – 2.00 41.22 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.5 (41.22-2.00) 97.5 (41.22-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 2.00Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.10.1	Depositor
R, R_{free}	0.179 , 0.218 0.186 , 0.224	Depositor DCC
R_{free} test set	1798 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	24.7	Xtriage
Anisotropy	0.343	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.012 for -h,h+k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4388	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/4170	1.20	4/5633 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1950	ASN	CA-C-N	6.11	128.75	120.38
1	A	1950	ASN	C-N-CA	6.11	128.75	120.38
1	A	2122	ASP	CA-CB-CG	5.97	118.57	112.60
1	A	2085	ILE	N-CA-CB	5.47	118.79	110.58

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4087	0	4069	19	0
2	A	28	2	42	3	0
3	A	271	0	0	1	0
All	All	4386	2	4111	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2317:VAL:HG12	1:A:2326:ILE:HG22	1.80	0.64
1:A:2149:PHE:CZ	2:A:2407:EDO:H21	2.33	0.63
1:A:1995:LEU:HG	1:A:1999:MET:HE2	1.85	0.58
1:A:1952:GLU:H	1:A:1952:GLU:CD	2.15	0.55
1:A:2074:ALA:HB2	1:A:2089:VAL:HG13	1.87	0.55
1:A:2299:ASP:HB2	1:A:2306:LYS:HE3	1.90	0.54
1:A:2186:TYR:OH	2:A:2407:EDO:H12	2.08	0.53
1:A:2063:LYS:HG2	1:A:2071:THR:HG21	1.95	0.48
1:A:2067:ILE:HG23	1:A:2175[A]:CYS:SG	2.54	0.48
1:A:1972:GLU:C	3:A:2505:HOH:O	2.56	0.48
1:A:2111:THR:OG1	1:A:2138:GLY:HA3	2.16	0.46
1:A:2072:LYS:HD2	1:A:2092:GLN:HB3	1.99	0.45
1:A:2189:VAL:HG22	1:A:2193:ASP:HB2	1.98	0.45
1:A:2085:ILE:HD13	1:A:2124:ILE:HD12	1.99	0.43
1:A:2084:GLU:O	1:A:2088:THR:HG23	2.19	0.43
1:A:2180:GLU:HG2	1:A:2185:LEU:HD12	2.02	0.42
1:A:2083:ARG:HB2	1:A:2120:GLU:HB2	2.02	0.42
1:A:1923:MET:HE3	1:A:1930:ILE:HA	2.03	0.40
1:A:2149:PHE:CE1	2:A:2407:EDO:H21	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	504/507 (99%)	488 (97%)	14 (3%)	2 (0%)	30 27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2113	GLY
1	A	1887	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	464/466 (100%)	453 (98%)	11 (2%)	43	47

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1873	LEU
1	A	1886	VAL
1	A	1949	GLU
1	A	1952	GLU
1	A	1972	GLU
1	A	2009	ILE
1	A	2010	GLN
1	A	2247	LEU
1	A	2277	ILE
1	A	2303	LYS
1	A	2304	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1874	ASN
1	A	1941	GLN
1	A	2001	GLN
1	A	2077	ASN
1	A	2188	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	2404	-	3,3,3	0.76	0	2,2,2	0.49	0
2	EDO	A	2405	-	3,3,3	0.47	0	2,2,2	0.66	0
2	EDO	A	2402	-	3,3,3	0.59	0	2,2,2	0.38	0
2	EDO	A	2407	-	3,3,3	0.61	0	2,2,2	0.51	0
2	EDO	A	2401	-	3,3,3	0.48	0	2,2,2	0.58	0
2	EDO	A	2406	-	3,3,3	0.52	0	2,2,2	0.36	0
2	EDO	A	2403	-	3,3,3	0.48	0	2,2,2	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	2404	-	-	0/1/1/1	-
2	EDO	A	2405	-	-	0/1/1/1	-
2	EDO	A	2402	-	-	1/1/1/1	-
2	EDO	A	2407	-	-	1/1/1/1	-
2	EDO	A	2401	-	-	0/1/1/1	-
2	EDO	A	2406	-	-	0/1/1/1	-
2	EDO	A	2403	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2402	EDO	O1-C1-C2-O2
2	A	2407	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2407	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	503/507 (99%)	0.39	41 (8%) 17 16	10, 37, 74, 111	5 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1933	GLY	3.6
1	A	2034	SER	3.5
1	A	1886	VAL	3.5
1	A	1880	LEU	3.4
1	A	2035	TYR	3.4
1	A	2303	LYS	3.2
1	A	2076	ILE	3.1
1	A	1881	LYS	3.0
1	A	2208	ASN	3.0
1	A	1852	THR	2.8
1	A	2077	ASN	2.8
1	A	1870	ALA	2.7
1	A	1872	THR	2.7
1	A	2328	THR	2.7
1	A	1878	ALA	2.6
1	A	1879	THR	2.6
1	A	1885	PRO	2.6
1	A	2032	ILE	2.6
1	A	1875	GLN	2.6
1	A	1873	LEU	2.5
1	A	1883	LYS	2.4
1	A	2333	LYS	2.4
1	A	1896	LYS	2.4
1	A	2014	PRO	2.4
1	A	2112	ALA	2.3
1	A	1887	ASP	2.3
1	A	2011	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2304	GLU	2.3
1	A	2054	LEU	2.3
1	A	2004	SER	2.2
1	A	2059	ILE	2.2
1	A	2280	LYS	2.2
1	A	2334	PRO	2.2
1	A	1890	THR	2.1
1	A	2016	LEU	2.1
1	A	1893	TYR	2.1
1	A	2300	PRO	2.1
1	A	2335	ILE	2.1
1	A	1868	LYS	2.1
1	A	2283	PRO	2.0
1	A	1884	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	A	2402	4/4	0.81	0.20	36,40,44,47	0
2	EDO	A	2405	4/4	0.85	0.16	44,46,46,47	0
2	EDO	A	2406	4/4	0.85	0.21	62,63,63,63	0
2	EDO	A	2403	4/4	0.93	0.09	38,40,45,49	0
2	EDO	A	2404	4/4	0.95	0.09	24,29,30,34	0
2	EDO	A	2407	4/4	0.95	0.09	26,28,29,30	0
2	EDO	A	2401	4/4	0.96	0.08	21,22,22,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.