



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 10:31 PM UTC

PDB ID : 6EJD / pdb_00006ejd
Title : Human Xylosyltransferase 1 in complex with peptide QEEEGSGGPQGG
Authors : Briggs, D.C.; Hohenester, E.
Deposited on : 2017-09-20
Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	751	9% 88% • 8%
2	B	12	42% 67% 8% 25%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10788 atoms, of which 5255 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xylosyltransferase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	693	Total	C	H	N	O	S	0	0	0
			10696	3513	5223	951	977	32			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	209	ALA	-	expression tag	UNP Q86Y38
A	210	PRO	-	expression tag	UNP Q86Y38
A	211	LEU	-	expression tag	UNP Q86Y38
A	212	VAL	-	expression tag	UNP Q86Y38
A	213	HIS	-	expression tag	UNP Q86Y38
A	214	HIS	-	expression tag	UNP Q86Y38
A	215	HIS	-	expression tag	UNP Q86Y38
A	216	HIS	-	expression tag	UNP Q86Y38
A	217	HIS	-	expression tag	UNP Q86Y38
A	218	HIS	-	expression tag	UNP Q86Y38
A	219	ALA	-	expression tag	UNP Q86Y38
A	220	LEU	-	expression tag	UNP Q86Y38
A	221	ASP	-	expression tag	UNP Q86Y38
A	222	GLU	-	expression tag	UNP Q86Y38
A	223	ASN	-	expression tag	UNP Q86Y38
A	224	LEU	-	expression tag	UNP Q86Y38
A	225	TYR	-	expression tag	UNP Q86Y38
A	226	PHE	-	expression tag	UNP Q86Y38
A	227	GLN	-	expression tag	UNP Q86Y38
A	228	GLY	-	expression tag	UNP Q86Y38
A	229	ALA	-	expression tag	UNP Q86Y38
A	230	LEU	-	expression tag	UNP Q86Y38
A	231	ALA	-	expression tag	UNP Q86Y38


- Molecule 2 is a protein called Protein AMBP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	9	Total	C	H	N	O	0	0	0
			81	28	32	9	12			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	218	PRO	GLY	engineered mutation	UNP P02760
B	220	GLY	LEU	engineered mutation	UNP P02760
B	221	GLY	VAL	engineered mutation	UNP P02760

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O P	0	0
			5	4 1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total	O	0	0
			6	6		

- Molecule 1: Xylosyltransferase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.54Å 87.35Å 153.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.83 – 2.68 61.83 – 2.68	Depositor EDS
% Data completeness (in resolution range)	99.3 (61.83-2.68) 99.3 (61.83-2.68)	Depositor EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.69Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.244 , 0.279 0.244 , 0.280	Depositor DCC
R_{free} test set	1258 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	35.8	Xtriage
Anisotropy	0.778	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	10788	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.17	0/5634	0.41	0/7674
2	B	0.21	0/49	0.46	0/65
All	All	0.17	0/5683	0.41	0/7739

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5473	5223	5217	16	0
2	B	49	32	32	0	0
3	A	5	0	0	0	0
4	A	6	0	0	0	0
All	All	5533	5255	5249	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:864:ARG:O	1:A:866:ALA:N	2.24	0.70
1:A:576:PRO:O	1:A:749:LYS:NZ	2.34	0.55
1:A:953:LYS:NZ	1:A:959:ARG:OXT	2.46	0.48
1:A:864:ARG:C	1:A:866:ALA:H	2.23	0.47
1:A:449:LYS:HE3	1:A:547:ASP:OD1	2.15	0.47
1:A:350:TYR:HA	1:A:356:TYR:OH	2.15	0.46
1:A:864:ARG:C	1:A:866:ALA:N	2.73	0.46
1:A:927:CYS:HB3	1:A:930:MET:HE2	1.97	0.46
1:A:425:ALA:HB1	1:A:599:LYS:HG2	2.00	0.44
1:A:864:ARG:O	1:A:865:ASN:C	2.61	0.43
1:A:637:GLU:CG	1:A:638:PRO:HD3	2.49	0.42
1:A:876:ASN:HB2	1:A:877:PRO:HD3	2.02	0.42
1:A:517:MET:HE2	1:A:517:MET:HA	2.01	0.41
1:A:931:GLN:OE1	1:A:936:THR:HG22	2.20	0.41
1:A:722:LYS:NZ	1:A:911:GLY:O	2.53	0.41
1:A:637:GLU:HG2	1:A:638:PRO:HD3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	687/751 (92%)	673 (98%)	13 (2%)	1 (0%)	48	71
2	B	7/12 (58%)	6 (86%)	0	1 (14%)	0	0
All	All	694/763 (91%)	679 (98%)	13 (2%)	2 (0%)	36	57

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	865	ASN
2	B	218	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	570/656 (87%)	559 (98%)	11 (2%)	50	75
2	B	3/7 (43%)	3 (100%)	0	100	100
All	All	573/663 (86%)	562 (98%)	11 (2%)	50	75

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	265	ILE
1	A	321	MET
1	A	449	LYS
1	A	461	LYS
1	A	590	THR
1	A	727	ILE
1	A	786	ASP
1	A	797	ILE
1	A	846	ASN
1	A	848	GLN
1	A	892	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	343	GLN
1	A	585	HIS
1	A	687	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	1001	-	4,4,4	0.97	0	6,6,6	0.45	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	693/751 (92%)	0.84	65 (9%) 14 12	17, 39, 66, 89	0
2	B	9/12 (75%)	2.72	5 (55%) 0 0	47, 58, 68, 74	0
All	All	702/763 (92%)	0.86	70 (9%) 12 10	17, 39, 66, 89	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	733	ASP	5.4
1	A	864	ARG	4.9
2	B	218	PRO	4.7
2	B	219	GLN	4.5
1	A	829	TRP	4.4
2	B	212	GLU	4.3
1	A	863	LEU	4.2
1	A	734	PHE	3.9
1	A	561	CYS	3.6
1	A	453	ARG	3.6
1	A	865	ASN	3.6
1	A	563	CYS	3.6
1	A	511	ASP	3.4
2	B	211	GLU	3.4
1	A	727	ILE	3.4
1	A	870	GLN	3.4
1	A	866	ALA	3.3
1	A	304	GLU	3.2
1	A	308	ASN	3.2
1	A	590	THR	3.1
2	B	217	GLY	3.1
1	A	776	PRO	3.1
1	A	869	GLU	3.1
1	A	271	ALA	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	257	CYS	3.0
1	A	558	LYS	2.9
1	A	471	CYS	2.9
1	A	858	LEU	2.8
1	A	879	LEU	2.8
1	A	811	PRO	2.7
1	A	557	ARG	2.7
1	A	892	ARG	2.7
1	A	873	GLN	2.6
1	A	831	PRO	2.6
1	A	319	GLU	2.5
1	A	876	ASN	2.5
1	A	277	ARG	2.5
1	A	883	ILE	2.5
1	A	775	GLY	2.5
1	A	285	CYS	2.4
1	A	253	GLN	2.4
1	A	822	THR	2.4
1	A	923	GLY	2.4
1	A	886	ALA	2.4
1	A	854	GLU	2.4
1	A	878	VAL	2.4
1	A	893	ARG	2.4
1	A	832	VAL	2.3
1	A	794	THR	2.3
1	A	454	ASP	2.3
1	A	306	LYS	2.3
1	A	423	SER	2.3
1	A	565	TYR	2.2
1	A	478	LEU	2.2
1	A	489	VAL	2.2
1	A	295	GLU	2.2
1	A	877	PRO	2.2
1	A	452	GLY	2.2
1	A	868	MET	2.2
1	A	297	VAL	2.2
1	A	874	SER	2.2
1	A	756	PHE	2.2
1	A	572	CYS	2.1
1	A	897	SER	2.1
1	A	547	ASP	2.1
1	A	457	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	670	ASP	2.0
1	A	275	HIS	2.0
1	A	276	CYS	2.0
1	A	585	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PO4	A	1001	5/5	0.76	0.28	109,110,116,120	0

6.5 Other polymers [i](#)

There are no such residues in this entry.