



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:22 PM UTC

PDB ID : 2ELA / pdb_00002ela
Title : Crystal Structure of the PTB domain of human APPL1
Authors : Li, J.; Mao, X.; Dong, L.Q.; Liu, F.; Tong, L.
Deposited on : 2007-03-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	175	8% 50% 23% • 25%
1	B	175	8% 53% 23% • 19%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2418 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adapter protein containing PH domain, PTB domain and leucine zipper motif 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	131	Total	C	N	O	S	0	0	0
			1048	667	182	191	8			
1	B	141	Total	C	N	O	S	0	0	0
			1134	716	201	209	8			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	472	MET	-	cloning artifact	UNP Q9UKG1
A	473	GLY	-	cloning artifact	UNP Q9UKG1
A	474	SER	-	cloning artifact	UNP Q9UKG1
A	475	SER	-	cloning artifact	UNP Q9UKG1
A	476	HIS	-	cloning artifact	UNP Q9UKG1
A	477	HIS	-	cloning artifact	UNP Q9UKG1
A	478	HIS	-	cloning artifact	UNP Q9UKG1
A	479	HIS	-	cloning artifact	UNP Q9UKG1
A	480	HIS	-	cloning artifact	UNP Q9UKG1
A	481	HIS	-	cloning artifact	UNP Q9UKG1
A	482	SER	-	cloning artifact	UNP Q9UKG1
A	483	SER	-	cloning artifact	UNP Q9UKG1
A	484	GLY	-	cloning artifact	UNP Q9UKG1
A	485	LEU	-	cloning artifact	UNP Q9UKG1
A	486	VAL	-	cloning artifact	UNP Q9UKG1
A	487	PRO	-	cloning artifact	UNP Q9UKG1
A	488	ARG	-	cloning artifact	UNP Q9UKG1
A	489	GLY	-	cloning artifact	UNP Q9UKG1
A	490	SER	-	cloning artifact	UNP Q9UKG1
A	491	HIS	-	cloning artifact	UNP Q9UKG1
A	492	MET	-	cloning artifact	UNP Q9UKG1
B	472	MET	-	cloning artifact	UNP Q9UKG1
B	473	GLY	-	cloning artifact	UNP Q9UKG1
B	474	SER	-	cloning artifact	UNP Q9UKG1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	475	SER	-	cloning artifact	UNP Q9UKG1
B	476	HIS	-	cloning artifact	UNP Q9UKG1
B	477	HIS	-	cloning artifact	UNP Q9UKG1
B	478	HIS	-	cloning artifact	UNP Q9UKG1
B	479	HIS	-	cloning artifact	UNP Q9UKG1
B	480	HIS	-	cloning artifact	UNP Q9UKG1
B	481	HIS	-	cloning artifact	UNP Q9UKG1
B	482	SER	-	cloning artifact	UNP Q9UKG1
B	483	SER	-	cloning artifact	UNP Q9UKG1
B	484	GLY	-	cloning artifact	UNP Q9UKG1
B	485	LEU	-	cloning artifact	UNP Q9UKG1
B	486	VAL	-	cloning artifact	UNP Q9UKG1
B	487	PRO	-	cloning artifact	UNP Q9UKG1
B	488	ARG	-	cloning artifact	UNP Q9UKG1
B	489	GLY	-	cloning artifact	UNP Q9UKG1
B	490	SER	-	cloning artifact	UNP Q9UKG1
B	491	HIS	-	cloning artifact	UNP Q9UKG1
B	492	MET	-	cloning artifact	UNP Q9UKG1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	108	Total O 108 108	0	0
2	B	128	Total O 128 128	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adapter protein containing PH domain, PTB domain and leucine zipper motif 1



- Molecule 1: Adapter protein containing PH domain, PTB domain and leucine zipper motif 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.53Å 61.59Å 60.74Å 90.00° 101.22° 90.00°	Depositor
Resolution (Å)	29.79 – 2.00 29.79 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.3 (29.79-2.00) 95.3 (29.79-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 1.88Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.222 , 0.259 0.222 , 0.259	Depositor DCC
R_{free} test set	1859 reflections (7.37%)	wwPDB-VP
Wilson B-factor (Å ²)	17.5	Xtriage
Anisotropy	0.643	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2418	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1066	0.87	2/1441 (0.1%)
1	B	0.42	0/1152	0.92	6/1553 (0.4%)
All	All	0.42	0/2218	0.90	8/2994 (0.3%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	517	HIS	CA-C-N	8.21	128.31	119.28
1	B	517	HIS	C-N-CA	8.21	128.31	119.28
1	B	513	LYS	N-CA-C	-6.69	104.07	111.36
1	B	551	CYS	N-CA-C	6.17	116.58	107.88
1	B	576	THR	N-CA-C	-6.11	100.67	110.14
1	A	576	THR	N-CA-C	-6.03	100.79	110.14
1	A	551	CYS	N-CA-C	5.97	116.30	107.88
1	B	515	ASP	N-CA-C	-5.88	100.61	109.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1048	0	1060	38	0
1	B	1134	0	1146	48	0
2	A	108	0	0	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	128	0	0	14	0
All	All	2418	0	2206	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:498:LEU:HD11	1:A:500:GLN:HE21	1.27	0.97
1:A:521:VAL:HG22	2:A:161:HOH:O	1.74	0.86
1:B:624:GLN:HG2	2:B:146:HOH:O	1.78	0.84
1:A:505:ARG:HH22	1:A:609:ASN:ND2	1.77	0.82
1:B:541:THR:HG21	1:B:563:ARG:HH22	1.47	0.79
1:A:565:THR:HG21	1:B:511:GLU:OE2	1.84	0.77
1:B:512:VAL:HG22	2:B:149:HOH:O	1.88	0.74
1:A:553:LYS:HG2	1:A:565:THR:HG22	1.70	0.73
1:A:505:ARG:HH22	1:A:609:ASN:HD22	1.38	0.71
1:B:505:ARG:HH22	1:B:609:ASN:ND2	1.86	0.70
1:A:590:THR:HB	2:A:128:HOH:O	1.89	0.70
1:A:600:SER:HB3	2:A:128:HOH:O	1.92	0.68
1:B:602:VAL:HA	2:B:149:HOH:O	1.94	0.68
1:A:539:ARG:NH1	1:B:539:ARG:HG3	2.13	0.64
1:B:645:VAL:HG12	2:B:132:HOH:O	1.99	0.62
1:A:554:LEU:HD23	1:A:566:PHE:HE1	1.65	0.62
1:B:512:VAL:HG23	1:B:514:SER:HB2	1.83	0.60
1:B:644:ARG:HB3	2:B:107:HOH:O	2.02	0.60
1:B:538:PHE:H	1:B:539:ARG:NH1	2.00	0.59
1:B:555:ILE:HD13	1:B:562:THR:HA	1.83	0.59
1:A:517:HIS:ND1	1:A:518:PRO:HD2	2.19	0.58
1:B:541:THR:CG2	1:B:563:ARG:HH22	2.16	0.57
1:B:538:PHE:H	1:B:539:ARG:HH12	1.53	0.56
1:B:637:GLU:HB2	2:B:129:HOH:O	2.06	0.56
1:B:582:ARG:HD2	2:B:11:HOH:O	2.06	0.56
1:B:499:HIS:HB2	1:B:546:LEU:HD11	1.88	0.55
1:B:545:LEU:HD23	1:B:546:LEU:N	2.22	0.55
1:A:511:GLU:HB3	1:B:562:THR:HG23	1.89	0.54
1:B:638:LYS:O	1:B:642:ILE:HG12	2.09	0.53
1:B:499:HIS:CG	1:B:546:LEU:HD11	2.44	0.53
1:B:556:ASP:HB2	1:B:563:ARG:HD2	1.92	0.52
1:A:546:LEU:CD1	1:A:548:THR:HG23	2.40	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:505:ARG:HH22	1:B:609:ASN:HD22	1.55	0.51
1:B:599:LEU:HD23	1:B:599:LEU:N	2.26	0.51
1:B:588:LEU:HD12	2:B:48:HOH:O	2.11	0.51
1:A:517:HIS:HD2	1:A:519:ASP:CG	2.18	0.50
1:A:552:LEU:HD13	1:A:568:LEU:HD13	1.94	0.49
1:B:639:GLN:O	1:B:643:GLU:HG3	2.13	0.48
1:A:588:LEU:HD23	2:A:3:HOH:O	2.12	0.48
1:A:497:ILE:HG22	1:A:498:LEU:N	2.29	0.47
1:A:517:HIS:CE1	1:A:518:PRO:HD2	2.50	0.47
1:A:539:ARG:HD3	2:A:130:HOH:O	2.14	0.47
1:A:565:THR:HG21	1:B:511:GLU:CD	2.41	0.46
1:A:554:LEU:O	1:A:563:ARG:HG2	2.15	0.46
1:A:501:LEU:HD22	1:A:546:LEU:HD23	1.97	0.46
1:B:628:HIS:HD2	2:B:213:HOH:O	1.99	0.46
1:A:568:LEU:N	1:A:569:PRO:HD2	2.31	0.45
1:B:510:MET:HE1	1:B:527:GLN:OE1	2.16	0.45
1:A:578:GLN:HG2	2:A:31:HOH:O	2.17	0.45
1:B:571:VAL:O	1:B:623:LYS:HE3	2.16	0.45
1:A:582:ARG:CZ	1:A:610:ASN:HB2	2.47	0.45
1:B:515:ASP:HB2	2:B:133:HOH:O	2.17	0.45
1:B:539:ARG:H	1:B:539:ARG:HH11	1.65	0.45
1:A:539:ARG:NH1	2:A:42:HOH:O	2.43	0.45
1:B:543:SER:HA	1:B:557:PRO:HD3	1.99	0.45
1:B:559:THR:O	1:B:560:GLN:HB2	2.17	0.44
1:B:515:ASP:OD1	1:B:516:ASP:N	2.42	0.44
1:A:587:VAL:HG23	2:A:161:HOH:O	2.18	0.44
1:B:517:HIS:HA	1:B:518:PRO:HD3	1.76	0.44
1:A:499:HIS:CE1	1:A:501:LEU:HD21	2.53	0.43
1:A:517:HIS:CG	1:A:518:PRO:HD2	2.53	0.43
1:B:645:VAL:HG11	2:B:199:HOH:O	2.17	0.43
1:A:515:ASP:O	1:A:516:ASP:HB3	2.19	0.43
1:A:559:THR:O	1:A:560:GLN:HB2	2.17	0.43
1:A:586:PHE:C	2:A:161:HOH:O	2.61	0.43
1:A:513:LYS:HE2	2:A:173:HOH:O	2.16	0.43
1:A:539:ARG:CZ	1:B:539:ARG:HG3	2.49	0.43
1:A:552:LEU:HD22	1:A:571:VAL:HG21	2.01	0.42
1:A:546:LEU:HD13	1:A:548:THR:HG23	2.00	0.42
1:B:536:ASN:ND2	1:B:538:PHE:HE1	2.18	0.42
1:A:498:LEU:HD23	1:A:622:ALA:HB2	2.02	0.42
1:B:644:ARG:HH11	1:B:644:ARG:HG3	1.84	0.42
1:B:582:ARG:O	1:B:608:SER:N	2.50	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:614:LYS:HE3	2:B:236:HOH:O	2.20	0.41
1:B:512:VAL:HG13	2:B:149:HOH:O	2.21	0.41
1:B:568:LEU:HD11	1:B:622:ALA:HB1	2.03	0.41
1:B:512:VAL:N	2:B:149:HOH:O	2.52	0.40
1:B:545:LEU:C	1:B:545:LEU:CD2	2.95	0.40
1:B:568:LEU:HD22	1:B:623:LYS:HE2	2.03	0.40
1:A:511:GLU:HB3	1:B:562:THR:CG2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/175 (73%)	120 (94%)	5 (4%)	2 (2%)	7	3
1	B	137/175 (78%)	133 (97%)	3 (2%)	1 (1%)	18	14
All	All	264/350 (75%)	253 (96%)	8 (3%)	3 (1%)	11	7

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	516	ASP
1	A	515	ASP
1	B	514	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	120/159 (76%)	117 (98%)	3 (2%)	42	45
1	B	129/159 (81%)	125 (97%)	4 (3%)	35	37
All	All	249/318 (78%)	242 (97%)	7 (3%)	38	41

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	536	ASN
1	A	621	LEU
1	A	633	ARG
1	B	539	ARG
1	B	545	LEU
1	B	568	LEU
1	B	599	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	500	GLN
1	A	517	HIS
1	A	535	HIS
1	A	536	ASN
1	A	609	ASN
1	A	610	ASN
1	A	624	GLN
1	B	499	HIS
1	B	517	HIS
1	B	535	HIS
1	B	536	ASN
1	B	544	HIS
1	B	558	GLN
1	B	560	GLN
1	B	609	ASN
1	B	628	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	131/175 (74%)	0.38	14 (10%) 11 10	13, 21, 44, 53	0
1	B	141/175 (80%)	0.59	14 (9%) 13 11	13, 23, 41, 50	0
All	All	272/350 (77%)	0.49	28 (10%) 12 11	13, 21, 43, 53	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	645	VAL	6.3
1	B	516	ASP	4.6
1	B	514	SER	4.3
1	B	599	LEU	4.2
1	A	633	ARG	4.2
1	B	591	SER	3.8
1	B	515	ASP	3.7
1	A	516	ASP	3.5
1	B	592	SER	3.4
1	A	497	ILE	3.0
1	A	515	ASP	3.0
1	A	514	SER	3.0
1	B	641	GLU	3.0
1	A	500	GLN	2.9
1	A	517	HIS	2.9
1	B	642	ILE	2.8
1	B	644	ARG	2.7
1	B	590	THR	2.6
1	A	591	SER	2.6
1	A	558	GLN	2.6
1	A	590	THR	2.4
1	B	499	HIS	2.3
1	A	582	ARG	2.3
1	B	637	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	578	GLN	2.1
1	A	599	LEU	2.0
1	A	498	LEU	2.0
1	B	638	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.