



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 09:07 PM UTC

PDB ID : 4EL4 / pdb_00004el4
Title : Crystal structure of the catalytic domain of botulinum neurotoxin BoNT/A C134S/C165S double mutant
Authors : Stura, E.A.; Vera, L.; Ptchelkine, D.; Dive, V.
Deposited on : 2012-04-10
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1216 (1.20-1.20)
Clashscore	190562	1265 (1.20-1.20)
Ramachandran outliers	187476	1226 (1.20-1.20)
Sidechain outliers	187428	1226 (1.20-1.20)
RSRZ outliers	180081	1214 (1.20-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	15% 78% 13% ...

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	IMD	A	504	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4195 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Botulinum neurotoxin A light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	425	Total	C	N	O	S	0	19	0
			3603	2315	592	689	7			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP P10845
A	-18	GLY	-	expression tag	UNP P10845
A	-17	SER	-	expression tag	UNP P10845
A	-16	SER	-	expression tag	UNP P10845
A	-15	HIS	-	expression tag	UNP P10845
A	-14	HIS	-	expression tag	UNP P10845
A	-13	HIS	-	expression tag	UNP P10845
A	-12	HIS	-	expression tag	UNP P10845
A	-11	HIS	-	expression tag	UNP P10845
A	-10	HIS	-	expression tag	UNP P10845
A	-9	SER	-	expression tag	UNP P10845
A	-8	SER	-	expression tag	UNP P10845
A	-7	GLY	-	expression tag	UNP P10845
A	-6	LEU	-	expression tag	UNP P10845
A	-5	VAL	-	expression tag	UNP P10845
A	-4	PRO	-	expression tag	UNP P10845
A	-3	ARG	-	expression tag	UNP P10845
A	-2	GLY	-	expression tag	UNP P10845
A	-1	SER	-	expression tag	UNP P10845
A	0	HIS	-	expression tag	UNP P10845
A	27	ALA	VAL	variant	UNP P10845
A	134	SER	CYS	engineered mutation	UNP P10845
A	165	SER	CYS	engineered mutation	UNP P10845

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is IMIDAZOLE (CCD ID: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			5	3	2		
4	A	1	Total	C	N	0	0
			5	3	2		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	565	Total	O	0	1
			565	565		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Botulinum neurotoxin A light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.30Å 66.30Å 64.90Å 90.00° 99.10° 90.00°	Depositor
Resolution (Å)	46.08 – 1.20 46.08 – 1.20	Depositor EDS
% Data completeness (in resolution range)	99.3 (46.08-1.20) 99.3 (46.08-1.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 1.20Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.167 , 0.202 0.168 , 0.203	Depositor DCC
R_{free} test set	6386 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	11.3	Xtriage
Anisotropy	0.179	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 56.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4195	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IMD, ZN, EDO, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	2/3688 (0.1%)	0.98	10/4987 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	132	THR	CA-C	6.24	1.61	1.52
1	A	174	ASN	CA-CB	5.88	1.60	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	GLU	N-CA-C	-10.16	101.10	113.19
1	A	62	PRO	N-CA-C	9.61	122.43	110.70
1	A	61	PRO	N-CA-C	7.67	120.05	110.70
1	A	25	PRO	CA-C-N	-6.62	112.07	123.25
1	A	25	PRO	C-N-CA	-6.62	112.07	123.25
1	A	61	PRO	CA-C-N	6.13	126.70	120.38
1	A	61	PRO	C-N-CA	6.13	126.70	120.38
1	A	310	LEU	O-C-N	-5.68	116.10	122.12
1	A	27	ALA	N-CA-C	-5.67	106.21	113.01
1	A	312	TYR	N-CA-C	-5.54	105.12	111.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3603	0	3512	104	0
2	A	1	0	0	0	0
3	A	12	0	15	1	0
4	A	10	0	10	8	0
5	A	4	0	6	0	0
6	A	565	0	0	19	0
All	All	4195	0	3543	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:PRO:HG2	1:A:62:PRO:HD3	1.14	1.07
1:A:238:ASN:H	4:A:504:IMD:H2	1.20	1.04
1:A:62:PRO:CB	1:A:63:PRO:HD3	1.91	0.99
1:A:62:PRO:HB2	1:A:63:PRO:CD	1.98	0.93
1:A:61:PRO:CG	1:A:62:PRO:HD3	1.98	0.93
1:A:238:ASN:H	4:A:504:IMD:C2	1.81	0.92
1:A:62:PRO:CB	1:A:63:PRO:CD	2.48	0.92
1:A:250:TYR:O	1:A:425:PHE:HD2	1.54	0.90
1:A:61:PRO:HG2	1:A:62:PRO:CD	2.02	0.88
1:A:423:PHE:HB3	1:A:425:PHE:CE1	2.10	0.86
1:A:62:PRO:HB3	1:A:63:PRO:HD3	1.57	0.84
1:A:238:ASN:HB2	4:A:504:IMD:H2	1.62	0.82
1:A:423:PHE:HB3	1:A:425:PHE:CD1	2.16	0.81
1:A:366:TYR:O	4:A:503:IMD:H5	1.81	0.81
1:A:67:GLN:O	1:A:68:VAL:CG1	2.30	0.80
1:A:425:PHE:CD1	1:A:425:PHE:N	2.51	0.79
1:A:171[B]:GLU:HG3	1:A:172:VAL:HG23	1.65	0.79
1:A:238:ASN:N	4:A:504:IMD:H2	1.98	0.77
1:A:162:GLN:HE22	1:A:164[A]:GLU:CD	1.94	0.75
1:A:67:GLN:O	1:A:68:VAL:HG12	1.87	0.74
1:A:67:GLN:C	1:A:68:VAL:HG12	2.12	0.74
1:A:63:PRO:C	1:A:65:ALA:H	1.94	0.74
1:A:68:VAL:CG2	6:A:1085:HOH:O	2.35	0.74
1:A:311[A]:GLN:H	1:A:311[A]:GLN:HE21	1.36	0.74
1:A:64:GLU:O	1:A:65:ALA:CB	2.38	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:PRO:HB2	1:A:63:PRO:HD2	1.72	0.70
1:A:238:ASN:CB	4:A:504:IMD:H2	2.22	0.70
1:A:67:GLN:C	1:A:69:PRO:N	2.46	0.69
1:A:61:PRO:N	1:A:62:PRO:HD2	2.07	0.69
1:A:318:LYS:HG3	1:A:319:GLU:N	2.08	0.68
1:A:250:TYR:O	1:A:425:PHE:CD2	2.43	0.68
1:A:282:PHE:CE1	4:A:504:IMD:H5	2.28	0.68
1:A:61:PRO:CG	1:A:62:PRO:CD	2.69	0.68
1:A:26:ASN:H	1:A:26:ASN:HD22	1.42	0.66
1:A:274:ILE:HA	6:A:973:HOH:O	1.94	0.66
1:A:67:GLN:O	1:A:69:PRO:N	2.30	0.64
1:A:282:PHE:CD1	4:A:504:IMD:H5	2.33	0.64
1:A:269[A]:HIS:ND1	6:A:875:HOH:O	2.30	0.63
1:A:68:VAL:N	1:A:69:PRO:CD	2.56	0.63
1:A:65:ALA:HB3	1:A:67:GLN:HG2	1.82	0.61
1:A:67:GLN:C	1:A:69:PRO:CD	2.74	0.61
1:A:356:LYS:NZ	6:A:1078:HOH:O	2.33	0.61
1:A:425:PHE:H	1:A:425:PHE:HD1	1.48	0.60
1:A:275:ASP:O	1:A:279[B]:GLU:HG3	2.02	0.60
1:A:241[A]:ARG:NH1	1:A:278:GLN:OE1	2.23	0.59
1:A:67:GLN:C	1:A:68:VAL:CG1	2.73	0.59
1:A:311[A]:GLN:H	1:A:311[A]:GLN:NE2	2.00	0.59
1:A:292:ASP:HB3	6:A:1165:HOH:O	2.03	0.59
1:A:123:ILE:HG22	1:A:125:THR:HG22	1.85	0.59
1:A:67:GLN:O	1:A:68:VAL:HG13	2.02	0.59
1:A:315:ASN:HB3	6:A:748:HOH:O	2.02	0.58
1:A:64:GLU:O	1:A:65:ALA:HB3	2.05	0.57
1:A:68:VAL:O	1:A:70:VAL:N	2.30	0.57
1:A:26:ASN:HD22	1:A:26:ASN:N	1.99	0.56
1:A:68:VAL:C	1:A:70:VAL:H	2.13	0.55
1:A:61:PRO:N	1:A:62:PRO:CD	2.70	0.55
1:A:70:VAL:HG21	1:A:370[B]:ASP:OD1	2.06	0.55
1:A:105:ARG:NH2	6:A:951:HOH:O	2.40	0.52
1:A:370[A]:ASP:CG	6:A:832:HOH:O	2.52	0.52
1:A:411:MET:HA	1:A:411:MET:HE2	1.92	0.51
1:A:63:PRO:C	1:A:65:ALA:N	2.59	0.51
1:A:254:SER:HB2	1:A:425:PHE:CE2	2.47	0.50
1:A:415:LYS:HB3	3:A:506:GOL:H32	1.94	0.50
1:A:279[B]:GLU:OE1	6:A:1107:HOH:O	2.20	0.50
1:A:326:ASP:OD2	1:A:326:ASP:C	2.55	0.50
1:A:194[A]:PHE:HD2	6:A:832:HOH:O	1.93	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:GLN:C	1:A:69:PRO:HD3	2.37	0.49
1:A:66:LYS:O	1:A:67:GLN:O	2.30	0.49
1:A:26:ASN:HA	6:A:1080:HOH:O	2.13	0.49
1:A:423:PHE:HB3	1:A:425:PHE:HE1	1.72	0.48
1:A:249:ALA:O	1:A:253:MET:HG3	2.13	0.48
1:A:23:LYS:HD2	1:A:30:MET:HG3	1.96	0.48
1:A:326:ASP:N	6:A:1052:HOH:O	2.46	0.48
1:A:285:TYR:CE2	1:A:289:LYS:HE3	2.49	0.47
1:A:68:VAL:C	1:A:70:VAL:N	2.73	0.46
1:A:315:ASN:O	1:A:319:GLU:HG3	2.15	0.46
1:A:336:LEU:O	1:A:340:LYS:HG3	2.16	0.46
1:A:278:GLN:HE21	1:A:278:GLN:HB2	1.62	0.45
1:A:292:ASP:CB	6:A:1165:HOH:O	2.63	0.45
1:A:68:VAL:HG22	6:A:1085:HOH:O	2.06	0.45
1:A:256:LEU:HD11	1:A:425:PHE:CE2	2.51	0.45
1:A:256:LEU:HD11	1:A:425:PHE:CZ	2.52	0.44
1:A:124:ASP:CG	6:A:880:HOH:O	2.61	0.44
1:A:307:THR:HG23	1:A:308:ALA:N	2.33	0.44
1:A:311[A]:GLN:HE21	1:A:311[A]:GLN:N	2.10	0.44
1:A:41:LYS:HE3	1:A:115:ILE:HD11	1.99	0.43
1:A:105:ARG:NH2	6:A:950:HOH:O	2.31	0.43
1:A:139:GLN:NE2	1:A:143:SER:OG	2.52	0.42
1:A:162:GLN:NE2	1:A:164[A]:GLU:OE2	2.43	0.42
1:A:162:GLN:HE22	1:A:164[A]:GLU:CG	2.32	0.42
1:A:170:HIS:CD2	1:A:173:LEU:H	2.38	0.42
1:A:64:GLU:O	1:A:65:ALA:HB2	2.16	0.42
1:A:23:LYS:HE2	1:A:30:MET:O	2.20	0.41
1:A:194[A]:PHE:CD2	6:A:832:HOH:O	2.57	0.41
1:A:330[B]:LYS:NZ	6:A:974:HOH:O	2.54	0.41
1:A:303:ILE:HD13	1:A:310:LEU:HA	2.03	0.41
1:A:381:LYS:NZ	6:A:629:HOH:O	2.53	0.41
1:A:119:GLY:HA3	1:A:180:TYR:CE2	2.56	0.40
1:A:126:GLU:HG2	1:A:302:SER:OG	2.21	0.40
1:A:67:GLN:HA	1:A:69:PRO:HD3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	442/445 (99%)	429 (97%)	8 (2%)	5 (1%)	11 1

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	62	PRO
1	A	65	ALA
1	A	67	GLN
1	A	68	VAL
1	A	69	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	399/397 (100%)	376 (94%)	23 (6%)	18 1

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	23	LYS
1	A	26	ASN
1	A	58	ASP
1	A	64	GLU
1	A	67	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	122	THR
1	A	124	ASP
1	A	171[A]	GLU
1	A	171[B]	GLU
1	A	173	LEU
1	A	277	LEU
1	A	278	GLN
1	A	301	LYS
1	A	303	ILE
1	A	311[A]	GLN
1	A	311[B]	GLN
1	A	318	LYS
1	A	330[A]	LYS
1	A	330[B]	LYS
1	A	340	LYS
1	A	424	GLU
1	A	425	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	A	40	ASN
1	A	139	GLN
1	A	170	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	IMD	A	504	-	5,5,5	0.54	0	5,5,5	0.81	0
4	IMD	A	503	-	5,5,5	0.83	0	5,5,5	0.52	0
3	GOL	A	506	-	5,5,5	0.18	0	5,5,5	0.61	0
3	GOL	A	502	-	5,5,5	1.11	1 (20%)	5,5,5	0.39	0
5	EDO	A	505	-	3,3,3	1.30	0	2,2,2	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IMD	A	504	-	-	-	0/1/1/1
4	IMD	A	503	-	-	-	0/1/1/1
3	GOL	A	506	-	-	2/4/4/4	-
3	GOL	A	502	-	-	2/4/4/4	-
5	EDO	A	505	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	GOL	O2-C2	-2.31	1.36	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	506	GOL	O1-C1-C2-C3
3	A	502	GOL	O1-C1-C2-O2
3	A	506	GOL	O1-C1-C2-O2
5	A	505	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	504	IMD	7	0
4	A	503	IMD	1	0
3	A	506	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	425/445 (95%)	0.65	67 (15%) 5 6	5, 15, 42, 71	19 (4%)

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	68	VAL	7.3
1	A	62	PRO	7.1
1	A	123	ILE	5.6
1	A	425	PHE	5.5
1	A	25	PRO	5.5
1	A	245	VAL	5.3
1	A	69	PRO	5.1
1	A	27	ALA	4.9
1	A	125	THR	4.6
1	A	309	SER	4.5
1	A	327	THR	4.4
1	A	122	THR	4.4
1	A	28	GLY	4.3
1	A	26	ASN	4.2
1	A	422	LEU	4.1
1	A	65	ALA	4.0
1	A	63	PRO	3.8
1	A	310	LEU	3.8
1	A	423	PHE	3.6
1	A	277	LEU	3.6
1	A	307	THR	3.6
1	A	67	GLN	3.5
1	A	61	PRO	3.5
1	A	329	GLY	3.5
1	A	303	ILE	3.4
1	A	304	VAL	3.4
1	A	276	SER	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	421	GLY	3.3
1	A	249	ALA	3.3
1	A	308	ALA	3.2
1	A	306	THR	3.2
1	A	131	ASP	3.1
1	A	326	ASP	3.0
1	A	330[A]	LYS	2.8
1	A	273	PHE	2.8
1	A	121	SER	2.7
1	A	58	ASP	2.7
1	A	256	LEU	2.7
1	A	323	LEU	2.7
1	A	250	TYR	2.6
1	A	173	LEU	2.6
1	A	60	ASN	2.6
1	A	305	GLY	2.5
1	A	246	ASN	2.5
1	A	252	GLU	2.5
1	A	312	TYR	2.5
1	A	275	ASP	2.5
1	A	124	ASP	2.5
1	A	1	MET	2.5
1	A	11[A]	LYS	2.3
1	A	240[A]	ASN	2.3
1	A	66	LYS	2.3
1	A	334	ASP	2.3
1	A	251	TYR	2.3
1	A	24	ILE	2.3
1	A	311[A]	GLN	2.2
1	A	293	ILE	2.2
1	A	274	ILE	2.2
1	A	300	ALA	2.2
1	A	70	VAL	2.1
1	A	254	SER	2.1
1	A	142	GLY	2.1
1	A	328	SER	2.1
1	A	279[A]	GLU	2.1
1	A	424	GLU	2.0
1	A	287	TYR	2.0
1	A	316	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	A	505	4/4	0.82	0.17	13,19,24,31	0
3	GOL	A	506	6/6	0.84	0.12	41,43,45,45	0
3	GOL	A	502	6/6	0.84	0.16	28,34,36,36	0
4	IMD	A	504	5/5	0.89	0.11	21,22,31,32	0
4	IMD	A	503	5/5	0.90	0.11	23,24,29,29	0
2	ZN	A	501	1/1	1.00	0.02	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.