



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:29 PM UTC

PDB ID : 5ELD / pdb_00005eld
Title : Cholera toxin classical B-pentamer in complex with A Lewis-y
Authors : Heggelund, J.E.; Burschowsky, D.; Krengel, U.
Deposited on : 2015-11-04
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	103	7% 96% •
1	B	103	10% 93% 6% •
1	C	103	95% 5%
1	D	103	3% 98% •
1	E	103	% 94% 5% •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	5	20% 80%
2	G	5	20% 80%
2	H	5	60% 40%
2	I	5	100%

2 Entry composition [i](#)


There are 9 unique types of molecules in this entry. The entry contains 5418 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cholera enterotoxin B subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	103	Total	C	N	O	S	0	8	0
			857	544	147	159	7			
1	B	103	Total	C	N	O	S	0	12	0
			879	559	149	164	7			
1	C	103	Total	C	N	O	S	0	9	0
			864	549	145	163	7			
1	D	103	Total	C	N	O	S	0	5	0
			842	532	147	156	7			
1	E	103	Total	C	N	O	S	0	5	0
			841	533	146	156	6			

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	F	5	Total	C	N	O	0	5	0
			120	68	4	48			
2	G	5	Total	C	N	O	0	5	0
			120	68	4	48			
2	H	5	Total	C	N	O	0	5	0
			120	68	4	48			
2	I	5	Total	C	N	O	0	0	0
			60	34	2	24			

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is beta-D-galactopyranose (CCD ID: GAL) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		
4	C	1	Total	C	O	0	0
			12	6	6		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	C	O	0	1
			12	6	6		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Na	0	0
			1	1		

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	C	1	Total	C	N	O	S	6	0
			12	6	1	4	1		

- Molecule 7 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			10	6	4		
7	D	1	Total	C	O	0	0
			10	6	4		
7	E	1	Total	C	O	0	0
			10	6	4		

- Molecule 8 is alpha-D-galactopyranose (CCD ID: GLA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	E	1	Total	C	O	0	1
			12	6	6		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	122	Total 122	O 122	0	0
9	B	114	Total 114	O 114	0	0
9	C	127	Total 127	O 127	0	0
9	D	128	Total 128	O 128	0	0
9	E	119	Total 119	O 119	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cholera enterotoxin B subunit



- Molecule 1: Cholera enterotoxin B subunit



- Molecule 1: Cholera enterotoxin B subunit



- Molecule 1: Cholera enterotoxin B subunit



- Molecule 1: Cholera enterotoxin B subunit



- Molecule 2: alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-alpha-D-glucopyran

ose

Chain F:  20% 80%

NDG1
GAL2
FUC3
A2G4
FUC5

• Molecule 2: alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-alpha-D-glucopyranose

Chain G:  20% 80%


NDG1
GAL2
FUC3
A2G4
FUC5

• Molecule 2: alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-alpha-D-glucopyranose

Chain H:  60% 40%

NDG1
GAL2
FUC3
A2G4
FUC5

• Molecule 2: alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-alpha-D-glucopyranose

Chain I:  100%

NDG1
GAL2
FUC3
A2G4
FUC5

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.70Å 69.81Å 134.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.48 – 1.40 48.48 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.1 (48.48-1.40) 99.3 (48.48-1.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.171 , 0.193 0.171 , 0.193	Depositor DCC
R_{free} test set	6438 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	13.8	Xtriage
Anisotropy	0.224	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 29.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.024 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5418	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, A2G, FUC, GAL, NDG, MES, PGE, GLA, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	0/895	0.93	0/1205
1	B	0.94	0/923	0.96	1/1243 (0.1%)
1	C	0.79	0/907	0.83	0/1224
1	D	0.91	0/871	0.87	0/1172
1	E	0.91	0/870	0.92	0/1170
All	All	0.90	0/4466	0.91	1/6014 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	59	ASP	N-CA-C	6.95	118.93	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	857	0	894	3	0
1	B	879	0	924	5	0
1	C	864	0	883	3	0
1	D	842	0	870	2	0
1	E	841	0	874	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	120	0	100	0	0
2	G	120	0	100	6	0
2	H	120	0	100	1	0
2	I	60	0	50	0	0
3	A	7	0	10	0	0
3	D	7	0	10	0	0
4	A	12	0	12	0	0
4	C	12	0	12	1	0
4	E	12	0	12	0	0
5	B	1	0	0	0	0
6	C	12	0	9	0	0
7	C	10	0	14	0	0
7	D	10	0	14	0	0
7	E	10	0	14	0	0
8	E	12	0	12	0	0
9	A	122	0	0	0	0
9	B	114	0	0	0	0
9	C	127	0	0	1	0
9	D	128	0	0	0	0
9	E	119	0	0	1	0
All	All	5418	0	4914	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:3[A]:FUC:O2	2:G:4[A]:A2G:H3	1.62	0.96
1:D:67[B]:ARG:NH1	1:E:70:ASP:OD1	2.08	0.84
1:C:56[B]:GLN:NE2	4:C:203:GAL:O1	2.22	0.73
1:B:91[B]:LYS:HD2	1:B:91[B]:LYS:O	1.90	0.72
2:G:2[A]:GAL:H3	2:G:4[A]:A2G:C5	2.23	0.68
1:A:67[B]:ARG:NH1	1:B:70:ASP:OD1	2.26	0.68
2:G:3[A]:FUC:O2	2:G:4[A]:A2G:C3	2.42	0.66
1:E:81[A]:LYS:HG2	9:E:319:HOH:O	2.08	0.54
1:C:16:GLN:NE2	1:C:89:ASN:HD22	2.09	0.51
1:B:91[B]:LYS:HD2	1:B:91[B]:LYS:C	2.36	0.50
1:B:9[B]:CYS:SG	1:B:86:CYS:SG	3.08	0.48
2:G:2[A]:GAL:C3	2:G:4[A]:A2G:C5	2.93	0.46
1:A:74[B]:ILE:HD12	9:C:425:HOH:O	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:59:ASP:HA	1:E:62:LYS:HG2	1.99	0.45
2:G:3[A]:FUC:HO2	2:G:4[A]:A2G:H3	1.76	0.44
1:C:31:LEU:C	1:C:31:LEU:HD12	2.42	0.44
1:B:35:ARG:HA	1:B:37:MET:HE3	2.00	0.44
1:E:31:LEU:HD12	1:E:31:LEU:C	2.44	0.42
2:H:2[B]:GAL:H3	2:H:4[B]:A2G:C5	2.51	0.41
1:D:62[B]:LYS:HB2	1:D:62[B]:LYS:HE3	1.80	0.41
1:A:32:ALA:HB1	1:E:12:TYR:CZ	2.56	0.40
2:G:1[A]:NDG:H6C1	2:G:2[A]:GAL:C1	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/103 (106%)	107 (98%)	2 (2%)	0	100	100
1	B	113/103 (110%)	110 (97%)	3 (3%)	0	100	100
1	C	110/103 (107%)	109 (99%)	1 (1%)	0	100	100
1	D	106/103 (103%)	105 (99%)	1 (1%)	0	100	100
1	E	106/103 (103%)	104 (98%)	2 (2%)	0	100	100
All	All	544/515 (106%)	535 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/89 (109%)	95 (98%)	2 (2%)	47	15
1	B	101/89 (114%)	100 (99%)	1 (1%)	68	40
1	C	98/89 (110%)	96 (98%)	2 (2%)	48	17
1	D	94/89 (106%)	94 (100%)	0	100	100
1	E	94/89 (106%)	92 (98%)	2 (2%)	47	15
All	All	484/445 (109%)	477 (99%)	7 (1%)	70	28

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81[A]	LYS
1	A	81[B]	LYS
1	B	59	ASP
1	C	29[A]	GLU
1	C	29[B]	GLU
1	E	81[A]	LYS
1	E	81[B]	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
1	B	49	GLN
1	C	3	GLN
1	C	16	GLN
1	C	49	GLN
1	C	103	ASN
1	D	3	GLN
1	D	13	HIS
1	D	49	GLN
1	E	49	GLN
1	E	103	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

35 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDG	F	1[A]	2	15,15,15	0.91	0	21,21,21	1.16	3 (14%)
2	NDG	F	1[B]	2	15,15,15	0.69	0	21,21,21	1.25	2 (9%)
2	GAL	F	2[A]	2	11,11,12	1.04	1 (9%)	15,15,17	1.01	1 (6%)
2	GAL	F	2[B]	2	11,11,12	1.03	2 (18%)	15,15,17	0.77	0
2	FUC	F	3[A]	2	10,10,11	1.18	0	14,14,16	0.76	0
2	FUC	F	3[B]	2	10,10,11	1.04	1 (10%)	14,14,16	0.86	0
2	A2G	F	4[A]	2	14,14,15	1.24	1 (7%)	17,19,21	1.34	2 (11%)
2	A2G	F	4[B]	2	14,14,15	1.41	2 (14%)	17,19,21	1.37	3 (17%)
2	FUC	F	5[A]	2	10,10,11	0.91	0	14,14,16	0.59	0
2	FUC	F	5[B]	2	10,10,11	0.81	0	14,14,16	0.91	0
2	NDG	G	1[A]	2	15,15,15	0.72	0	21,21,21	1.34	3 (14%)
2	NDG	G	1[B]	2	15,15,15	0.70	0	21,21,21	1.17	2 (9%)
2	GAL	G	2[A]	2	11,11,12	0.40	0	15,15,17	0.94	0
2	GAL	G	2[B]	2	11,11,12	1.37	2 (18%)	15,15,17	0.91	0
2	FUC	G	3[A]	2	10,10,11	0.57	0	14,14,16	1.06	1 (7%)
2	FUC	G	3[B]	2	10,10,11	0.71	0	14,14,16	0.86	1 (7%)
2	A2G	G	4[A]	2	14,14,15	0.74	0	17,19,21	1.20	2 (11%)
2	A2G	G	4[B]	2	14,14,15	1.44	2 (14%)	17,19,21	1.35	2 (11%)
2	FUC	G	5[A]	2	10,10,11	0.59	0	14,14,16	0.77	0
2	FUC	G	5[B]	2	10,10,11	1.05	0	14,14,16	0.76	0
2	NDG	H	1[A]	2	15,15,15	1.16	3 (20%)	21,21,21	1.32	3 (14%)
2	NDG	H	1[B]	2	15,15,15	1.05	1 (6%)	21,21,21	1.38	3 (14%)
2	GAL	H	2[A]	2	11,11,12	0.72	0	15,15,17	1.45	3 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GAL	H	2[B]	2	11,11,12	0.70	0	15,15,17	0.72	0
2	FUC	H	3[A]	2	10,10,11	1.15	1 (10%)	14,14,16	0.86	0
2	FUC	H	3[B]	2	10,10,11	0.64	0	14,14,16	1.00	0
2	A2G	H	4[A]	2	14,14,15	1.04	1 (7%)	17,19,21	1.57	3 (17%)
2	A2G	H	4[B]	2	14,14,15	0.95	0	17,19,21	0.97	0
2	FUC	H	5[A]	2	10,10,11	1.05	0	14,14,16	0.78	0
2	FUC	H	5[B]	2	10,10,11	1.02	1 (10%)	14,14,16	1.14	1 (7%)
2	NDG	I	1	2	15,15,15	0.72	0	21,21,21	0.96	1 (4%)
2	GAL	I	2	2	11,11,12	0.84	0	15,15,17	1.32	3 (20%)
2	FUC	I	3	2	10,10,11	0.71	0	14,14,16	1.11	1 (7%)
2	A2G	I	4	2	14,14,15	1.44	3 (21%)	17,19,21	1.92	3 (17%)
2	FUC	I	5	2	10,10,11	1.08	1 (10%)	14,14,16	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDG	F	1[A]	2	-	0/6/26/26	0/1/1/1
2	NDG	F	1[B]	2	-	0/6/26/26	0/1/1/1
2	GAL	F	2[A]	2	-	0/2/19/22	0/1/1/1
2	GAL	F	2[B]	2	-	1/2/19/22	0/1/1/1
2	FUC	F	3[A]	2	-	-	0/1/1/1
2	FUC	F	3[B]	2	-	-	0/1/1/1
2	A2G	F	4[A]	2	-	2/6/23/26	0/1/1/1
2	A2G	F	4[B]	2	-	2/6/23/26	0/1/1/1
2	FUC	F	5[A]	2	-	-	0/1/1/1
2	FUC	F	5[B]	2	-	-	0/1/1/1
2	NDG	G	1[A]	2	-	3/6/26/26	0/1/1/1
2	NDG	G	1[B]	2	-	0/6/26/26	0/1/1/1
2	GAL	G	2[A]	2	-	1/2/19/22	0/1/1/1
2	GAL	G	2[B]	2	-	0/2/19/22	0/1/1/1
2	FUC	G	3[A]	2	-	-	0/1/1/1
2	FUC	G	3[B]	2	-	-	0/1/1/1
2	A2G	G	4[A]	2	-	0/6/23/26	0/1/1/1
2	A2G	G	4[B]	2	-	2/6/23/26	0/1/1/1
2	FUC	G	5[A]	2	-	-	0/1/1/1
2	FUC	G	5[B]	2	-	-	0/1/1/1
2	NDG	H	1[A]	2	-	0/6/26/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDG	H	1[B]	2	-	0/6/26/26	0/1/1/1
2	GAL	H	2[A]	2	-	1/2/19/22	0/1/1/1
2	GAL	H	2[B]	2	-	0/2/19/22	0/1/1/1
2	FUC	H	3[A]	2	-	-	0/1/1/1
2	FUC	H	3[B]	2	-	-	0/1/1/1
2	A2G	H	4[A]	2	-	0/6/23/26	0/1/1/1
2	A2G	H	4[B]	2	-	0/6/23/26	0/1/1/1
2	FUC	H	5[A]	2	-	-	0/1/1/1
2	FUC	H	5[B]	2	-	-	0/1/1/1
2	NDG	I	1	2	-	0/6/26/26	0/1/1/1
2	GAL	I	2	2	-	0/2/19/22	0/1/1/1
2	FUC	I	3	2	-	-	0/1/1/1
2	A2G	I	4	2	-	0/6/23/26	0/1/1/1
2	FUC	I	5	2	-	-	0/1/1/1

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	4[B]	A2G	C1-C2	3.51	1.57	1.52
2	I	4	A2G	C1-C2	2.95	1.56	1.52
2	F	4[B]	A2G	O4-C4	2.93	1.50	1.43
2	I	4	A2G	O5-C5	2.87	1.49	1.43
2	H	1[B]	NDG	C1-C2	-2.71	1.49	1.52
2	I	5	FUC	O5-C1	-2.71	1.39	1.43
2	G	4[B]	A2G	O5-C5	2.66	1.48	1.43
2	F	4[A]	A2G	C1-C2	-2.64	1.48	1.52
2	F	4[B]	A2G	C4-C5	2.58	1.58	1.53
2	H	1[A]	NDG	C2-N2	2.35	1.49	1.45
2	G	2[B]	GAL	O2-C2	2.32	1.48	1.43
2	F	2[A]	GAL	O5-C1	-2.31	1.39	1.43
2	G	2[B]	GAL	O5-C1	-2.31	1.39	1.43
2	H	4[A]	A2G	C2-N2	2.24	1.50	1.46
2	F	2[B]	GAL	C2-C3	2.23	1.55	1.52
2	H	1[A]	NDG	O5-C1	2.21	1.48	1.42
2	H	5[B]	FUC	O4-C4	2.19	1.48	1.43
2	F	3[B]	FUC	O5-C1	2.18	1.47	1.43
2	I	4	A2G	C8-C7	2.15	1.55	1.50
2	H	3[A]	FUC	O2-C2	2.10	1.47	1.43
2	F	2[B]	GAL	O5-C1	-2.03	1.40	1.43
2	H	1[A]	NDG	O5-C5	-2.02	1.39	1.44

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	4	A2G	O5-C5-C6	5.86	119.07	107.66
2	H	2[A]	GAL	O5-C5-C6	-3.54	100.77	107.66
2	H	1[B]	NDG	C1-O5-C5	-3.32	107.22	113.65
2	H	1[B]	NDG	O6-C6-C5	-3.11	100.73	111.33
2	G	1[A]	NDG	C1-O5-C5	-3.08	107.70	113.65
2	H	2[A]	GAL	C3-C4-C5	-2.93	104.92	110.23
2	H	1[A]	NDG	C1-O5-C5	2.88	119.23	113.65
2	H	4[A]	A2G	O5-C1-C2	-2.83	106.91	111.29
2	H	1[B]	NDG	C1-C2-N2	-2.79	107.49	110.73
2	G	4[A]	A2G	C4-C3-C2	-2.79	106.93	111.02
2	H	4[A]	A2G	C1-O5-C5	2.77	115.90	112.19
2	H	5[B]	FUC	C6-C5-C4	-2.71	108.12	113.08
2	I	4	A2G	C6-C5-C4	-2.68	106.43	113.02
2	I	3	FUC	C1-C2-C3	2.67	113.54	109.64
2	I	2	GAL	C1-O5-C5	2.63	115.71	112.19
2	F	4[B]	A2G	O5-C1-C2	-2.62	107.23	111.29
2	G	4[B]	A2G	C1-O5-C5	-2.60	108.70	112.19
2	F	4[A]	A2G	C2-N2-C7	-2.58	119.44	122.90
2	H	1[A]	NDG	O6-C6-C5	-2.55	102.65	111.33
2	F	1[B]	NDG	O1-C1-O5	-2.47	103.07	110.41
2	G	1[A]	NDG	O6-C6-C5	-2.44	103.02	111.33
2	F	1[B]	NDG	C1-C2-N2	-2.44	107.90	110.73
2	F	1[A]	NDG	O5-C5-C6	2.42	112.44	106.44
2	H	4[A]	A2G	O7-C7-N2	2.41	126.24	121.98
2	F	4[B]	A2G	O5-C5-C4	2.40	116.67	110.83
2	G	4[B]	A2G	C3-C4-C5	-2.39	105.90	110.23
2	G	1[A]	NDG	O5-C1-C2	-2.39	107.12	109.52
2	H	2[A]	GAL	O6-C6-C5	-2.39	103.21	111.33
2	F	4[B]	A2G	O4-C4-C5	2.38	115.18	109.32
2	F	2[A]	GAL	O3-C3-C4	2.37	115.96	110.38
2	I	2	GAL	O5-C5-C6	2.33	112.19	107.66
2	F	4[A]	A2G	C3-C4-C5	-2.21	106.23	110.23
2	G	3[A]	FUC	C2-C3-C4	-2.14	107.09	110.86
2	F	1[A]	NDG	C1-C2-N2	-2.14	108.25	110.73
2	G	1[B]	NDG	C1-C2-N2	-2.14	108.25	110.73
2	F	1[A]	NDG	O5-C1-C2	-2.14	107.37	109.52
2	I	2	GAL	O6-C6-C5	-2.13	104.08	111.33
2	G	3[B]	FUC	O5-C5-C4	2.12	113.36	109.55
2	H	1[A]	NDG	C1-C2-C3	2.08	113.38	110.54
2	G	1[B]	NDG	O7-C7-C8	-2.07	118.37	122.05
2	I	1	NDG	O7-C7-C8	-2.05	118.40	122.05
2	I	4	A2G	O6-C6-C5	-2.02	104.46	111.33
2	G	4[A]	A2G	O6-C6-C5	-2.01	104.49	111.33

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	4[B]	A2G	O5-C5-C6-O6
2	G	1[A]	NDG	O5-C5-C6-O6
2	G	4[B]	A2G	C4-C5-C6-O6
2	F	4[B]	A2G	C4-C5-C6-O6
2	G	1[A]	NDG	C4-C5-C6-O6
2	G	4[B]	A2G	O5-C5-C6-O6
2	F	4[A]	A2G	O5-C5-C6-O6
2	G	2[A]	GAL	O5-C5-C6-O6
2	H	2[A]	GAL	O5-C5-C6-O6
2	F	2[B]	GAL	O5-C5-C6-O6
2	F	4[A]	A2G	C4-C5-C6-O6
2	G	1[A]	NDG	C3-C2-N2-C7


There are no ring outliers.


6 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	3[A]	FUC	3	0
2	H	2[B]	GAL	1	0
2	G	2[A]	GAL	3	0
2	H	4[B]	A2G	1	0
2	G	1[A]	NDG	1	0
2	G	4[A]	A2G	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	PGE	C	202	-	9,9,9	0.42	0	8,8,8	0.32	0
3	PEG	D	201	-	6,6,6	0.50	0	5,5,5	0.32	0
7	PGE	D	202	-	9,9,9	0.53	0	8,8,8	0.36	0
7	PGE	E	201	-	9,9,9	0.42	0	8,8,8	0.39	0
4	GAL	A	202	-	12,12,12	1.02	1 (8%)	17,17,17	0.71	0
4	GAL	C	203	-	12,12,12	1.21	0	17,17,17	0.87	0
6	MES	C	201	-	12,12,12	6.21	2 (16%)	15,16,16	3.40	8 (53%)
4	GAL	E	202[A]	-	12,12,12	0.65	0	17,17,17	0.96	0
8	GLA	E	203[B]	-	12,12,12	1.11	1 (8%)	17,17,17	0.98	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PEG	A	201	-	6,6,6	0.39	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	PGE	C	202	-	-	1/7/7/7	-
3	PEG	D	201	-	-	0/4/4/4	-
7	PGE	D	202	-	-	1/7/7/7	-
7	PGE	E	201	-	-	0/7/7/7	-
4	GAL	A	202	-	-	0/2/22/22	0/1/1/1
4	GAL	C	203	-	-	0/2/22/22	0/1/1/1
6	MES	C	201	-	-	5/6/14/14	0/1/1/1
4	GAL	E	202[A]	-	-	0/2/22/22	0/1/1/1
8	GLA	E	203[B]	-	-	0/2/22/22	0/1/1/1
3	PEG	A	201	-	-	1/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	201	MES	C7-C8	-18.53	1.03	1.52
6	C	201	MES	C8-S	-10.64	1.62	1.77
4	A	202	GAL	O3-C3	2.36	1.48	1.43
8	E	203[B]	GLA	O2-C2	2.30	1.48	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	201	MES	O3S-S-C8	7.42	120.53	106.00
6	C	201	MES	C8-C7-N4	6.66	137.56	112.36
6	C	201	MES	C2-C3-N4	4.40	116.80	110.12
6	C	201	MES	O3S-S-O2S	-3.53	102.56	111.40
6	C	201	MES	C6-C5-N4	3.14	114.89	110.12
6	C	201	MES	O1-C6-C5	2.88	117.97	111.77
6	C	201	MES	C6-O1-C2	2.66	118.49	109.88
6	C	201	MES	O2S-S-O1S	-2.58	105.42	113.82
8	E	203[B]	GLA	C1-O5-C5	2.45	118.39	113.65

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	201	MES	C8-C7-N4-C5
6	C	201	MES	C7-C8-S-O2S
6	C	201	MES	C7-C8-S-O3S
3	A	201	PEG	O1-C1-C2-O2
7	C	202	PGE	O1-C1-C2-O2
6	C	201	MES	C7-C8-S-O1S
6	C	201	MES	C8-C7-N4-C3
7	D	202	PGE	C6-C5-O3-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	203	GAL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	103/103 (100%)	0.01	7 (6%) 23 24	5, 13, 34, 49	8 (7%)
1	B	103/103 (100%)	0.06	10 (9%) 13 13	6, 11, 48, 79	12 (11%)
1	C	103/103 (100%)	-0.19	0 100 100	6, 13, 22, 30	9 (8%)
1	D	103/103 (100%)	0.01	3 (2%) 53 56	7, 14, 30, 39	5 (4%)
1	E	103/103 (100%)	-0.06	1 (0%) 79 83	7, 15, 24, 35	5 (4%)
All	All	515/515 (100%)	-0.04	21 (4%) 41 43	5, 13, 31, 79	39 (7%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	58	ILE	7.3
1	B	57	HIS	5.9
1	B	54	GLY	4.9
1	B	60	SER	4.5
1	B	56	GLN	4.3
1	A	58	ILE	4.2
1	B	55	SER	3.7
1	B	53	PRO	3.3
1	D	103	ASN	3.2
1	D	58	ILE	3.2
1	B	103	ASN	3.2
1	B	52	VAL	3.1
1	A	57	HIS	2.7
1	B	59	ASP	2.7
1	A	55	SER	2.5
1	A	60	SER	2.4
1	A	59	ASP	2.3
1	A	54	GLY	2.2
1	E	103	ASN	2.2
1	A	56	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	55	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	A2G	G	4[A]	14/15	0.61	0.28	31,32,34,34	14
2	A2G	G	4[B]	14/15	0.61	0.28	36,39,45,45	14
2	A2G	I	4	14/15	0.71	0.21	39,42,56,57	14
2	GAL	G	2[B]	11/12	0.73	0.18	22,24,27,34	11
2	GAL	G	2[A]	11/12	0.73	0.18	24,27,27,30	11
2	FUC	G	3[B]	10/11	0.83	0.15	21,24,25,27	10
2	NDG	G	1[A]	15/15	0.83	0.14	21,22,23,24	15
2	NDG	G	1[B]	15/15	0.83	0.14	14,17,21,24	15
2	FUC	G	3[A]	10/11	0.83	0.15	22,22,23,23	10
2	FUC	G	5[B]	10/11	0.85	0.15	11,12,13,13	10
2	FUC	G	5[A]	10/11	0.85	0.15	17,19,19,19	10
2	FUC	I	3	10/11	0.88	0.12	24,27,30,33	10
2	GAL	I	2	11/12	0.90	0.11	25,29,31,34	11
2	NDG	I	1	15/15	0.91	0.10	18,22,28,30	15
2	A2G	F	4[A]	14/15	0.95	0.07	12,14,15,16	14
2	A2G	F	4[B]	14/15	0.95	0.07	13,16,17,18	14
2	NDG	H	1[A]	15/15	0.96	0.06	12,14,16,17	15
2	NDG	H	1[B]	15/15	0.96	0.06	18,20,22,23	15
2	NDG	F	1[A]	15/15	0.96	0.05	12,14,15,17	15
2	NDG	F	1[B]	15/15	0.96	0.05	15,16,19,19	15
2	FUC	F	3[A]	10/11	0.96	0.05	13,14,16,16	10
2	FUC	F	3[B]	10/11	0.96	0.05	16,17,17,19	10
2	FUC	I	5	10/11	0.96	0.08	16,17,19,20	10
2	A2G	H	4[B]	14/15	0.97	0.05	20,20,21,21	14
2	FUC	F	5[A]	10/11	0.97	0.05	11,12,12,12	10
2	FUC	F	5[B]	10/11	0.97	0.05	15,16,16,17	10
2	FUC	H	3[A]	10/11	0.97	0.05	10,10,11,11	10

Continued on next page...

Continued from previous page...



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FUC	H	3[B]	10/11	0.97	0.05	16,16,17,17	10
2	A2G	H	4[A]	14/15	0.97	0.05	12,13,14,14	14
2	FUC	H	5[B]	10/11	0.98	0.04	17,18,19,19	10
2	GAL	F	2[A]	11/12	0.98	0.05	12,13,14,14	11
2	GAL	F	2[B]	11/12	0.98	0.05	14,15,15,16	11
2	GAL	H	2[A]	11/12	0.98	0.04	10,12,12,14	11
2	GAL	H	2[B]	11/12	0.98	0.04	16,18,18,19	11
2	FUC	H	5[A]	10/11	0.98	0.04	12,13,13,13	10

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain G:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	PGE	C	202	10/10	0.81	0.16	35,40,44,48	0
7	PGE	D	202	10/10	0.81	0.16	36,40,44,44	0
3	PEG	D	201	7/7	0.82	0.16	44,44,49,53	0
4	GAL	E	202[A]	12/12	0.82	0.11	18,19,20,20	12
4	GAL	A	202	12/12	0.83	0.14	26,28,31,34	12
3	PEG	A	201	7/7	0.83	0.16	38,40,43,44	0
7	PGE	E	201	10/10	0.83	0.14	37,42,46,50	0
4	GAL	C	203	12/12	0.84	0.13	19,27,31,34	12
6	MES	C	201	12/12	0.86	0.13	24,27,64,65	6
8	GLA	E	203[B]	12/12	0.94	0.07	16,18,19,19	12
5	NA	B	201	1/1	0.99	0.04	11,11,11,11	1

6.5 Other polymers [i](#)

There are no such residues in this entry.