



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:38 AM UTC

PDB ID : 4EMB / pdb_00004emb
Title : Crystal structure of a phosphoglycerate mutase gpmA from *Borrelia burgdorferi* B31
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2012-04-11
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	
1	B	274	
1	C	274	
1	D	274	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8028 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2,3-bisphosphoglycerate-dependent phosphoglycerate mutase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	Se	0	1	0
			1928	1231	321	372	1	3			
1	B	241	Total	C	N	O	S	Se	0	0	0
			1895	1213	316	363	1	2			
1	C	244	Total	C	N	O	S	Se	0	0	0
			1905	1221	320	361	1	2			
1	D	241	Total	C	N	O	S	Se	0	1	0
			1882	1210	315	354	1	2			

There are 104 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MSE	-	expression tag	UNP O51602
A	-19	ALA	-	expression tag	UNP O51602
A	-18	HIS	-	expression tag	UNP O51602
A	-17	HIS	-	expression tag	UNP O51602
A	-16	HIS	-	expression tag	UNP O51602
A	-15	HIS	-	expression tag	UNP O51602
A	-14	HIS	-	expression tag	UNP O51602
A	-13	HIS	-	expression tag	UNP O51602
A	-12	MSE	-	expression tag	UNP O51602
A	-11	GLY	-	expression tag	UNP O51602
A	-10	THR	-	expression tag	UNP O51602
A	-9	LEU	-	expression tag	UNP O51602
A	-8	GLU	-	expression tag	UNP O51602
A	-7	ALA	-	expression tag	UNP O51602
A	-6	GLN	-	expression tag	UNP O51602
A	-5	THR	-	expression tag	UNP O51602
A	-4	GLN	-	expression tag	UNP O51602
A	-3	GLY	-	expression tag	UNP O51602
A	-2	PRO	-	expression tag	UNP O51602
A	-1	GLY	-	expression tag	UNP O51602
A	0	SER	-	expression tag	UNP O51602

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP O51602
A	2	LEU	-	expression tag	UNP O51602
A	3	GLY	-	expression tag	UNP O51602
A	4	ASP	-	expression tag	UNP O51602
A	5	PHE	-	expression tag	UNP O51602
B	-20	MSE	-	expression tag	UNP O51602
B	-19	ALA	-	expression tag	UNP O51602
B	-18	HIS	-	expression tag	UNP O51602
B	-17	HIS	-	expression tag	UNP O51602
B	-16	HIS	-	expression tag	UNP O51602
B	-15	HIS	-	expression tag	UNP O51602
B	-14	HIS	-	expression tag	UNP O51602
B	-13	HIS	-	expression tag	UNP O51602
B	-12	MSE	-	expression tag	UNP O51602
B	-11	GLY	-	expression tag	UNP O51602
B	-10	THR	-	expression tag	UNP O51602
B	-9	LEU	-	expression tag	UNP O51602
B	-8	GLU	-	expression tag	UNP O51602
B	-7	ALA	-	expression tag	UNP O51602
B	-6	GLN	-	expression tag	UNP O51602
B	-5	THR	-	expression tag	UNP O51602
B	-4	GLN	-	expression tag	UNP O51602
B	-3	GLY	-	expression tag	UNP O51602
B	-2	PRO	-	expression tag	UNP O51602
B	-1	GLY	-	expression tag	UNP O51602
B	0	SER	-	expression tag	UNP O51602
B	1	MSE	-	expression tag	UNP O51602
B	2	LEU	-	expression tag	UNP O51602
B	3	GLY	-	expression tag	UNP O51602
B	4	ASP	-	expression tag	UNP O51602
B	5	PHE	-	expression tag	UNP O51602
C	-20	MSE	-	expression tag	UNP O51602
C	-19	ALA	-	expression tag	UNP O51602
C	-18	HIS	-	expression tag	UNP O51602
C	-17	HIS	-	expression tag	UNP O51602
C	-16	HIS	-	expression tag	UNP O51602
C	-15	HIS	-	expression tag	UNP O51602
C	-14	HIS	-	expression tag	UNP O51602
C	-13	HIS	-	expression tag	UNP O51602
C	-12	MSE	-	expression tag	UNP O51602
C	-11	GLY	-	expression tag	UNP O51602
C	-10	THR	-	expression tag	UNP O51602

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-9	LEU	-	expression tag	UNP O51602
C	-8	GLU	-	expression tag	UNP O51602
C	-7	ALA	-	expression tag	UNP O51602
C	-6	GLN	-	expression tag	UNP O51602
C	-5	THR	-	expression tag	UNP O51602
C	-4	GLN	-	expression tag	UNP O51602
C	-3	GLY	-	expression tag	UNP O51602
C	-2	PRO	-	expression tag	UNP O51602
C	-1	GLY	-	expression tag	UNP O51602
C	0	SER	-	expression tag	UNP O51602
C	1	MSE	-	expression tag	UNP O51602
C	2	LEU	-	expression tag	UNP O51602
C	3	GLY	-	expression tag	UNP O51602
C	4	ASP	-	expression tag	UNP O51602
C	5	PHE	-	expression tag	UNP O51602
D	-20	MSE	-	expression tag	UNP O51602
D	-19	ALA	-	expression tag	UNP O51602
D	-18	HIS	-	expression tag	UNP O51602
D	-17	HIS	-	expression tag	UNP O51602
D	-16	HIS	-	expression tag	UNP O51602
D	-15	HIS	-	expression tag	UNP O51602
D	-14	HIS	-	expression tag	UNP O51602
D	-13	HIS	-	expression tag	UNP O51602
D	-12	MSE	-	expression tag	UNP O51602
D	-11	GLY	-	expression tag	UNP O51602
D	-10	THR	-	expression tag	UNP O51602
D	-9	LEU	-	expression tag	UNP O51602
D	-8	GLU	-	expression tag	UNP O51602
D	-7	ALA	-	expression tag	UNP O51602
D	-6	GLN	-	expression tag	UNP O51602
D	-5	THR	-	expression tag	UNP O51602
D	-4	GLN	-	expression tag	UNP O51602
D	-3	GLY	-	expression tag	UNP O51602
D	-2	PRO	-	expression tag	UNP O51602
D	-1	GLY	-	expression tag	UNP O51602
D	0	SER	-	expression tag	UNP O51602
D	1	MSE	-	expression tag	UNP O51602
D	2	LEU	-	expression tag	UNP O51602
D	3	GLY	-	expression tag	UNP O51602
D	4	ASP	-	expression tag	UNP O51602
D	5	PHE	-	expression tag	UNP O51602

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	112	Total	O	0	0
			112	112		
4	B	101	Total	O	0	0
			101	101		
4	C	105	Total	O	0	0
			105	105		
4	D	87	Total	O	0	0
			87	87		

- Molecule 1: 2,3-bisphosphoglycerate-dependent phosphoglycerate mutase





- Chain B: 3% 84% 12%



- Chain C:  3% 87% 11%



- Chain D:  5% 84% 12%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.32Å 116.75Å 84.41Å 90.00° 107.67° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 50.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-2.30) 100.0 (50.00-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.80 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.176 , 0.201 0.178 , 0.200	Depositor DCC
R_{free} test set	3216 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8028	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	0/1970	0.90	1/2674 (0.0%)
1	B	0.96	0/1934	0.88	0/2624
1	C	1.00	3/1944 (0.2%)	0.88	0/2639
1	D	0.97	1/1924 (0.1%)	0.89	0/2611
All	All	0.98	4/7772 (0.1%)	0.89	1/10548 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	14	HIS	ND1-CE1	5.51	1.38	1.32
1	C	14	HIS	CE1-NE2	5.51	1.38	1.32
1	D	94	HIS	CG-CD2	5.47	1.41	1.35
1	C	14	HIS	CG-CD2	5.05	1.41	1.35

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	122	TYR	N-CA-C	-5.78	106.36	113.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1928	0	1861	4	0
1	B	1895	0	1836	7	0
1	C	1905	0	1848	3	0
1	D	1882	0	1826	5	0
2	A	4	0	6	1	0
2	B	4	0	6	1	0
2	C	4	0	6	0	0
3	D	1	0	0	0	0
4	A	112	0	0	0	0
4	B	101	0	0	1	0
4	C	105	0	0	0	0
4	D	87	0	0	0	0
All	All	8028	0	7389	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:THR:HG22	1:B:115:VAL:CG2	2.14	0.76
1:D:106:THR:HG22	1:D:115:VAL:CG2	2.20	0.71
1:B:106:THR:HG22	1:B:115:VAL:HG21	1.76	0.65
1:C:106:THR:HG22	1:C:115:VAL:CG2	2.26	0.65
1:D:106:THR:HG22	1:D:115:VAL:HG21	1.77	0.64
1:B:203:GLU:OE2	4:B:496:HOH:O	2.16	0.61
1:A:106:THR:HG22	1:A:115:VAL:CG2	2.30	0.61
1:B:136:HIS:CE1	1:B:138:ILE:HG23	2.38	0.58
1:A:26:THR:H	2:A:301:EDO:HO1	1.52	0.56
1:C:106:THR:HG22	1:C:115:VAL:HG21	1.87	0.56
1:C:19:TRP:CH2	1:C:29:THR:HG21	2.41	0.54
1:A:136:HIS:CE1	1:A:138:ILE:HG23	2.44	0.53
1:D:136:HIS:CE1	1:D:138:ILE:HG23	2.45	0.51
1:D:19:TRP:CH2	1:D:29:THR:HG21	2.47	0.50
1:B:19:TRP:CH2	1:B:29:THR:HG21	2.48	0.48
1:B:106:THR:CG2	1:B:115:VAL:CG2	2.89	0.47
1:A:106:THR:HG22	1:A:115:VAL:HG21	1.96	0.47
1:B:27:GLY:H	2:B:301:EDO:HO1	1.62	0.42
1:D:160:THR:O	1:D:164:VAL:HG23	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/274 (89%)	240 (98%)	4 (2%)	0	100	100
1	B	239/274 (87%)	235 (98%)	4 (2%)	0	100	100
1	C	242/274 (88%)	238 (98%)	4 (2%)	0	100	100
1	D	240/274 (88%)	236 (98%)	4 (2%)	0	100	100
All	All	965/1096 (88%)	949 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/234 (87%)	200 (98%)	4 (2%)	48	67
1	B	200/234 (86%)	197 (98%)	3 (2%)	57	75
1	C	199/234 (85%)	197 (99%)	2 (1%)	68	82
1	D	195/234 (83%)	193 (99%)	2 (1%)	68	82
All	All	798/936 (85%)	787 (99%)	11 (1%)	59	76

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ASN
1	A	115	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	236	GLU
1	A	243	MSE
1	B	23	ASN
1	B	115	VAL
1	B	148	LYS
1	C	115	VAL
1	C	148	LYS
1	D	115	VAL
1	D	148	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	301	-	3,3,3	0.67	0	2,2,2	0.39	0
2	EDO	C	301	-	3,3,3	0.50	0	2,2,2	0.46	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	301	-	3,3,3	0.56	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	301	-	-	1/1/1/1	-
2	EDO	C	301	-	-	0/1/1/1	-
2	EDO	B	301	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	EDO	O1-C1-C2-O2
2	B	301	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	EDO	1	0
2	B	301	EDO	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	242/274 (88%)	-0.31	1 (0%) 88 89	20, 35, 62, 82	1 (0%)
1	B	238/274 (86%)	-0.18	7 (2%) 53 56	20, 35, 73, 88	0
1	C	241/274 (87%)	-0.24	7 (2%) 53 56	20, 35, 67, 95	0
1	D	238/274 (86%)	-0.10	13 (5%) 30 32	20, 37, 77, 92	1 (0%)
All	All	959/1096 (87%)	-0.21	28 (2%) 53 56	20, 35, 71, 95	2 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	116	LEU	3.8
1	D	116	LEU	3.6
1	C	244	GLU	3.5
1	C	248	SER	3.3
1	D	23	ASN	3.3
1	D	245	SER	3.1
1	D	244	GLU	3.1
1	D	235	ASP	2.9
1	D	113	ASP	2.9
1	D	21	LYS	2.7
1	D	242	ALA	2.7
1	C	246	VAL	2.6
1	C	239	ILE	2.6
1	B	113	ASP	2.6
1	D	22	GLU	2.5
1	C	245	SER	2.5
1	C	236	GLU	2.4
1	B	241	LYS	2.4
1	D	236	GLU	2.4
1	D	112	GLU	2.3
1	B	5	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	238	LYS	2.3
1	D	20	ASN	2.2
1	A	235	ASP	2.2
1	B	239	ILE	2.1
1	D	18	GLU	2.1
1	C	247	ALA	2.0
1	B	134	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	D	301	1/1	0.86	0.16	81,81,81,81	0
2	EDO	B	301	4/4	0.92	0.15	50,54,60,61	0
2	EDO	A	301	4/4	0.93	0.16	38,45,45,47	0
2	EDO	C	301	4/4	0.94	0.15	35,40,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.