



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:33 AM UTC

PDB ID : 1ENA / pdb_00001ena
Title : CRYSTAL STRUCTURES OF THE BINARY CA²⁺ AND PDTP COMPLEXES AND THE TERNARY COMPLEX OF THE ASP 21->GLU MUTANT OF STAPHYLOCOCCAL NUCLEASE. IMPLICATIONS FOR CATALYSIS AND LIGAND BINDING
Authors : Libson, A.; Gittis, A.; Lattman, E.E.
Deposited on : 1994-02-14
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	135	47% 38% 11% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1119 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called STAPHYLOCOCCAL NUCLEASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	135	Total	C	N	O	S	0	0	0
			1084	692	190	198	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	GLU	ASP	conflict	UNP P00644

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	34	Total	O	0	0
			34	34		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: STAPHYLOCOCCAL NUCLEASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	47.29Å 47.29Å 63.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.15	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.15)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ, X-PLOR	Depositor
R, R_{free}	0.225 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1119	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.40	9/1105 (0.8%)	2.28	60/1480 (4.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	121	HIS	CD2-NE2	-7.25	1.29	1.37
1	A	35	ARG	CZ-NH1	7.24	1.42	1.32
1	A	46	HIS	CD2-NE2	-6.96	1.30	1.37
1	A	124	HIS	CD2-NE2	-6.77	1.30	1.37
1	A	8	HIS	CD2-NE2	-6.16	1.31	1.37
1	A	36	LEU	CA-CB	-6.06	1.45	1.53
1	A	18	ILE	CA-CB	5.64	1.61	1.54
1	A	37	LEU	CB-CG	-5.38	1.42	1.53
1	A	87	ARG	CZ-NH1	5.09	1.39	1.32

All (60) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	87	ARG	NE-CZ-NH2	-10.98	109.32	119.20
1	A	123	GLN	OE1-CD-NE2	-10.59	112.01	122.60
1	A	68	ASN	OD1-CG-ND2	-9.31	113.29	122.60
1	A	131	GLN	OE1-CD-NE2	-9.11	113.49	122.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	123	GLN	CG-CD-NE2	8.58	129.27	116.40
1	A	46	HIS	CB-CG-CD2	-8.56	120.06	131.20
1	A	83	ASP	CA-CB-CG	7.88	120.48	112.60
1	A	87	ARG	NE-CZ-NH1	7.35	128.85	121.50
1	A	15	ILE	N-CA-C	-7.26	104.66	111.48
1	A	46	HIS	CB-CG-ND1	7.25	133.58	122.70
1	A	91	TYR	O-C-N	-7.12	115.23	123.27
1	A	131	GLN	CG-CD-NE2	7.07	127.01	116.40
1	A	128	SER	N-CA-C	6.99	118.97	111.36
1	A	12	ALA	CA-C-O	-6.94	113.84	121.33
1	A	121	HIS	CB-CG-CD2	-6.92	122.21	131.20
1	A	30	GLN	CA-C-N	6.88	127.26	119.83
1	A	30	GLN	C-N-CA	6.88	127.26	119.83
1	A	124	HIS	CB-CG-CD2	-6.85	122.30	131.20
1	A	35	ARG	NE-CZ-NH2	-6.69	113.18	119.20
1	A	112	ALA	N-CA-C	6.66	118.26	108.14
1	A	68	ASN	CB-CG-ND2	6.55	126.22	116.40
1	A	100	ASN	N-CA-C	-6.50	104.20	111.28
1	A	67	GLU	N-CA-C	6.46	118.40	111.36
1	A	16	LYS	O-C-N	-6.37	116.16	123.42
1	A	100	ASN	CA-CB-CG	6.27	118.87	112.60
1	A	116	LYS	O-C-N	-6.27	115.72	121.30
1	A	31	PRO	N-CA-CB	6.17	108.82	103.27
1	A	102	ALA	CA-C-N	6.15	128.52	120.28
1	A	102	ALA	C-N-CA	6.15	128.52	120.28
1	A	92	ILE	CB-CG1-CD1	6.11	126.64	113.80
1	A	115	TYR	N-CA-C	6.10	118.78	107.99
1	A	53	LYS	CA-CB-CG	6.04	126.18	114.10
1	A	8	HIS	CB-CG-CD2	-6.02	123.37	131.20
1	A	99	VAL	CB-CA-C	-6.00	104.04	112.14
1	A	80	GLN	CG-CD-NE2	-5.92	107.51	116.40
1	A	104	VAL	N-CA-C	-5.88	105.12	110.82
1	A	14	LEU	CA-CB-CG	5.87	136.86	116.30
1	A	133	LYS	N-CA-C	-5.87	104.79	111.07
1	A	126	ARG	NE-CZ-NH1	5.70	127.20	121.50
1	A	36	LEU	CA-C-O	5.68	127.38	120.62
1	A	83	ASP	CA-C-N	5.61	128.07	120.38
1	A	83	ASP	C-N-CA	5.61	128.07	120.38
1	A	111	VAL	N-CA-C	-5.61	102.34	109.58
1	A	12	ALA	O-C-N	-5.57	116.90	123.25
1	A	94	ALA	N-CA-C	-5.53	98.20	107.99
1	A	138	ASN	O-C-N	-5.53	115.23	122.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	81	ARG	NE-CZ-NH2	-5.48	114.27	119.20
1	A	23	VAL	CA-CB-CG1	5.42	119.61	110.40
1	A	40	ASP	CA-C-O	-5.42	114.50	120.30
1	A	30	GLN	OE1-CD-NE2	-5.38	117.22	122.60
1	A	41	THR	CA-C-N	5.35	125.65	119.92
1	A	41	THR	C-N-CA	5.35	125.65	119.92
1	A	92	ILE	N-CA-CB	-5.35	104.57	110.99
1	A	89	LEU	CA-C-O	-5.29	114.79	120.46
1	A	27	TYR	N-CA-C	-5.23	100.17	109.06
1	A	111	VAL	CA-C-O	-5.14	115.73	121.23
1	A	140	TRP	CG-CD1-NE1	-5.14	103.52	110.20
1	A	8	HIS	CB-CG-ND1	5.14	130.41	122.70
1	A	35	ARG	CG-CD-NE	5.10	123.22	112.00
1	A	126	ARG	N-CA-C	-5.09	105.92	111.82

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	91	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1084	0	1120	32	0
2	A	1	0	0	0	0
3	A	34	0	0	2	0
All	All	1119	0	1120	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:MET:SD	1:A:99:VAL:HG13	2.16	0.86
1:A:46:HIS:CG	1:A:49:LYS:HD2	2.13	0.83
1:A:38:LEU:HD22	1:A:114:VAL:HG12	1.59	0.83
1:A:46:HIS:CD2	1:A:49:LYS:HD2	2.14	0.82
1:A:44:THR:HA	1:A:51:VAL:HG13	1.68	0.74
1:A:130:ALA:O	1:A:134:LYS:HG3	1.95	0.66
1:A:123:GLN:OE1	1:A:123:GLN:HA	1.99	0.62
1:A:46:HIS:HB3	1:A:49:LYS:HB2	1.83	0.60
1:A:46:HIS:CE1	1:A:49:LYS:HD2	2.36	0.59
1:A:71:LYS:HG3	1:A:95:ASP:HA	1.86	0.56
1:A:49:LYS:HE2	3:A:175:HOH:O	2.05	0.55
1:A:83:ASP:HB3	1:A:89:LEU:HD21	1.89	0.55
1:A:19:ASP:HA	1:A:59:SER:HB3	1.89	0.54
1:A:46:HIS:CE1	1:A:49:LYS:NZ	2.76	0.53
1:A:77:ASP:HB3	1:A:118:ASN:OD1	2.09	0.53
1:A:122:GLU:O	1:A:126:ARG:HB2	2.11	0.51
1:A:35:ARG:HG2	3:A:153:HOH:O	2.09	0.51
1:A:46:HIS:ND1	1:A:49:LYS:HD2	2.28	0.49
1:A:46:HIS:NE2	1:A:49:LYS:HD2	2.28	0.47
1:A:46:HIS:CD2	1:A:49:LYS:CD	2.92	0.46
1:A:60:ALA:O	1:A:64:LYS:HE2	2.15	0.46
1:A:116:LYS:HD3	1:A:117:PRO:HA	1.98	0.45
1:A:13:THR:OG1	1:A:26:MET:HB3	2.15	0.45
1:A:81:ARG:CG	1:A:81:ARG:HH11	2.31	0.44
1:A:46:HIS:CG	1:A:49:LYS:HB2	2.52	0.44
1:A:116:LYS:HD3	1:A:117:PRO:CA	2.49	0.43
1:A:72:ILE:HD13	1:A:94:ALA:HA	2.01	0.42
1:A:46:HIS:HB3	1:A:49:LYS:CB	2.47	0.42
1:A:131:GLN:HA	1:A:134:LYS:HZ3	1.85	0.42
1:A:35:ARG:HD3	1:A:35:ARG:C	2.46	0.41
1:A:46:HIS:CE1	1:A:49:LYS:HZ3	2.39	0.40
1:A:70:LYS:HG2	1:A:95:ASP:OD1	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/135 (98%)	126 (95%)	5 (4%)	2 (2%)	8	4

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	49	LYS
1	A	138	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/114 (100%)	95 (83%)	19 (17%)	2	1

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	28	LYS
1	A	30	GLN
1	A	35	ARG
1	A	45	LYS
1	A	49	LYS
1	A	51	VAL
1	A	53	LYS
1	A	57	GLU
1	A	59	SER
1	A	70	LYS
1	A	81	ARG
1	A	84	LYS
1	A	92	ILE
1	A	106	GLN
1	A	116	LYS
1	A	126	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	131	GLN
1	A	133	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	HIS
1	A	121	HIS
1	A	131	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.