



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:32 AM UTC

PDB ID : 6EN3 / pdb_00006en3
Title : Crystal structure of full length EndoS from Streptococcus pyogenes in complex with G2 oligosaccharide.
Authors : Trastoy, B.; Klontz, E.H.; Orwenyo, J.; Marina, A.; Wang, L.X.; Sundberg, E.J.; Guerin, M.E.
Deposited on : 2017-10-04
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1191	 2% 74% 22%
2	B	8	 38% 50% 12%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7451 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endo-beta-N-acetylglucosaminidase F2,Multifunctional-autoprocessing repeats-in-toxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	934	Total	C	N	O	S	0	0	0
			7344	4649	1240	1438	17			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	36	MET	-	initiating methionine	UNP J7M8R4
A	233	ALA	ASP	engineered mutation	UNP J7M8R4
A	235	LEU	GLU	engineered mutation	UNP J7M8R4
A	996	LEU	-	linker	UNP J7M8R4
A	997	GLY	-	linker	UNP J7M8R4
A	998	SER	-	linker	UNP J7M8R4
A	1205	LEU	-	expression tag	UNP Q9KS12
A	1206	GLU	-	expression tag	UNP Q9KS12
A	1207	GLY	-	expression tag	UNP Q9KS12
A	1208	GLY	-	expression tag	UNP Q9KS12
A	1209	SER	-	expression tag	UNP Q9KS12
A	1210	GLY	-	expression tag	UNP Q9KS12
A	1211	GLY	-	expression tag	UNP Q9KS12
A	1212	SER	-	expression tag	UNP Q9KS12
A	1213	GLY	-	expression tag	UNP Q9KS12
A	1214	ASN	-	expression tag	UNP Q9KS12
A	1215	SER	-	expression tag	UNP Q9KS12
A	1216	GLY	-	expression tag	UNP Q9KS12
A	1217	HIS	-	expression tag	UNP Q9KS12
A	1218	HIS	-	expression tag	UNP Q9KS12
A	1219	HIS	-	expression tag	UNP Q9KS12
A	1220	HIS	-	expression tag	UNP Q9KS12
A	1221	HIS	-	expression tag	UNP Q9KS12
A	1222	HIS	-	expression tag	UNP Q9KS12
A	1223	HIS	-	expression tag	UNP Q9KS12
A	1224	HIS	-	expression tag	UNP Q9KS12

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1225	HIS	-	expression tag	UNP Q9KS12
A	1226	HIS	-	expression tag	UNP Q9KS12

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	8	Total	C	N	O	0	0	0
			98	54	3	41			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ni	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	7	Total	O	0	0
			7	7		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endo-beta-N-acetylglucosaminidase F2,Multifunctional-autoprocessing repeats-in-toxin



- Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.64Å 90.08Å 104.37Å 90.00° 110.44° 90.00°	Depositor
Resolution (Å)	45.04 – 2.90 45.04 – 2.90	Depositor EDS
% Data completeness (in resolution range)	98.8 (45.04-2.90) 99.5 (45.04-2.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 2.91Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.209 , 0.239 0.213 , 0.236	Depositor DCC
R_{free} test set	1492 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	68.6	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7451	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, CA, NAG, GAL, BMA, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/7488	0.33	0/10121

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7344	0	7268	29	0
2	B	98	0	84	5	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	7	0	0	0	0
All	All	7451	0	7352	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:ARG:NH1	1:A:182:THR:O	2.01	0.92
2:B:7:NAG:O6	2:B:8:GAL:O5	2.17	0.62
1:A:391:LYS:O	1:A:503:LYS:NZ	2.34	0.60
1:A:152:ASP:OD1	1:A:153:TRP:N	2.39	0.55
1:A:191:GLY:HA2	1:A:194:SER:OG	2.12	0.50
1:A:402:TYR:OH	2:B:2:BMA:O2	2.23	0.50
1:A:150:PHE:CE1	2:B:1:NAG:O7	2.64	0.50
2:B:7:NAG:C6	2:B:8:GAL:C1	2.89	0.50
1:A:143:GLU:N	1:A:143:GLU:OE1	2.45	0.49
1:A:192:ASP:HB2	1:A:241:LYS:O	2.11	0.49
1:A:806:LYS:O	1:A:807:GLN:NE2	2.45	0.49
1:A:186:ARG:NH2	2:B:6:MAN:O3	2.46	0.49
1:A:894:THR:HG21	1:A:928:ASP:HA	1.95	0.48
1:A:775:ILE:O	1:A:861:PRO:HB2	2.15	0.47
1:A:866:ASP:OD1	1:A:867:GLU:N	2.48	0.47
1:A:941:SER:HA	1:A:952:LEU:HD22	1.99	0.45
1:A:119:ARG:HD2	1:A:121:TRP:CZ2	2.52	0.45
1:A:781:PRO:O	1:A:785:ARG:HG3	2.16	0.45
1:A:279:ASP:HA	1:A:301:LEU:O	2.17	0.44
1:A:122:HIS:NE2	1:A:350:GLU:OE2	2.47	0.44
1:A:234:VAL:HG23	1:A:234:VAL:O	2.16	0.44
1:A:710:GLU:O	1:A:711:ASN:HB2	2.19	0.43
1:A:718:ASP:OD1	1:A:722:GLY:N	2.52	0.43
1:A:906:GLY:O	1:A:907:ASP:C	2.62	0.42
1:A:68:LEU:O	1:A:72:GLY:HA2	2.21	0.41
1:A:169:VAL:HB	1:A:170:PRO:HD3	2.03	0.41
1:A:86:GLN:O	1:A:89:ASP:N	2.53	0.41
1:A:734:GLU:HA	1:A:758:VAL:HG23	2.02	0.41
1:A:719:SER:HB3	1:A:735:GLU:OE2	2.22	0.40
1:A:344:GLY:HA3	1:A:400:PHE:CZ	2.57	0.40
1:A:891:ASN:O	1:A:892:ASN:C	2.65	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	930/1191 (78%)	893 (96%)	37 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	796/1027 (78%)	794 (100%)	2 (0%)	86	96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	192	ASP
1	A	787	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	A	168	HIS
1	A	686	GLN
1	A	703	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2	15,15,15	0.13	0	21,21,21	2.25	2 (9%)
2	BMA	B	2	2	11,11,12	0.42	0	15,15,17	0.74	0
2	MAN	B	3	2	11,11,12	0.35	0	15,15,17	0.46	0
2	NAG	B	4	2	14,14,15	0.37	0	17,19,21	0.56	0
2	GAL	B	5	2	11,11,12	0.37	0	15,15,17	0.29	0
2	MAN	B	6	2	11,11,12	0.35	0	15,15,17	0.55	0
2	NAG	B	7	2	14,14,15	0.38	0	17,19,21	0.56	0
2	GAL	B	8	2	11,11,12	0.36	0	15,15,17	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2	-	3/6/26/26	0/1/1/1
2	BMA	B	2	2	-	0/2/19/22	0/1/1/1
2	MAN	B	3	2	-	1/2/19/22	0/1/1/1
2	NAG	B	4	2	-	2/6/23/26	0/1/1/1
2	GAL	B	5	2	-	0/2/19/22	0/1/1/1
2	MAN	B	6	2	-	0/2/19/22	0/1/1/1
2	NAG	B	7	2	-	4/6/23/26	0/1/1/1
2	GAL	B	8	2	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	O5-C1-C2	-8.83	100.64	109.52
2	B	1	NAG	C1-C2-C3	-3.44	105.85	110.54

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C3-C2-N2-C7
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2
2	B	7	NAG	C8-C7-N2-C2
2	B	7	NAG	O7-C7-N2-C2
2	B	4	NAG	C4-C5-C6-O6
2	B	4	NAG	O5-C5-C6-O6
2	B	3	MAN	O5-C5-C6-O6
2	B	7	NAG	C3-C2-N2-C7
2	B	7	NAG	C4-C5-C6-O6
2	B	8	GAL	C4-C5-C6-O6

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	NAG	1	0
2	B	7	NAG	2	0
2	B	6	MAN	1	0
2	B	2	BMA	1	0
2	B	8	GAL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	934/1191 (78%)	0.20	28 (2%)	52 43	38, 66, 117, 156	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	156	ASP	6.1
1	A	799	ILE	3.7
1	A	48	PRO	3.7
1	A	909	TYR	3.6
1	A	800	SER	3.4
1	A	804	ASP	3.3
1	A	794	SER	3.3
1	A	798	ASN	3.2
1	A	816	ASP	2.8
1	A	796	THR	2.7
1	A	953	ASP	2.7
1	A	907	ASP	2.7
1	A	868	LYS	2.7
1	A	916	GLU	2.6
1	A	719	SER	2.6
1	A	50	ILE	2.5
1	A	987	ILE	2.5
1	A	329	TRP	2.4
1	A	190	GLY	2.4
1	A	111	GLY	2.3
1	A	854	LEU	2.3
1	A	782	VAL	2.3
1	A	940	LEU	2.3
1	A	236	HIS	2.2
1	A	541	TYR	2.2
1	A	710	GLU	2.1
1	A	914	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	626	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GAL	B	8	11/12	0.47	0.19	136,154,168,172	0
2	GAL	B	5	11/12	0.72	0.15	95,116,139,142	0
2	NAG	B	7	14/15	0.80	0.18	60,115,134,150	0
2	NAG	B	1	15/15	0.88	0.18	49,71,101,113	0
2	MAN	B	6	11/12	0.91	0.12	59,69,82,84	0
2	NAG	B	4	14/15	0.92	0.10	49,55,69,81	0
2	MAN	B	3	11/12	0.94	0.10	49,49,56,57	0
2	BMA	B	2	11/12	0.96	0.09	49,49,62,79	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	A	1301	1/1	0.93	0.06	94,94,94,94	0
4	NI	A	1302	1/1	0.99	0.03	64,64,64,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.