



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 12:32 PM UTC

PDB ID : 6ENQ / pdb_00006enq
Title : Structure of human PPAR gamma LBD in complex with Lanifibranor (IVA337)
Authors : Boubia, B.; Poupardin, O.; Barth, M.; Amaudrut, J.; Broqua, P.; Tallandier, M.; Zeyer, D.
Deposited on : 2017-10-06
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	11% 66% 20% 5% 9%
1	B	282	15% 57% 28% • 12%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4221 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	0	0	0
			2044	1319	334	381	10			
1	B	247	Total	C	N	O	S	0	0	0
			1977	1277	321	369	10			

- Molecule 2 is 4-[1-(1,3-benzothiazol-6-ylsulfonyl)-5-chloro-indol-2-yl]butanoic acid (CCD ID: BJB) (formula: C₁₉H₁₅ClN₂O₄S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O S	0	0
			28	19	1	2	4 2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	104	Total	O	0	0
			104	104		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	68	Total	O	0	0
			68	68		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor gamma



- Molecule 1: Peroxisome proliferator-activated receptor gamma



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	92.88Å 60.00Å 117.49Å 90.00° 102.93° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20 50.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	92.8 (50.00-2.20) 97.7 (50.00-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.14Å)	Xtriage
Refinement program	CNX 2005	Depositor
R, R_{free}	0.227 , 0.271 0.228 , 0.230	Depositor DCC
R_{free} test set	1687 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	40.2	Xtriage
Anisotropy	0.403	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4221	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BJB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2078	0.83	5/2801 (0.2%)
1	B	0.40	0/2009	0.74	0/2707
All	All	0.43	0/4087	0.79	5/5508 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	244	LYS	N-CA-C	6.37	117.53	108.74
1	A	205	ASN	N-CA-C	-5.55	99.19	109.06
1	A	207	GLU	N-CA-C	-5.43	99.24	110.80
1	A	208	SER	N-CA-C	-5.14	107.00	113.28
1	A	206	PRO	CA-C-O	-5.09	116.04	123.07

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	239	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2044	0	2105	69	0
1	B	1977	0	2026	61	0
2	A	28	0	0	4	0
3	A	104	0	0	5	0
3	B	68	0	0	2	0
All	All	4221	0	4131	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (129) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:GLY:HA3	1:A:240:LYS:HD2	1.31	1.08
1:A:239:GLY:HA3	1:A:240:LYS:CD	2.00	0.92
1:B:335:ASN:HD22	1:B:337:ASP:H	1.12	0.90
1:B:335:ASN:ND2	1:B:337:ASP:H	1.76	0.82
1:B:349:THR:HG22	1:B:351:GLU:H	1.42	0.82
1:A:205:ASN:HD21	1:A:207:GLU:HB3	1.51	0.75
1:A:474:LYS:O	1:A:475:ASP:HB2	1.85	0.73
1:A:276:GLU:OE1	1:A:357:ARG:NH1	2.22	0.72
1:A:252:MET:HE1	1:A:277:VAL:HG11	1.73	0.70
1:A:395:GLY:HA2	1:A:400:LEU:CD1	2.22	0.70
1:A:457:LYS:NZ	1:A:457:LYS:HA	2.08	0.69
1:A:457:LYS:HD3	1:A:465:LEU:HD11	1.76	0.68
1:A:205:ASN:ND2	1:A:207:GLU:HB3	2.09	0.67
1:B:466:HIS:N	1:B:467:PRO:HD2	2.10	0.66
1:A:285:CYS:SG	2:A:501:BJB:O29	2.53	0.66
1:B:465:LEU:C	1:B:467:PRO:HD2	2.20	0.66
1:B:349:THR:HG22	1:B:351:GLU:N	2.11	0.63
1:A:252:MET:CE	1:A:277:VAL:HG11	2.29	0.63
1:A:279:ILE:O	1:A:283:GLN:HG2	2.00	0.62
1:B:335:ASN:HD22	1:B:337:ASP:N	1.93	0.61
1:A:235:ALA:HA	1:A:240:LYS:HG3	1.81	0.60
1:B:358:LYS:HD2	1:B:358:LYS:H	1.65	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:457:LYS:HE2	1:A:465:LEU:HD13	1.83	0.60
1:B:335:ASN:HD22	1:B:335:ASN:C	2.10	0.59
1:A:239:GLY:H	1:A:240:LYS:HG2	1.65	0.59
1:A:370:PHE:HB2	1:A:445:ILE:HD11	1.85	0.58
1:A:453:LEU:HD12	1:A:465:LEU:HD21	1.85	0.57
1:B:451:GLN:O	1:B:454:GLN:HG2	2.03	0.57
1:B:466:HIS:O	1:B:469:LEU:N	2.37	0.57
1:B:447:THR:O	1:B:450:VAL:HG22	2.04	0.57
1:A:325:ILE:HD11	1:A:392:ILE:CG1	2.35	0.57
1:B:328:THR:OG1	1:B:442:LEU:HD11	2.05	0.57
1:A:360:PHE:O	1:A:363:PHE:HD2	1.89	0.56
1:A:354:LYS:HA	1:A:361:GLY:O	2.06	0.56
1:A:457:LYS:HD3	1:A:465:LEU:CD1	2.35	0.56
1:B:252:MET:O	1:B:256:MET:HG2	2.06	0.55
1:B:454:GLN:O	1:B:457:LYS:HB3	2.06	0.55
1:A:471:GLU:O	1:A:474:LYS:HG2	2.06	0.55
1:B:452:LEU:O	1:B:456:ILE:HG12	2.06	0.55
1:A:457:LYS:NZ	3:A:602:HOH:O	2.40	0.54
1:B:466:HIS:N	1:B:467:PRO:CD	2.70	0.54
1:A:239:GLY:N	1:A:240:LYS:HG2	2.23	0.54
1:A:433:ALA:O	1:A:437:GLN:HG3	2.08	0.54
1:B:358:LYS:HD2	1:B:358:LYS:N	2.22	0.54
1:A:325:ILE:HD11	1:A:392:ILE:HG13	1.90	0.53
1:B:255:LEU:O	1:B:259:GLU:HB2	2.07	0.53
1:B:343:GLU:O	1:B:343:GLU:HG3	2.08	0.53
1:A:370:PHE:CB	1:A:445:ILE:HD11	2.37	0.53
1:A:232:LYS:HE3	1:A:243:ASP:OD1	2.09	0.53
1:A:205:ASN:ND2	1:A:207:GLU:CB	2.71	0.53
1:B:301:LYS:HD3	3:B:525:HOH:O	2.08	0.53
1:A:474:LYS:O	1:A:475:ASP:CB	2.57	0.52
1:B:226:PHE:CG	1:B:329:MET:HE1	2.44	0.52
1:A:305:GLY:HA2	1:A:308:ASN:HD22	1.75	0.52
1:B:365:GLU:N	1:B:366:PRO:HD2	2.25	0.52
1:A:285:CYS:SG	2:A:501:BJB:C9	2.98	0.51
1:A:348:MET:SD	1:A:353:LEU:HD21	2.50	0.51
1:A:212:ARG:O	1:A:216:LYS:HG3	2.10	0.51
1:B:208:SER:O	1:B:212:ARG:HG2	2.10	0.51
1:B:348:MET:SD	1:B:353:LEU:HD21	2.51	0.51
1:B:234:ARG:HD2	3:B:560:HOH:O	2.10	0.51
1:A:205:ASN:HB2	1:A:206:PRO:HD2	1.92	0.51
1:A:235:ALA:O	1:A:240:LYS:HG2	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:404:LYS:N	1:A:405:PRO:HD2	2.26	0.50
1:B:454:GLN:HG3	1:B:455:VAL:N	2.25	0.50
1:A:391:ILE:HG12	1:A:439:MET:HE3	1.94	0.50
1:B:336:LYS:NZ	1:B:336:LYS:HB3	2.26	0.50
1:B:319:LYS:HD3	1:B:320:TYR:CE1	2.47	0.50
1:B:255:LEU:HD22	1:B:281:ILE:HD11	1.93	0.49
2:A:501:BJB:C35	2:A:501:BJB:C3	2.90	0.49
1:B:282:PHE:O	1:B:286:GLN:HG2	2.11	0.49
1:A:357:ARG:HG2	1:A:358:LYS:O	2.13	0.49
1:A:457:LYS:HA	1:A:457:LYS:HZ3	1.78	0.49
1:A:464:SER:C	1:A:465:LEU:HD12	2.38	0.48
1:B:276:GLU:O	1:B:280:ARG:HG3	2.14	0.48
1:B:318:LEU:O	1:B:322:VAL:HG13	2.14	0.48
1:A:427:GLU:HB2	3:A:657:HOH:O	2.13	0.47
1:A:457:LYS:NZ	1:A:463:MET:HB3	2.29	0.47
1:A:253:ASN:HD22	1:A:253:ASN:N	2.11	0.47
1:A:257:MET:HA	1:A:257:MET:HE2	1.95	0.47
1:A:207:GLU:H	1:A:209:ALA:H	1.63	0.47
1:A:434:LYS:HE2	1:B:411:ASP:OD2	2.15	0.47
1:B:339:VAL:HG22	1:B:340:LEU:O	2.15	0.47
1:A:276:GLU:CD	1:A:357:ARG:NH1	2.72	0.47
1:B:435:LEU:O	1:B:438:LYS:HB2	2.15	0.47
1:A:242:THR:O	1:A:243:ASP:O	2.33	0.46
1:B:404:LYS:HB3	1:B:405:PRO:HD3	1.97	0.46
1:A:251:ASP:OD1	1:A:251:ASP:C	2.58	0.46
1:A:437:GLN:OE1	1:B:439:MET:HE1	2.16	0.46
1:A:290:VAL:HG13	1:A:468:LEU:HD23	1.97	0.46
1:A:453:LEU:CD1	1:A:465:LEU:HD21	2.46	0.46
1:B:379:LEU:HD11	1:B:435:LEU:HD13	1.98	0.45
1:B:430:GLN:O	1:B:434:LYS:HD2	2.15	0.45
1:A:452:LEU:O	1:A:456:ILE:HG13	2.16	0.45
1:B:245:SER:HA	1:B:246:PRO:HD3	1.73	0.45
1:B:446:VAL:O	1:B:450:VAL:HG13	2.17	0.45
1:B:462:ASP:C	1:B:463:MET:HG3	2.42	0.45
1:A:234:ARG:O	1:A:238:THR:HB	2.16	0.45
1:B:335:ASN:ND2	1:B:337:ASP:N	2.56	0.45
1:A:289:SER:OG	2:A:501:BJB:N26	2.50	0.44
1:B:324:GLU:OE2	1:B:443:ARG:HD2	2.15	0.44
1:A:277:VAL:O	1:A:281:ILE:HG13	2.18	0.44
1:A:238:THR:HG22	1:A:240:LYS:HD3	1.99	0.44
1:A:353:LEU:HD13	1:A:364:MET:HG3	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:463:MET:HB2	1:B:466:HIS:CD2	2.53	0.43
1:B:255:LEU:HD23	1:B:277:VAL:HG13	2.01	0.43
1:B:290:VAL:HG21	1:B:473:TYR:CD1	2.54	0.43
1:B:335:ASN:ND2	1:B:338:GLY:H	2.17	0.42
1:B:356:LEU:O	1:B:357:ARG:O	2.38	0.42
1:A:363:PHE:CZ	1:A:364:MET:HG2	2.55	0.42
1:B:276:GLU:HB3	1:B:279:ILE:HB	2.01	0.42
3:A:681:HOH:O	1:B:434:LYS:HE2	2.18	0.42
1:B:216:LYS:HE3	1:B:220:ASP:OD2	2.19	0.42
1:B:233:ALA:O	1:B:237:LEU:HG	2.19	0.42
1:B:256:MET:C	1:B:258:GLY:H	2.28	0.42
1:A:239:GLY:HA3	1:A:240:LYS:CG	2.50	0.42
1:A:402:ASN:HB2	3:A:635:HOH:O	2.20	0.41
1:A:277:VAL:CG2	1:A:278:ALA:N	2.82	0.41
1:A:207:GLU:O	1:A:211:LEU:HG	2.18	0.41
1:B:365:GLU:N	1:B:366:PRO:CD	2.84	0.41
1:A:234:ARG:HA	1:A:234:ARG:HD2	1.78	0.41
1:B:329:MET:O	1:B:332:SER:HB2	2.20	0.41
1:B:368:PHE:O	1:B:372:VAL:HG23	2.21	0.41
1:B:466:HIS:O	1:B:467:PRO:C	2.64	0.41
1:A:364:MET:HB2	3:A:633:HOH:O	2.21	0.40
1:A:290:VAL:O	1:A:294:GLN:HG3	2.21	0.40
1:A:457:LYS:HE2	1:A:465:LEU:CD1	2.48	0.40
1:B:290:VAL:HG21	1:B:473:TYR:HD1	1.87	0.40
1:B:313:ASP:O	1:B:317:LEU:HG	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	252/282 (89%)	243 (96%)	7 (3%)	2 (1%)	16 16

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	241/282 (86%)	231 (96%)	8 (3%)	2 (1%)	16	16
All	All	493/564 (87%)	474 (96%)	15 (3%)	4 (1%)	16	16

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	357	ARG
1	A	207	GLU
1	A	243	ASP
1	B	474	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/255 (90%)	216 (94%)	13 (6%)	18	23
1	B	221/255 (87%)	208 (94%)	13 (6%)	18	22
All	All	450/510 (88%)	424 (94%)	26 (6%)	18	22

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	204	LEU
1	A	208	SER
1	A	234	ARG
1	A	238	THR
1	A	244	LYS
1	A	277	VAL
1	A	357	ARG
1	A	358	LYS
1	A	443	ARG
1	A	452	LEU
1	A	453	LEU
1	A	457	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	470	GLN
1	B	210	ASP
1	B	224	LYS
1	B	228	LEU
1	B	232	LYS
1	B	252	MET
1	B	318	LEU
1	B	335	ASN
1	B	336	LYS
1	B	362	ASP
1	B	412	ASN
1	B	427	GLU
1	B	438	LYS
1	B	443	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	205	ASN
1	A	253	ASN
1	A	308	ASN
1	A	314	GLN
1	A	323	HIS
1	A	402	ASN
1	A	410	GLN
1	A	415	GLN
1	A	444	GLN
1	A	454	GLN
1	B	308	ASN
1	B	335	ASN
1	B	412	ASN
1	B	430	GLN
1	B	437	GLN
1	B	470	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BJB	A	501	-	31,31,31	2.25	7 (22%)	39,46,46	3.86	6 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BJB	A	501	-	-	8/18/18/18	0/4/4/4

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	BJB	C25-N26	6.27	1.38	1.29
2	A	501	BJB	C2-N1	-6.23	1.37	1.42
2	A	501	BJB	S14-N1	-3.46	1.62	1.68
2	A	501	BJB	C3-C2	3.43	1.40	1.35
2	A	501	BJB	C31-C2	2.78	1.53	1.50
2	A	501	BJB	C18-C17	2.37	1.43	1.40
2	A	501	BJB	O41-C38	-2.13	1.23	1.30

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	BJB	S24-C25-N26	-16.04	106.94	117.01
2	A	501	BJB	C18-N26-C25	15.19	117.49	109.83
2	A	501	BJB	C15-S14-N1	6.72	113.28	104.90
2	A	501	BJB	C16-C17-C18	-3.52	119.38	121.99
2	A	501	BJB	C16-C17-S24	2.51	130.97	126.86
2	A	501	BJB	C4-C3-C2	-2.00	106.06	108.41

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	BJB	C20-C15-S14-N1
2	A	501	BJB	C20-C15-S14-O30
2	A	501	BJB	C16-C15-S14-O30
2	A	501	BJB	C16-C15-S14-N1
2	A	501	BJB	C3-C2-C31-C32
2	A	501	BJB	C32-C35-C38-O42
2	A	501	BJB	C32-C35-C38-O41
2	A	501	BJB	N1-C2-C31-C32

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	BJB	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	256/282 (90%)	0.59	32 (12%) 8 6	28, 45, 79, 100	0
1	B	247/282 (87%)	0.85	42 (17%) 4 3	28, 49, 105, 128	0
All	All	503/564 (89%)	0.72	74 (14%) 6 4	28, 47, 95, 128	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	459	THR	4.9
1	B	257	MET	4.9
1	B	210	ASP	4.7
1	A	243	ASP	4.6
1	A	258	GLY	4.5
1	B	463	MET	4.4
1	B	363	PHE	4.1
1	A	471	GLU	4.1
1	B	461	THR	4.0
1	B	475	ASP	3.8
1	B	279	ILE	3.8
1	B	462	ASP	3.8
1	A	394	SER	3.8
1	A	361	GLY	3.6
1	B	465	LEU	3.6
1	A	239	GLY	3.6
1	B	209	ALA	3.4
1	B	343	GLU	3.3
1	B	469	LEU	3.3
1	B	252	MET	3.2
1	A	254	SER	3.2
1	B	253	ASN	3.2
1	B	251	ASP	3.1
1	A	282	PHE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	460	GLU	3.0
1	B	248	VAL	3.0
1	B	238	THR	2.9
1	B	254	SER	2.9
1	A	256	MET	2.9
1	A	240	LYS	2.8
1	A	277	VAL	2.8
1	B	258	GLY	2.8
1	B	468	LEU	2.8
1	B	473	TYR	2.8
1	B	278	ALA	2.8
1	B	455	VAL	2.8
1	B	358	LYS	2.7
1	A	280	ARG	2.7
1	A	242	THR	2.7
1	A	472	ILE	2.7
1	A	257	MET	2.7
1	A	285	CYS	2.7
1	B	464	SER	2.7
1	A	278	ALA	2.6
1	B	361	GLY	2.6
1	A	204	LEU	2.6
1	A	363	PHE	2.6
1	B	277	VAL	2.6
1	A	474	LYS	2.5
1	B	346	GLY	2.5
1	B	345	GLN	2.5
1	A	279	ILE	2.5
1	B	255	LEU	2.4
1	A	205	ASN	2.4
1	B	256	MET	2.4
1	B	359	PRO	2.4
1	B	467	PRO	2.4
1	B	353	LEU	2.4
1	A	457	LYS	2.3
1	A	284	GLY	2.3
1	A	241	THR	2.3
1	A	461	THR	2.3
1	B	357	ARG	2.3
1	A	286	GLN	2.3
1	B	466	HIS	2.3
1	B	245	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	450	VAL	2.2
1	A	276	GLU	2.2
1	B	474	LYS	2.1
1	A	395	GLY	2.1
1	A	238	THR	2.1
1	A	359	PRO	2.1
1	A	255	LEU	2.1
1	B	457	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BJB	A	501	28/28	0.62	0.19	48,56,66,70	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BJB A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.