



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 10:20 PM UTC

PDB ID : 7EOX / pdb_00007eox
Title : Protease structure from Euphorbia resinifera
Authors : Siritapetawee, J.; Attarataya, J.; Charoenwattanasatien, R.
Deposited on : 2021-04-24
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	654	13% 90% 6% .
1	B	654	2% 92% . .
2	C	7	14% 86%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 10263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	634	Total	C	N	O	S	0	0	0
			4719	2963	783	956	17			
1	B	634	Total	C	N	O	S	0	0	0
			4719	2963	783	956	17			

- Molecule 2 is an oligosaccharide called beta-D-xylopyranose-(1-2)-[alpha-D-mannopyranose-(1-3)][alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	7	Total	C	N	O	0	0	0
			80	45	2	33			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	I	0	0
			2	2		
4	B	2	Total	I	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	277	Total	O	0	0
			277	277		
5	B	422	Total	O	0	0
			422	422		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protease



• Molecule 1: Protease



• Molecule 2: beta-D-xylopyranose-(1-2)-[alpha-D-mannopyranose-(1-3)][alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.67Å 112.88Å 201.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.01 – 1.70 60.01 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.6 (60.01-1.70) 98.6 (60.01-1.70)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.203 , 0.237 0.214 , 0.245	Depositor DCC
R_{free} test set	8569 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	22.7	Xtriage
Anisotropy	0.148	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 26.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10263	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, NAG, XYP, BMA, FUC, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.10	1/4836 (0.0%)	1.34	10/6601 (0.2%)
1	B	1.08	2/4836 (0.0%)	1.26	9/6601 (0.1%)
All	All	1.09	3/9672 (0.0%)	1.30	19/13202 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	477	ILE	C-O	6.36	1.30	1.24
1	B	289	VAL	N-CA	5.07	1.50	1.46
1	A	289	VAL	N-CA	5.04	1.50	1.46

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	ASP	CA-CB-CG	8.94	121.54	112.60
1	B	1	ASP	CA-CB-CG	8.30	120.90	112.60
1	B	654	TRP	CA-C-O	-8.12	107.00	120.80
1	A	654	TRP	CA-C-O	-7.20	108.56	120.80
1	A	407	ALA	CA-C-N	6.71	126.71	120.34
1	A	407	ALA	C-N-CA	6.71	126.71	120.34
1	A	578	VAL	N-CA-C	-6.20	106.73	112.43
1	B	479	ASP	CA-CB-CG	6.12	118.72	112.60
1	A	167	ASP	CA-CB-CG	5.80	118.40	112.60
1	B	392	GLN	CB-CG-CD	5.48	121.91	112.60
1	B	209	GLY	CA-C-O	-5.38	118.58	122.45
1	B	471	ASP	CA-CB-CG	5.36	117.96	112.60
1	A	106	HIS	CA-C-N	5.32	125.89	119.98
1	A	106	HIS	C-N-CA	5.32	125.89	119.98
1	B	641	ASP	CA-C-N	5.24	125.89	120.03
1	B	641	ASP	C-N-CA	5.24	125.89	120.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	184	ASP	CA-CB-CG	5.22	117.82	112.60
1	B	433	THR	N-CA-C	-5.05	106.38	112.54
1	A	198	VAL	CA-C-O	-5.04	116.02	121.27

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4719	0	4495	17	0
1	B	4719	0	4493	9	0
2	C	80	0	60	0	0
3	A	14	0	13	0	0
3	B	28	0	26	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	277	0	0	1	0
5	B	422	0	0	5	0
All	All	10263	0	9087	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:THR:O	1:A:112:THR:HB	1.76	0.85
1:A:112:THR:HG21	1:A:408:GLY:H	1.44	0.80
1:A:112:THR:HG21	1:A:408:GLY:N	2.15	0.60
1:A:112:THR:HG22	1:A:407:ALA:HB1	1.84	0.60
1:B:247:ARG:NH1	5:B:801:HOH:O	2.23	0.57
1:B:198:VAL:HG22	5:B:1139:HOH:O	2.08	0.53
1:A:411:GLY:O	1:A:424:ILE:HD12	2.09	0.52
1:A:112:THR:CG2	1:A:408:GLY:H	2.18	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:ASP:HB2	1:A:406:LEU:O	2.10	0.51
1:A:145:LYS:HE2	1:A:148:TRP:CZ3	2.47	0.48
1:B:35:VAL:HB	1:B:143:VAL:HG21	1.96	0.47
1:B:129:LYS:HE2	5:B:846:HOH:O	2.15	0.47
1:B:286:SER:O	1:B:290:GLN:HG2	2.16	0.46
1:A:535:GLU:O	1:A:537:PRO:HD3	2.16	0.45
1:A:322:THR:CG2	1:A:339:THR:CG2	2.96	0.44
1:B:199:LYS:HB3	5:B:811:HOH:O	2.17	0.44
1:A:22:TYR:HB2	1:A:138:LYS:HG3	2.00	0.44
1:B:318:MET:O	1:B:337:ALA:HA	2.18	0.43
1:A:175:SER:HB3	1:A:434:SER:HA	2.01	0.43
1:A:199:LYS:HB3	5:A:886:HOH:O	2.18	0.43
1:B:269:TYR:OH	1:B:297:ASP:OD2	2.36	0.43
1:A:54:TRP:CZ2	1:A:56:GLY:HA3	2.55	0.41
1:B:434:SER:HB2	5:B:911:HOH:O	2.20	0.41
1:A:540:ASP:HB2	1:A:543:ASP:OD2	2.21	0.41
1:A:4:THR:N	1:A:5:PRO:CD	2.85	0.40
1:A:488:PRO:HD2	1:A:634:PHE:CD1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	628/654 (96%)	604 (96%)	23 (4%)	1 (0%)	43	28
1	B	628/654 (96%)	608 (97%)	19 (3%)	1 (0%)	43	28
All	All	1256/1308 (96%)	1212 (96%)	42 (3%)	2 (0%)	43	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	330	GLY
1	B	330	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	517/528 (98%)	504 (98%)	13 (2%)	42	24
1	B	517/528 (98%)	508 (98%)	9 (2%)	53	38
All	All	1034/1056 (98%)	1012 (98%)	22 (2%)	47	30

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	ASP
1	A	17	TRP
1	A	112	THR
1	A	121	ASN
1	A	131	GLN
1	A	137	THR
1	A	199	LYS
1	A	298	ASN
1	A	361	LYS
1	A	392	GLN
1	A	417	ASP
1	A	563	GLU
1	A	575	VAL
1	B	17	TRP
1	B	53	SER
1	B	129	LYS
1	B	199	LYS
1	B	298	ASN
1	B	392	GLN
1	B	417	ASP
1	B	563	GLU
1	B	609	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	196	ASN
1	A	544	ASN
1	B	19	ASN
1	B	131	GLN
1	B	196	ASN
1	B	497	HIS
1	B	591	ASN
1	B	611	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.60	0	17,19,21	1.41	1 (5%)
2	NAG	C	2	2	14,14,15	0.66	0	17,19,21	1.42	2 (11%)
2	BMA	C	3	2	11,11,12	0.83	0	15,15,17	1.31	2 (13%)
2	XYP	C	4	2	9,9,10	0.69	0	10,12,14	1.12	1 (10%)
2	MAN	C	5	2	11,11,12	0.51	0	15,15,17	1.00	1 (6%)
2	MAN	C	6	2	11,11,12	0.75	0	15,15,17	1.00	0
2	FUC	C	7	2	10,10,11	0.40	0	14,14,16	1.11	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	XYP	C	4	2	-	-	0/1/1/1
2	MAN	C	5	2	-	2/2/19/22	0/1/1/1
2	MAN	C	6	2	-	0/2/19/22	0/1/1/1
2	FUC	C	7	2	-	-	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	C1-C2-N2	-3.27	105.28	110.43
2	C	2	NAG	C1-O5-C5	3.02	116.23	112.19
2	C	3	BMA	O2-C2-C3	2.96	116.29	110.15
2	C	2	NAG	O6-C6-C5	-2.42	103.10	111.33
2	C	4	XYP	C4-C3-C2	-2.22	108.28	110.92
2	C	7	FUC	O2-C2-C3	-2.12	105.75	110.15
2	C	5	MAN	O5-C5-C6	2.07	111.69	107.66
2	C	3	BMA	O2-C2-C1	-2.00	104.64	109.22

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	5	MAN	C4-C5-C6-O6
2	C	5	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	701	1	14,14,15	1.14	1 (7%)	17,19,21	0.89	0
3	NAG	B	701	1	14,14,15	0.82	1 (7%)	17,19,21	1.47	3 (17%)
3	NAG	B	702	1	14,14,15	1.34	1 (7%)	17,19,21	1.32	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	701	1	-	0/6/23/26	0/1/1/1
3	NAG	B	701	1	-	2/6/23/26	0/1/1/1
3	NAG	B	702	1	-	2/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	702	NAG	C1-C2	3.72	1.57	1.52
3	A	701	NAG	C1-C2	2.43	1.55	1.52
3	B	701	NAG	C1-C2	2.27	1.55	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	701	NAG	C4-C3-C2	-3.18	106.36	111.02
3	B	702	NAG	O3-C3-C4	2.70	116.74	110.38
3	B	701	NAG	C1-O5-C5	2.24	115.19	112.19
3	B	702	NAG	O5-C1-C2	-2.21	107.87	111.29
3	B	701	NAG	O3-C3-C2	2.21	114.00	109.40
3	B	702	NAG	C4-C3-C2	-2.01	108.08	111.02

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	701	NAG	O5-C5-C6-O6
3	B	701	NAG	C4-C5-C6-O6
3	B	702	NAG	C4-C5-C6-O6
3	B	702	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	634/654 (96%)	0.82	84 (13%) 7 7	16, 29, 51, 76	0
1	B	634/654 (96%)	0.16	16 (2%) 58 62	15, 23, 41, 71	0
All	All	1268/1308 (96%)	0.49	100 (7%) 18 20	15, 26, 48, 76	0

All (100) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	83	ILE	6.3
1	A	566	THR	4.7
1	B	565	ASP	4.1
1	A	58	CYS	3.9
1	A	260	LEU	3.8
1	A	150	ALA	3.8
1	A	81	LYS	3.7
1	A	64	PHE	3.7
1	A	103	ILE	3.7
1	A	66	ALA	3.6
1	A	73	LEU	3.6
1	A	565	ASP	3.6
1	B	150	ALA	3.6
1	A	564	GLY	3.5
1	A	54	TRP	3.5
1	A	151	GLY	3.4
1	A	60	THR	3.3
1	B	98	ASP	3.3
1	B	290	GLN	3.3
1	A	42	PHE	3.2
1	A	409	TRP	3.2
1	A	364	PHE	3.2
1	B	259	PRO	3.1
1	A	242	VAL	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	536	VAL	3.1
1	A	264	PHE	3.1
1	B	536	VAL	3.1
1	B	46	ASP	3.1
1	A	98	ASP	3.1
1	A	335	PHE	3.0
1	A	57	THR	3.0
1	A	259	PRO	3.0
1	A	148	TRP	3.0
1	A	538	PRO	3.0
1	A	36	TRP	2.9
1	B	83	ILE	2.9
1	A	121	ASN	2.9
1	A	425	VAL	2.9
1	B	198	VAL	2.9
1	A	418	LEU	2.9
1	A	45	SER	2.9
1	B	624	LEU	2.8
1	A	50	ILE	2.8
1	A	99	SER	2.8
1	A	63	ASP	2.7
1	A	424	ILE	2.7
1	A	46	ASP	2.7
1	A	624	LEU	2.7
1	A	52	SER	2.7
1	A	265	PHE	2.7
1	A	255	TYR	2.7
1	A	33	THR	2.6
1	A	334	LEU	2.6
1	A	365	LEU	2.6
1	A	422	ASP	2.6
1	A	261	GLN	2.6
1	A	82	GLY	2.6
1	A	427	PHE	2.6
1	A	69	CYS	2.5
1	A	143	VAL	2.5
1	A	56	GLY	2.5
1	A	61	SER	2.5
1	A	369	ILE	2.5
1	A	363	VAL	2.5
1	A	122	ALA	2.4
1	A	239	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1	ASP	2.4
1	A	412	ALA	2.4
1	A	256	THR	2.4
1	A	22	TYR	2.4
1	A	62	ASP	2.4
1	B	258	GLU	2.4
1	A	546	LYS	2.4
1	A	366	GLY	2.3
1	A	563	GLU	2.3
1	A	263	GLU	2.3
1	A	51	PRO	2.3
1	A	128	ALA	2.3
1	B	261	GLN	2.3
1	B	371	ASN	2.3
1	A	368	VAL	2.3
1	A	367	THR	2.3
1	A	248	THR	2.2
1	A	149	SER	2.2
1	A	361	LYS	2.2
1	A	246	GLY	2.2
1	A	43	SER	2.2
1	A	356	GLN	2.2
1	B	631	ALA	2.2
1	A	542	CYS	2.2
1	A	124	PHE	2.2
1	A	372	SER	2.2
1	B	542	CYS	2.1
1	A	411	GLY	2.1
1	A	74	ILE	2.1
1	A	47	MET	2.1
1	A	34	GLY	2.1
1	B	287	SER	2.1
1	A	355	SER	2.0
1	A	159	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	C	5	11/12	0.80	0.15	50,54,57,64	0
2	FUC	C	7	10/11	0.87	0.12	35,38,40,41	0
2	NAG	C	1	14/15	0.93	0.09	27,28,32,36	0
2	NAG	C	2	14/15	0.93	0.10	25,30,37,47	0
2	MAN	C	6	11/12	0.94	0.09	25,26,30,31	0
2	XYP	C	4	9/10	0.95	0.09	31,31,36,37	0
2	BMA	C	3	11/12	0.95	0.08	27,30,37,43	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	B	702	14/15	0.80	0.15	40,46,51,51	0
3	NAG	B	701	14/15	0.81	0.14	44,50,54,57	0
3	NAG	A	701	14/15	0.82	0.13	37,44,47,52	0
4	IOD	A	702	1/1	0.99	0.03	25,25,25,25	0
4	IOD	A	703	1/1	0.99	0.04	31,31,31,31	0
4	IOD	B	704	1/1	0.99	0.03	28,28,28,28	0
4	IOD	B	703	1/1	1.00	0.02	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.