



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:35 AM UTC

PDB ID : 2EPK / pdb_00002epk
Title : N-acetyl-B-D-glucosaminidase (GCNA) from Streptococcus gordonii
Authors : Langley, D.B.
Deposited on : 2007-03-30
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	627	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5241 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-acetyl-beta-D-glucosaminidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	X	613	Total	C	N	O	S	Se	0	1	0
			4989	3185	838	945	7	14			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	X	217	Total	O	0	0
			217	217		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-acetyl-beta-D-glucosaminidase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	123.89Å 124.51Å 148.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.97 – 1.85 39.97 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.6 (39.97-1.85) 99.7 (39.97-1.85)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.215 , 0.235 0.201 , 0.219	Depositor DCC
R_{free} test set	4876 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.217	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.014 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5241	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	X	0.61	0/5084	0.76	1/6844 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	X	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	X	129	ILE	N-CA-C	-6.99	97.55	107.75

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	X	129	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	4989	0	4874	25	0
2	X	35	0	0	0	0
3	X	217	0	0	0	0
All	All	5241	0	4874	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:7:LEU:HD23	1:X:31:LEU:HD11	1.79	0.64
1:X:130:GLU:C	1:X:131:ASN:HD22	2.11	0.58
1:X:275:MSE:HE3	1:X:323:ASN:HB3	1.91	0.53
1:X:275:MSE:HE3	1:X:323:ASN:CB	2.39	0.52
1:X:37:THR:HG23	1:X:81:GLU:HG2	1.92	0.50
1:X:543:PHE:CE2	1:X:559:ILE:HD13	2.47	0.49
1:X:509:ILE:HD12	1:X:577:LEU:HD11	1.96	0.47
1:X:110:VAL:CG1	1:X:114:MSE:HE3	2.47	0.45
1:X:509:ILE:HD13	1:X:524:LEU:HD13	1.98	0.45
1:X:7:LEU:CD2	1:X:31:LEU:HD11	2.46	0.45
1:X:315:GLU:CD	1:X:315:GLU:H	2.25	0.45
1:X:505:ILE:O	1:X:509:ILE:HG12	2.17	0.44
1:X:130:GLU:C	1:X:131:ASN:ND2	2.75	0.44
1:X:7:LEU:HD22	1:X:11:GLN:HB3	1.99	0.43
1:X:95:ARG:O	1:X:96:ASN:HB2	2.18	0.43
1:X:95:ARG:O	1:X:96:ASN:CB	2.67	0.43
1:X:375:GLY:O	1:X:376:ASP:C	2.62	0.43
1:X:32:VAL:O	1:X:51:SER:HA	2.20	0.42
1:X:409:PHE:CZ	1:X:413:THR:HG21	2.55	0.41
1:X:444:TYR:CD2	1:X:455:LEU:HD22	2.55	0.41
1:X:242:ARG:O	1:X:246:MSE:HG2	2.21	0.41
1:X:107:MSE:HG3	1:X:111:LEU:HD22	2.01	0.41
1:X:334:ALA:HA	1:X:370:ILE:O	2.20	0.40
1:X:482:ALA:HB3	1:X:486:ALA:HA	2.02	0.40
1:X:125:ASP:HA	1:X:135:PHE:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	610/627 (97%)	593 (97%)	14 (2%)	3 (0%)	24 13

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	130	GLU
1	X	96	ASN
1	X	376	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	532/528 (101%)	520 (98%)	12 (2%)	44 30

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	15	LEU
1	X	95	ARG
1	X	111	LEU
1	X	130	GLU
1	X	131	ASN
1	X	226	HIS
1	X	234	LEU
1	X	244	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	315	GLU
1	X	331	ILE
1	X	415	LEU
1	X	624	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	54	GLN
1	X	96	ASN
1	X	131	ASN
1	X	132	GLN
1	X	148	GLN
1	X	208	GLN
1	X	241	ASN
1	X	248	GLN
1	X	318	ASN
1	X	473	GLN
1	X	515	ASN
1	X	546	GLN
1	X	574	GLN
1	X	613	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	X	742	-	4,4,4	0.20	0	6,6,6	0.25	0
2	SO4	X	740	-	4,4,4	0.25	0	6,6,6	0.20	0
2	SO4	X	744	-	4,4,4	0.24	0	6,6,6	0.10	0
2	SO4	X	741	-	4,4,4	0.23	0	6,6,6	0.17	0
2	SO4	X	746	-	4,4,4	0.24	0	6,6,6	0.15	0
2	SO4	X	743	-	4,4,4	0.24	0	6,6,6	0.15	0
2	SO4	X	745	-	4,4,4	0.26	0	6,6,6	0.06	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	599/627 (95%)	0.43	52 (8%) 16 16	14, 31, 53, 66	2 (0%)

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	625	TYR	7.4
1	X	35	SER	4.6
1	X	70	ARG	4.3
1	X	131	ASN	3.8
1	X	226	HIS	3.6
1	X	274	LEU	3.5
1	X	626	THR	3.4
1	X	289	GLU	3.4
1	X	23	ASN	3.3
1	X	515	ASN	3.3
1	X	288	PRO	3.2
1	X	33	THR	3.2
1	X	291	THR	3.2
1	X	273	LYS	3.2
1	X	624	ILE	3.1
1	X	26	GLU	3.1
1	X	519	GLU	3.0
1	X	511	LYS	3.0
1	X	130	GLU	3.0
1	X	240	GLN	2.9
1	X	514	ARG	2.8
1	X	180	LYS	2.8
1	X	531	GLN	2.7
1	X	37	THR	2.7
1	X	584	GLN	2.6
1	X	516	GLY	2.5
1	X	77	GLN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	527	LYS	2.5
1	X	523	ALA	2.5
1	X	484	GLU	2.4
1	X	24	LEU	2.4
1	X	234	LEU	2.4
1	X	415	LEU	2.4
1	X	297	ARG	2.4
1	X	290	GLU	2.3
1	X	526	GLU	2.3
1	X	512	ALA	2.3
1	X	238	GLY	2.3
1	X	594	GLU	2.3
1	X	604	ASP	2.3
1	X	225	ALA	2.2
1	X	330	ASP	2.2
1	X	36	ALA	2.2
1	X	72	GLY	2.1
1	X	239	PHE	2.1
1	X	623	THR	2.1
1	X	331	ILE	2.1
1	X	509	ILE	2.1
1	X	75	GLU	2.1
1	X	12	GLU	2.0
1	X	241	ASN	2.0
1	X	310	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	X	744	5/5	0.59	0.16	108,108,109,109	0
2	SO4	X	746	5/5	0.78	0.13	80,80,83,86	0
2	SO4	X	745	5/5	0.79	0.14	92,95,96,96	0
2	SO4	X	741	5/5	0.79	0.15	72,73,78,79	0
2	SO4	X	743	5/5	0.88	0.10	46,50,59,59	0
2	SO4	X	740	5/5	0.96	0.08	43,44,46,49	0
2	SO4	X	742	5/5	0.96	0.08	42,44,45,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.