



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:32 PM UTC

PDB ID : 7EPF / pdb_00007epf
Title : Crystal structure of mGlu2 bound to NAM597
Authors : Du, J.; Wang, D.; Lin, S.; Han, S.; Wu, B.; Zhao, Q.
Deposited on : 2021-04-26
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2893 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metabotropic glutamate receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	0	0
			2833	1841	467	504	21			


- Molecule 2 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: $C_{17}H_{21}N_4O_9P$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
3	A	1	29	21	2	5	1	0	0

- Molecule 1: Metabotropic glutamate receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	58.50Å 86.44Å 99.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.59 – 2.70 43.59 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (43.59-2.70) 97.6 (43.59-2.70)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.19.2	Depositor
R, R_{free}	0.258 , 0.286 0.259 , 0.288	Depositor DCC
R_{free} test set	746 reflections (4.58%)	wwPDB-VP
Wilson B-factor (Å ²)	85.0	Xtriage
Anisotropy	0.296	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 57.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2893	wwPDB-VP
Average B, all atoms (Å ²)	96.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: J9U, FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	0/2894	1.43	11/3950 (0.3%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1071	ILE	N-CA-CB	6.02	115.14	110.45
1	A	1118	ILE	CA-C-N	5.90	128.55	120.46
1	A	1118	ILE	C-N-CA	5.90	128.55	120.46
1	A	721	CYS	CA-C-N	5.64	128.15	120.54
1	A	721	CYS	C-N-CA	5.64	128.15	120.54
1	A	1046	PHE	CA-CB-CG	5.59	119.39	113.80
1	A	637	LEU	CA-C-N	5.43	126.00	119.98
1	A	637	LEU	C-N-CA	5.43	126.00	119.98
1	A	638	GLY	N-CA-C	5.26	119.05	112.73
1	A	730	GLY	CA-C-N	5.04	127.03	120.28
1	A	730	GLY	C-N-CA	5.04	127.03	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2833	0	2785	17	0
2	A	31	0	19	0	0
3	A	29	0	0	0	0
All	All	2893	0	2804	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:658:ALA:O	1:A:662:GLY:HA3	1.83	0.78
1:A:1053:LEU:HD12	1:A:1088:ALA:HB3	1.72	0.71
1:A:609:LEU:HD11	1:A:805:VAL:HG11	1.85	0.58
1:A:661:PHE:CD2	1:A:661:PHE:N	2.73	0.57
1:A:784:SER:HA	1:A:790:GLN:NE2	2.26	0.51
1:A:775:ALA:O	1:A:778:PRO:HD2	2.13	0.48
1:A:1052:VAL:HG13	1:A:1087:VAL:HG12	1.97	0.47
1:A:661:PHE:H	1:A:661:PHE:HD2	1.63	0.46
1:A:658:ALA:O	1:A:662:GLY:CA	2.60	0.46
1:A:742:CYS:HB3	1:A:766:MET:HE2	1.99	0.44
1:A:1094:ASP:HB3	1:A:1097:TRP:HD1	1.82	0.44
1:A:776:PHE:HB3	1:A:780:PHE:CE2	2.55	0.42
1:A:581:ALA:O	1:A:585:VAL:HG23	2.19	0.42
1:A:691:LEU:HA	1:A:694:VAL:HG12	2.02	0.41
1:A:1053:LEU:CD1	1:A:1088:ALA:HB3	2.44	0.41
1:A:660:ILE:O	1:A:672:ARG:NH2	2.52	0.40
1:A:654:THR:CG2	1:A:745:TYR:HB3	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/446 (86%)	368 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/360 (78%)	270 (96%)	12 (4%)	26	54

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	565	ASP
1	A	580	LEU
1	A	1002	LYS
1	A	1032	VAL
1	A	1080	THR
1	A	1130	ARG
1	A	1135	ASP
1	A	672	ARG
1	A	694	VAL
1	A	777	LEU
1	A	784	SER
1	A	806	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1120	GLN
1	A	723	HIS
1	A	790	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FMN	A	1201	-	33,33,33	0.41	0	48,50,50	0.71	1 (2%)
3	J9U	A	1202	-	30,32,32	2.08	11 (36%)	38,47,47	1.16	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	1201	-	-	5/18/18/18	0/3/3/3
3	J9U	A	1202	-	-	3/12/25/25	0/4/4/4

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1202	J9U	C17-N4	6.76	1.44	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1202	J9U	C13-C2	2.57	1.45	1.41
3	A	1202	J9U	C15-C14	2.51	1.56	1.51
3	A	1202	J9U	C4-N1	2.50	1.37	1.33
3	A	1202	J9U	C7-C12	2.43	1.43	1.39
3	A	1202	J9U	C15-N2	2.32	1.51	1.47
3	A	1202	J9U	F-C10	-2.27	1.30	1.36
3	A	1202	J9U	C9-C10	2.26	1.41	1.37
3	A	1202	J9U	C18-C17	2.21	1.41	1.37
3	A	1202	J9U	C2-N	2.15	1.37	1.34
3	A	1202	J9U	C16-C17	2.05	1.52	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1202	J9U	C19-N3-N4	3.81	105.97	104.41
2	A	1201	FMN	C4'-C3'-C2'	2.96	118.49	113.57
3	A	1202	J9U	O-C4-C3	2.30	121.58	119.65

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1201	FMN	C1'-C2'-C3'-C4'
2	A	1201	FMN	O2'-C2'-C3'-C4'
2	A	1201	FMN	O2'-C2'-C3'-O3'
2	A	1201	FMN	C1'-C2'-C3'-O3'
3	A	1202	J9U	C17-C16-N2-C1
3	A	1202	J9U	N-C3-C4-N1
3	A	1202	J9U	N-C3-C4-O
2	A	1201	FMN	C2'-C3'-C4'-O4'


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	387/446 (86%)	0.71	32 (8%) 17 14	70, 95, 118, 137	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	724	ARG	3.8
1	A	593	ASN	3.8
1	A	727	SER	3.6
1	A	1061	ASP	3.2
1	A	725	ASP	3.2
1	A	1048	GLY	3.2
1	A	1127	GLY	3.2
1	A	1129	PRO	3.0
1	A	755	ASN	2.9
1	A	1064	ILE	2.8
1	A	804	VAL	2.8
1	A	726	ALA	2.7
1	A	566	ALA	2.6
1	A	594	ALA	2.5
1	A	798	VAL	2.5
1	A	738	LEU	2.4
1	A	1128	ASP	2.4
1	A	1046	PHE	2.4
1	A	634	LEU	2.4
1	A	602	GLY	2.4
1	A	682	ILE	2.3
1	A	623	PHE	2.3
1	A	807	GLY	2.3
1	A	825	VAL	2.3
1	A	776	PHE	2.3
1	A	756	PHE	2.2
1	A	1078	GLU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	808	CYS	2.1
1	A	631	VAL	2.1
1	A	1100	PHE	2.1
1	A	1132	ALA	2.1
1	A	744	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FMN	A	1201	31/31	0.88	0.10	95,101,107,108	0
3	J9U	A	1202	29/29	0.90	0.12	82,92,111,112	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FMN A 1201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.