



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 04:03 PM EDT

PDB ID : 3EQI / pdb_00003eqi
Title : X-ray structure of the human mitogen-activated protein kinase kinase 1 (MEK1) in a binary complex with ADP and MG2P
Authors : Fischmann, T.O.
Deposited on : 2008-09-30
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2626 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dual specificity mitogen-activated protein kinase kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	315	2466	1575	420	455	16	0	0	2

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	GLY	-	expression tag	UNP Q02750
A	298	ASN	SER	engineered mutation	UNP Q02750
A	299	LYS	SER	engineered mutation	UNP Q02750
A	300	PHE	TYR	engineered mutation	UNP Q02750

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	27	10	5	10	2	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	130	Total	O	0	0
			130	130		

4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	76.89Å 76.89Å 222.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.90 50.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	5.0 (50.00-1.90) 93.7 (50.00-1.90)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 1.91Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.159 , (Not available) 0.221 , 0.263	Depositor DCC
R_{free} test set	1483 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.9	Xtriage
Anisotropy	0.750	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2626	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, CA, MG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2514	0.78	1/3384 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	60	VAL	N-CA-C	9.47	119.43	110.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2466	0	2498	80	0
2	A	27	0	12	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	1	0	0	0	0
6	A	130	0	0	1	0
All	All	2626	0	2510	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:LEU:HD22	1:A:103:ILE:HD11	1.32	1.06
1:A:79:GLY:HA2	1:A:223:PHE:HD2	1.27	1.00
1:A:79:GLY:HA2	1:A:223:PHE:CD2	2.12	0.83
1:A:103:ILE:CG2	1:A:107:ILE:HD12	2.10	0.81
1:A:103:ILE:HG22	1:A:107:ILE:HD12	1.65	0.76
1:A:270:GLU:O	1:A:274:MET:HG3	1.88	0.73
1:A:108:ARG:HD3	1:A:134:TYR:CG	2.25	0.72
1:A:308:MET:HE1	1:A:316:TYR:HB2	1.72	0.70
1:A:308:MET:HE2	1:A:312:GLU:C	2.17	0.69
1:A:110:GLN:HA	1:A:113:ARG:NH1	2.10	0.67
1:A:238:THR:HG22	1:A:239:HIS:H	1.59	0.67
1:A:73:GLU:HG2	1:A:83:PHE:CE2	2.30	0.66
1:A:379:ILE:HD11	1:A:381:LEU:HD12	1.79	0.65
1:A:70:LYS:NZ	1:A:83:PHE:HE2	1.97	0.63
1:A:189:ARG:HD3	1:A:240:TYR:OH	1.99	0.63
1:A:320:GLU:O	1:A:344:LYS:NZ	2.29	0.62
1:A:127:VAL:HG21	1:A:207:CYS:HB3	1.81	0.62
1:A:127:VAL:HG21	1:A:197:LEU:HD12	1.82	0.61
1:A:127:VAL:HG23	1:A:206:LEU:O	2.01	0.60
1:A:58:GLN:CA	1:A:58:GLN:HE21	2.14	0.60
1:A:134:TYR:OH	1:A:137:GLY:HA2	2.01	0.59
1:A:238:THR:HG22	1:A:239:HIS:N	2.18	0.59
1:A:242:VAL:HG21	1:A:352:LEU:HD11	1.85	0.58
1:A:108:ARG:HD3	1:A:134:TYR:CD1	2.39	0.58
1:A:104:LYS:HG3	1:A:107:ILE:HG13	1.84	0.57
1:A:188:HIS:O	1:A:189:ARG:HB2	2.05	0.57
1:A:79:GLY:CA	1:A:223:PHE:HD2	2.11	0.57
1:A:160:ARG:NH1	1:A:274:MET:O	2.37	0.57
1:A:70:LYS:HZ2	1:A:83:PHE:HE2	1.52	0.57
1:A:88:LYS:HB2	1:A:89:PRO:HD3	1.86	0.57
1:A:103:ILE:CD1	1:A:218:SER:HB3	2.35	0.56
1:A:260:ARG:NH1	1:A:266:PRO:HG3	2.21	0.56
1:A:201:ARG:HD3	6:A:510:HOH:O	2.07	0.54
1:A:267:ASP:N	1:A:270:GLU:HG3	2.22	0.54
1:A:49:ARG:HG2	1:A:201:ARG:HH22	1.72	0.54
1:A:47:ARG:NH1	1:A:47:ARG:HG2	2.22	0.53
1:A:260:ARG:CD	1:A:274:MET:HE1	2.38	0.53
1:A:47:ARG:CG	1:A:47:ARG:HH11	2.22	0.53
1:A:260:ARG:HD3	1:A:274:MET:HE1	1.91	0.52
1:A:223:PHE:O	1:A:224:VAL:HG23	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:LYS:HE3	1:A:83:PHE:CD2	2.46	0.51
1:A:223:PHE:CD1	1:A:223:PHE:N	2.78	0.51
1:A:184:HIS:O	1:A:186:ILE:HG13	2.11	0.50
1:A:79:GLY:O	1:A:99:ILE:HG23	2.13	0.49
1:A:112:ILE:O	1:A:116:GLN:HG2	2.12	0.49
1:A:128:GLY:O	1:A:143:MET:HB3	2.13	0.48
1:A:219:MET:O	1:A:223:PHE:HB2	2.13	0.48
1:A:47:ARG:HG2	1:A:47:ARG:HH11	1.79	0.48
1:A:50:LEU:HD13	1:A:50:LEU:HA	1.75	0.48
1:A:70:LYS:NZ	1:A:83:PHE:CE2	2.80	0.47
1:A:272:GLU:O	1:A:276:GLY:HA2	2.15	0.47
1:A:242:VAL:HG21	1:A:352:LEU:CD1	2.45	0.46
1:A:110:GLN:HA	1:A:113:ARG:HH11	1.80	0.46
1:A:272:GLU:O	1:A:272:GLU:HG2	2.15	0.45
1:A:127:VAL:CG2	1:A:207:CYS:HB3	2.44	0.45
1:A:179:TYR:CE2	1:A:183:LYS:HD2	2.52	0.45
1:A:308:MET:HE2	1:A:312:GLU:O	2.17	0.45
1:A:70:LYS:CE	1:A:83:PHE:CE2	3.00	0.44
1:A:47:ARG:NH1	1:A:47:ARG:CG	2.79	0.44
1:A:114:GLU:HB2	1:A:212:SER:HB2	1.99	0.43
1:A:104:LYS:H	1:A:104:LYS:HG2	1.47	0.43
1:A:103:ILE:HD13	1:A:218:SER:HB3	2.00	0.43
1:A:105:PRO:O	1:A:109:ASN:ND2	2.51	0.43
1:A:238:THR:CG2	1:A:239:HIS:H	2.24	0.43
1:A:60:VAL:HG12	1:A:61:GLY:O	2.19	0.42
1:A:110:GLN:O	1:A:114:GLU:HG3	2.18	0.42
1:A:209:PHE:O	1:A:211:VAL:HG23	2.20	0.42
1:A:265:PRO:HA	1:A:266:PRO:HD3	1.78	0.42
1:A:103:ILE:HD11	1:A:218:SER:HB3	2.02	0.42
1:A:233:GLU:HG3	1:A:239:HIS:O	2.20	0.42
1:A:308:MET:HG2	1:A:312:GLU:HB2	2.02	0.42
1:A:134:TYR:HD2	1:A:139:ILE:HG12	1.84	0.42
1:A:345:ASN:HA	1:A:346:PRO:HD2	1.84	0.42
1:A:233:GLU:HG2	1:A:234:ARG:N	2.35	0.42
1:A:168:LYS:HD3	1:A:168:LYS:HA	1.83	0.41
1:A:188:HIS:CD2	1:A:209:PHE:HB3	2.55	0.41
1:A:215:LEU:O	1:A:219:MET:HB2	2.20	0.41
1:A:216:ILE:HG22	1:A:217:ASP:N	2.36	0.41
1:A:100:HIS:O	1:A:223:PHE:HZ	2.02	0.41
1:A:247:TRP:CD1	1:A:247:TRP:C	2.98	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/360 (86%)	296 (95%)	14 (4%)	1 (0%)	36	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	221	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/310 (88%)	245 (90%)	27 (10%)	7	3

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	LEU
1	A	43	ASP
1	A	47	ARG
1	A	48	LYS
1	A	50	LEU
1	A	58	GLN
1	A	78	ASN
1	A	88	LYS
1	A	98	LEU
1	A	103	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	104	LYS
1	A	113	ARG
1	A	138	GLU
1	A	139	ILE
1	A	143	MET
1	A	156	LYS
1	A	181	ARG
1	A	187	MET
1	A	201	ARG
1	A	216	ILE
1	A	219	MET
1	A	223	PHE
1	A	260	ARG
1	A	270	GLU
1	A	313	LEU
1	A	324	LYS
1	A	344	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	GLN
1	A	58	GLN
1	A	78	ASN
1	A	116	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADP	A	2	3	28,29,29	1.08	3 (10%)	43,45,45	1.48	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	2	3	-	2/16/32/32	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2	ADP	C8-N7	2.61	1.36	1.31
2	A	2	ADP	C2-N1	2.47	1.38	1.33
2	A	2	ADP	C6-N6	-2.46	1.27	1.34

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2	ADP	C2-N1-C6	5.04	127.01	118.73
2	A	2	ADP	N3-C2-N1	-3.60	123.13	128.58
2	A	2	ADP	O2A-PA-O3A	3.02	115.45	107.27
2	A	2	ADP	C5-C6-N1	-2.81	110.38	117.51
2	A	2	ADP	C5-C6-N6	2.69	129.95	123.29
2	A	2	ADP	C6-C5-C4	2.27	120.28	117.18

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2	ADP	C5'-O5'-PA-O1A
2	A	2	ADP	PA-O3A-PB-O3B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	315/360 (87%)	0.88	48 (15%) 5 5	23, 43, 79, 96	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	CYS	6.1
1	A	223	PHE	6.0
1	A	239	HIS	5.7
1	A	79	GLY	5.3
1	A	382	ASN	5.0
1	A	78	ASN	4.0
1	A	43	ASP	3.9
1	A	189	ARG	3.6
1	A	210	GLY	3.5
1	A	275	PHE	3.4
1	A	101	LEU	3.4
1	A	220	ALA	3.4
1	A	240	TYR	3.3
1	A	219	MET	3.3
1	A	224	VAL	3.2
1	A	237	GLY	3.2
1	A	103	ILE	2.9
1	A	40	LEU	2.8
1	A	39	GLU	2.8
1	A	77	GLY	2.8
1	A	328	GLY	2.7
1	A	48	LYS	2.7
1	A	268	ALA	2.5
1	A	216	ILE	2.5
1	A	186	ILE	2.4
1	A	42	LEU	2.4
1	A	273	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	272	GLU	2.3
1	A	381	LEU	2.3
1	A	271	LEU	2.3
1	A	76	ALA	2.2
1	A	134	TYR	2.2
1	A	222	SER	2.2
1	A	117	VAL	2.2
1	A	238	THR	2.2
1	A	217	ASP	2.1
1	A	71	ILE	2.1
1	A	55	THR	2.1
1	A	225	GLY	2.1
1	A	347	ALA	2.1
1	A	267	ASP	2.1
1	A	104	LYS	2.0
1	A	183	LYS	2.0
1	A	343	ILE	2.0
1	A	137	GLY	2.0
1	A	106	ALA	2.0
1	A	100	HIS	2.0
1	A	236	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ADP	A	2	27/27	0.85	0.12	33,47,58,59	0
5	NA	A	6	1/1	0.97	0.15	31,31,31,31	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CA	A	5	1/1	0.98	0.07	37,37,37,37	1
3	MG	A	3	1/1	0.99	0.07	38,38,38,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.