



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 11:37 PM UTC

PDB ID : 5EQA / pdb\_00005eqa  
Title : Crystal structure of *Medicago truncatula* Histidinol-Phosphate Phosphatase (MtHPP) with intermolecular cross-link between Lys158 and Cys245  
Authors : Ruszkowski, M.; Dauter, Z.  
Deposited on : 2015-11-12  
Resolution : 1.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2531 (1.34-1.30)
Clashscore	190562	2585 (1.34-1.30)
Ramachandran outliers	187476	2528 (1.34-1.30)
Sidechain outliers	187428	2528 (1.34-1.30)
RSRZ outliers	180081	2528 (1.34-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	 86% 5% • 8%
1	B	277	 84% 6% • 9%
1	C	277	 82% 10% 8%
1	D	277	 82% 9% 8%

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 9078 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inositol monophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	254	2003	1288	331	379	5	0	7	0
1	B	253	1994	1281	329	379	5	0	7	0
1	C	256	2000	1282	333	380	5	0	4	0
1	D	254	1997	1278	334	380	5	0	4	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	SER	-	expression tag	UNP G7J7Q5
A	51	ASN	-	expression tag	UNP G7J7Q5
A	52	ALA	-	expression tag	UNP G7J7Q5
B	50	SER	-	expression tag	UNP G7J7Q5
B	51	ASN	-	expression tag	UNP G7J7Q5
B	52	ALA	-	expression tag	UNP G7J7Q5
C	50	SER	-	expression tag	UNP G7J7Q5
C	51	ASN	-	expression tag	UNP G7J7Q5
C	52	ALA	-	expression tag	UNP G7J7Q5
D	50	SER	-	expression tag	UNP G7J7Q5
D	51	ASN	-	expression tag	UNP G7J7Q5
D	52	ALA	-	expression tag	UNP G7J7Q5

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total 3	Cl 3	0	0
2	B	3	Total 3	Cl 3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total 1	Cl 1	0	0
2	D	2	Total 2	Cl 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	308	Total 309	O 309	0	1
3	B	312	Total 315	O 315	0	3
3	C	211	Total 213	O 213	0	2
3	D	238	Total 238	O 238	0	0


### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Inositol monophosphatase




- Molecule 1: Inositol monophosphatase



- Molecule 1: Inositol monophosphatase



- Molecule 1: Inositol monophosphatase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.24Å 89.39Å 92.24Å 90.00° 96.93° 90.00°	Depositor
Resolution (Å)	36.20 – 1.32 36.20 – 1.32	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.20-1.32) 99.9 (36.20-1.32)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.11 (at 1.32Å)	Xtrriage
Refinement program	REFMAC 5.8.0131	Depositor
R, $R_{free}$	0.124 , 0.165 0.124 , 0.164	Depositor DCC
$R_{free}$ test set	2342 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.6	Xtrriage
Anisotropy	0.679	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	9078	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MLZ, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.35	6/2055 (0.3%)	1.12	3/2793 (0.1%)
1	B	1.34	3/2046 (0.1%)	1.13	0/2782
1	C	1.25	3/2046 (0.1%)	1.22	9/2780 (0.3%)
1	D	1.31	6/2040 (0.3%)	1.18	5/2771 (0.2%)
All	All	1.31	18/8187 (0.2%)	1.16	17/11126 (0.2%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	205[A]	ARG	CA-C	8.44	1.63	1.52
1	D	205[B]	ARG	CA-C	8.44	1.63	1.52
1	C	127	GLY	N-CA	7.70	1.52	1.44
1	D	240	ILE	CA-CB	7.41	1.63	1.54
1	A	146	ASP	C-O	-7.27	1.19	1.25
1	D	316	ILE	CA-CB	6.41	1.62	1.54
1	D	115	VAL	CA-CB	5.88	1.61	1.54
1	C	114[A]	SER	N-CA	5.70	1.53	1.46
1	C	114[B]	SER	N-CA	5.70	1.53	1.46
1	B	126	TYR	C-O	-5.50	1.17	1.23
1	A	63	LEU	C-O	5.48	1.30	1.24
1	D	130	LYS	CA-C	-5.45	1.45	1.52
1	A	316	ILE	CA-CB	5.37	1.60	1.54
1	A	245	CYS	C-O	-5.28	1.17	1.24
1	B	176	GLY	N-CA	5.25	1.52	1.45
1	A	308	VAL	C-O	5.24	1.29	1.24
1	B	248	TYR	CA-C	5.12	1.59	1.52
1	A	202	VAL	C-O	5.10	1.29	1.23

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	245	CYS	N-CA-C	15.71	133.66	112.30
1	D	245	CYS	N-CA-C	15.11	132.85	112.30
1	D	244	GLY	N-CA-C	13.63	134.49	115.27
1	C	244	GLY	N-CA-C	12.58	133.00	115.27
1	C	114[A]	SER	N-CA-C	8.19	124.88	111.37
1	C	114[B]	SER	N-CA-C	8.19	124.88	111.37
1	A	213	ALA	N-CA-C	7.55	121.15	110.50
1	C	243	TYR	N-CA-C	7.28	120.47	109.41
1	C	225	GLY	N-CA-C	6.97	120.59	112.50
1	C	129	GLU	N-CA-C	6.79	118.37	110.97
1	D	243	TYR	N-CA-C	6.17	118.78	109.41
1	D	205[A]	ARG	CA-C-O	5.95	128.30	121.28
1	D	205[B]	ARG	CA-C-O	5.95	128.30	121.28
1	A	314	LYS	N-CA-C	5.78	119.15	111.75
1	A	130	LYS	N-CA-C	5.60	120.10	112.88
1	C	208	ALA	N-CA-C	5.47	117.24	111.28
1	C	246	ASP	N-CA-C	5.12	116.09	108.31

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2003	0	1999	11	0
1	B	1994	0	1987	15	0
1	C	2000	0	1986	16	0
1	D	1997	0	1974	16	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
2	C	1	0	0	0	0
2	D	2	0	0	0	0
3	A	309	0	0	2	0
3	B	315	0	0	10	0
3	C	213	0	0	0	0
3	D	238	0	0	3	0
All	All	9078	0	7946	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:214:TYR:CE2	1:C:240:ILE:HD11	2.04	0.92
1:B:218[B]:THR:HG21	3:B:712[B]:HOH:O	1.86	0.74
1:A:284[B]:VAL:HG21	1:A:314:LYS:CD	2.25	0.66
1:B:129:GLU:HG2	3:B:724:HOH:O	2.00	0.62
1:B:191[B]:THR:HG21	3:B:575:HOH:O	2.00	0.60
1:C:214:TYR:CZ	1:C:240:ILE:HD11	2.36	0.59
1:A:284[B]:VAL:HG21	1:A:314:LYS:HD3	1.85	0.58
1:D:207:CYS:SG	3:D:684:HOH:O	2.56	0.58
1:B:207:CYS:O	1:B:315:GLN:NE2	2.36	0.57
1:B:314:LYS:HE3	3:B:767:HOH:O	2.04	0.57
1:D:191[B]:THR:HG22	3:D:508:HOH:O	2.05	0.56
1:B:126:TYR:HE1	3:B:551:HOH:O	1.88	0.55
1:A:284[B]:VAL:CG2	1:A:314:LYS:HD3	2.37	0.55
1:A:207:CYS:HB3	1:A:316:ILE:HG13	1.88	0.54
1:B:191[B]:THR:HG22	3:B:519:HOH:O	2.07	0.54
1:A:284[B]:VAL:CG2	1:A:314:LYS:CD	2.85	0.54
1:C:214:TYR:CE2	1:C:240:ILE:CD1	2.85	0.54
1:B:218[A]:THR:HG23	3:B:540:HOH:O	2.08	0.53
1:B:284[B]:VAL:HG21	1:B:314:LYS:HG3	1.90	0.53
1:B:99:SER:HB2	3:B:755:HOH:O	2.09	0.53
1:C:216:TYR:CE1	1:D:158:MLZ:HCM2	2.45	0.52
1:A:284[B]:VAL:HG21	1:A:314:LYS:HD2	1.91	0.51
1:D:239:LYS:HG2	1:D:240:ILE:HG12	1.93	0.51
1:C:239:LYS:HG2	1:C:240:ILE:HD13	1.94	0.49
1:A:214:TYR:CE1	1:A:240:ILE:HD12	2.49	0.48
1:C:152:LYS:HE3	1:D:240:ILE:CD1	2.44	0.48
1:D:128:GLU:N	3:D:503:HOH:O	2.47	0.47
1:B:236:ASP:CG	3:B:502:HOH:O	2.58	0.46
1:A:284[B]:VAL:CG2	1:A:314:LYS:HD2	2.44	0.46
1:C:158:MLZ:HCM2	1:D:216:TYR:CE1	2.52	0.45
1:D:239:LYS:HG2	1:D:240:ILE:CG1	2.46	0.45
1:B:110:GLU:CD	3:B:501:HOH:O	2.60	0.44
1:C:224:SER:O	1:C:228:GLU:HB2	2.18	0.43
1:D:263[B]:GLU:HG2	1:D:266:LEU:HG	2.00	0.43
1:D:284:VAL:HG21	1:D:314:LYS:HG2	2.00	0.43
1:D:126:TYR:CD1	1:D:126:TYR:C	2.97	0.43
1:A:130:LYS:HA	3:A:503:HOH:O	2.18	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:263[B]:GLU:HG3	3:A:636:HOH:O	2.18	0.43
1:B:205:ARG:NH2	1:B:214:TYR:CD2	2.77	0.43
1:C:311:ALA:HB3	1:C:317:HIS:HB2	2.00	0.43
1:D:275:ILE:HB	1:D:276:PRO:HD3	2.00	0.42
1:B:174:ILE:HA	1:B:191[B]:THR:CG2	2.50	0.42
1:C:152:LYS:HE3	1:D:240:ILE:HD13	2.00	0.42
1:D:239:LYS:C	1:D:240:ILE:HG12	2.44	0.42
1:C:175:LEU:HD23	1:C:175:LEU:C	2.44	0.42
1:A:183:LEU:HD22	1:B:183:LEU:HD22	2.00	0.41
1:C:224:SER:O	1:C:228:GLU:N	2.46	0.41
1:C:183:LEU:HD22	1:D:183:LEU:HD22	2.03	0.41
1:C:214:TYR:CD2	1:C:240:ILE:HD11	2.54	0.41
1:D:210:LEU:O	1:D:238:VAL:HA	2.21	0.40
1:C:294:ARG:HH11	1:C:294:ARG:HG2	1.86	0.40
1:C:146:ASP:OD2	1:C:270:ASP:OD1	2.39	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/277 (92%)	250 (98%)	6 (2%)	0	100	100
1	B	255/277 (92%)	248 (97%)	7 (3%)	0	100	100
1	C	255/277 (92%)	249 (98%)	6 (2%)	0	100	100
1	D	253/277 (91%)	244 (96%)	8 (3%)	1 (0%)	30	9
All	All	1019/1108 (92%)	991 (97%)	27 (3%)	1 (0%)	48	18

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	225	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/232 (94%)	214 (99%)	3 (1%)	59	22
1	B	216/232 (93%)	212 (98%)	4 (2%)	50	14
1	C	216/232 (93%)	211 (98%)	5 (2%)	44	9
1	D	214/232 (92%)	212 (99%)	2 (1%)	70	40
All	All	863/928 (93%)	849 (98%)	14 (2%)	53	18

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	LYS
1	A	180	GLN
1	A	314	LYS
1	B	87	LYS
1	B	103	ILE
1	B	314	LYS
1	B	315	GLN
1	C	98	LEU
1	C	135	LYS
1	C	136	GLN
1	C	240	ILE
1	C	314	LYS
1	D	63	LEU
1	D	240	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	136	GLN
1	C	136	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	64	ASN
1	D	212	GLN
1	D	292	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	MLZ	B	158	1	8,9,10	1.04	0	4,9,11	0.39	0
1	MLZ	A	158	1	8,9,10	1.02	1 (12%)	4,9,11	0.80	0
1	MLZ	C	158	1	8,9,10	1.16	0	4,9,11	0.61	0
1	MLZ	D	158	1	8,9,10	0.51	0	4,9,11	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLZ	B	158	1	-	0/7/8/10	-
1	MLZ	A	158	1	-	0/7/8/10	-
1	MLZ	C	158	1	-	0/7/8/10	-
1	MLZ	D	158	1	-	0/7/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	158	MLZ	CB-CA	-2.33	1.50	1.53

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	158	MLZ	1	0
1	D	158	MLZ	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	253/277 (91%)	-0.16	2 (0%) 82 87	7, 17, 37, 60	7 (2%)
1	B	252/277 (90%)	-0.16	3 (1%) 76 81	7, 16, 35, 50	7 (2%)
1	C	255/277 (92%)	0.34	17 (6%) 24 30	12, 23, 48, 78	4 (1%)
1	D	253/277 (91%)	0.13	13 (5%) 33 42	10, 20, 42, 59	4 (1%)
All	All	1013/1108 (91%)	0.04	35 (3%) 47 55	7, 19, 41, 78	22 (2%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	63	LEU	5.6
1	C	98	LEU	5.5
1	C	223	PHE	5.1
1	D	63	LEU	4.2
1	C	231	PHE	4.1
1	C	224	SER	4.1
1	D	244	GLY	3.6
1	C	63	LEU	3.6
1	C	222	LEU	3.5
1	C	218	THR	3.4
1	D	222	LEU	3.4
1	D	243	TYR	3.4
1	C	244	GLY	3.2
1	D	223	PHE	3.1
1	C	217	THR	3.0
1	A	208	ALA	2.9
1	C	216	TYR	2.9
1	B	208	ALA	2.9
1	B	64	ASN	2.7
1	C	135	LYS	2.6
1	C	240	ILE	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	132	TRP	2.6
1	C	129	GLU	2.5
1	C	243	TYR	2.5
1	A	63	LEU	2.5
1	D	220	PRO	2.5
1	D	225	GLY	2.4
1	D	224	SER	2.4
1	D	297	ALA	2.3
1	C	221	HIS	2.2
1	D	231	PHE	2.2
1	C	208	ALA	2.2
1	D	219	SER	2.1
1	D	240	ILE	2.0
1	D	221	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	MLZ	C	158	10/11	0.97	0.06	16,17,21,21	0
1	MLZ	B	158	10/11	0.98	0.04	10,12,12,13	0
1	MLZ	D	158	10/11	0.98	0.06	14,16,21,23	0
1	MLZ	A	158	10/11	0.99	0.04	11,12,12,13	0

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CL	A	402	1/1	0.99	0.13	29,29,29,29	0
2	CL	B	401	1/1	0.99	0.10	26,26,26,26	0
2	CL	B	402	1/1	0.99	0.12	29,29,29,29	0
2	CL	C	401	1/1	0.99	0.09	31,31,31,31	0
2	CL	D	401	1/1	0.99	0.10	29,29,29,29	0
2	CL	D	402	1/1	0.99	0.06	33,33,33,33	0
2	CL	A	401	1/1	1.00	0.07	32,32,32,32	0
2	CL	A	403	1/1	1.00	0.09	20,20,20,20	0
2	CL	B	403	1/1	1.00	0.12	22,22,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.