



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 05:06 PM EDT

PDB ID : 5ERP / pdb_00005erp
Title : Crystal structure of human Desmocollin-2 ectodomain fragment EC2-5
Authors : Harrison, O.J.; Brasch, J.; Shapiro, L.
Deposited on : 2015-11-14
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	11% 86% 11% ..
1	B	451	3% 85% 12% ..
2	C	4	75% 25%
2	D	4	50% 50%
2	E	4	50% 50%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 14287 atoms, of which 6991 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Desmocollin-2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	443	Total	C	H	N	O	S	0	0	0
			6802	2140	3358	575	710	19			
1	B	438	Total	C	H	N	O	S	0	0	0
			6733	2121	3323	569	701	19			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	546	HIS	-	expression tag	UNP Q02487
A	547	HIS	-	expression tag	UNP Q02487
A	548	HIS	-	expression tag	UNP Q02487
A	549	HIS	-	expression tag	UNP Q02487
A	550	HIS	-	expression tag	UNP Q02487
A	551	HIS	-	expression tag	UNP Q02487
B	546	HIS	-	expression tag	UNP Q02487
B	547	HIS	-	expression tag	UNP Q02487
B	548	HIS	-	expression tag	UNP Q02487
B	549	HIS	-	expression tag	UNP Q02487
B	550	HIS	-	expression tag	UNP Q02487
B	551	HIS	-	expression tag	UNP Q02487

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	4	Total	C	H	N	O	0	0	0
			100	28	50	2	20			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	4	Total	C	H	N	O	0	0	0
			100	28	50	2	20			
2	E	4	Total	C	H	N	O	0	0	0
			100	28	50	2	20			

- Molecule 3 is alpha-D-mannopyranose (CCD ID: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			22	6	11	5		
3	A	1	Total	C	H	O	0	0
			22	6	11	5		
3	A	1	Total	C	H	O	0	0
			22	6	11	5		
3	A	1	Total	C	H	O	0	0
			22	6	11	5		
3	B	1	Total	C	H	O	0	0
			22	6	11	5		
3	B	1	Total	C	H	O	0	0
			22	6	11	5		
3	B	1	Total	C	H	O	0	0
			22	6	11	5		
3	B	1	Total	C	H	O	0	0
			22	6	11	5		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
4	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
4	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	10	Total	Ca	0	0
			10	10		
5	B	11	Total	Ca	0	0
			11	11		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			10	2	6	2		
7	A	1	Total	C	H	O	0	0
			10	2	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	59	Total	O	0	0
			59	59		
8	B	52	Total	O	0	0
			52	52		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Desmocollin-2



- Molecule 1: Desmocollin-2



- Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  50% 50%

MAG1
MAG2
BGL3
MAN4

- Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50% 50%

MAG1
MAG2
BGL3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	146.22Å 94.17Å 126.31Å 90.00° 93.66° 90.00°	Depositor
Resolution (Å)	79.12 – 2.70 79.12 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.4 (79.12-2.70) 86.7 (79.12-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 2.69Å)	Xtriage
Refinement program	PHENIX dev_1810	Depositor
R, R_{free}	0.224 , 0.262 0.233 , 0.269	Depositor DCC
R_{free} test set	1951 reflections (4.23%)	wwPDB-VP
Wilson B-factor (Å ²)	67.3	Xtriage
Anisotropy	0.402	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14287	wwPDB-VP
Average B, all atoms (Å ²)	110.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, MAN, SO4, BMA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	1/3505 (0.0%)	0.70	3/4778 (0.1%)
1	B	0.34	0/3470	0.68	0/4729
All	All	0.35	1/6975 (0.0%)	0.69	3/9507 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	497	ALA	CA-CB	6.50	1.63	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	310	ARG	N-CA-C	5.35	117.53	111.11
1	A	411	ASN	N-CA-C	5.19	117.70	109.24
1	A	497	ALA	N-CA-C	-5.18	102.02	110.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3444	3358	3355	28	0
1	B	3410	3323	3321	27	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	50	50	43	3	0
2	D	50	50	43	0	0
2	E	50	50	43	2	0
3	A	44	44	39	1	0
3	B	44	44	40	2	0
4	A	28	28	26	1	0
4	B	14	14	13	0	0
5	A	10	0	0	0	0
5	B	11	0	0	0	0
6	A	5	0	0	0	0
6	B	5	0	0	0	0
7	A	12	18	18	0	0
7	B	8	12	11	1	0
8	A	59	0	0	0	0
8	B	52	0	0	0	0
All	All	7296	6991	6952	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:801:MAN:O6	3:A:801:MAN:C6	1.65	1.43
2:E:3:BMA:O2	2:E:4:MAN:O2	2.00	0.76
1:B:336:CYS:N	1:B:424:CYS:SG	2.64	0.71
1:A:454:MET:O	1:A:502:TYR:OH	2.13	0.67
1:B:192:ASP:OD1	1:B:200:LEU:N	2.29	0.65

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:228:SER:O	1:B:489:ARG:NH2[3_555]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/451 (98%)	416 (94%)	25 (6%)	0	100	100
1	B	434/451 (96%)	410 (94%)	24 (6%)	0	100	100
All	All	875/902 (97%)	826 (94%)	49 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	402/410 (98%)	391 (97%)	11 (3%)	39	69
1	B	398/410 (97%)	384 (96%)	14 (4%)	32	61
All	All	800/820 (98%)	775 (97%)	25 (3%)	35	65

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	217	LEU
1	B	273	THR
1	B	527	THR
1	B	272	VAL
1	B	323	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	407	ASN
1	B	299	GLN
1	A	504	ASN
1	A	503	GLN
1	B	196	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.19	0	17,19,21	0.61	0
2	NAG	C	2	2	14,14,15	0.31	0	17,19,21	1.14	2 (11%)
2	BMA	C	3	2	11,11,12	0.67	0	15,15,17	0.70	0
2	MAN	C	4	2	11,11,12	0.64	0	15,15,17	0.91	1 (6%)
2	NAG	D	1	1,2	14,14,15	1.86	3 (21%)	17,19,21	1.05	1 (5%)
2	NAG	D	2	2	14,14,15	0.41	0	17,19,21	0.61	0
2	BMA	D	3	2	11,11,12	0.78	0	15,15,17	0.83	0
2	MAN	D	4	2	11,11,12	0.75	0	15,15,17	0.97	2 (13%)
2	NAG	E	1	1,2	14,14,15	0.31	0	17,19,21	0.62	0
2	NAG	E	2	2	14,14,15	0.24	0	17,19,21	0.56	0
2	BMA	E	3	2	11,11,12	1.49	3 (27%)	15,15,17	1.45	2 (13%)
2	MAN	E	4	2	11,11,12	1.00	1 (9%)	15,15,17	2.03	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	6/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	1/2/19/22	0/1/1/1
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
2	BMA	D	3	2	-	2/2/19/22	0/1/1/1
2	MAN	D	4	2	-	1/2/19/22	0/1/1/1
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1
2	BMA	E	3	2	-	2/2/19/22	0/1/1/1
2	MAN	E	4	2	-	2/2/19/22	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	NAG	O5-C1	5.61	1.53	1.43
2	E	3	BMA	O3-C3	3.02	1.50	1.43
2	D	1	NAG	C1-C2	2.83	1.56	1.52
2	E	4	MAN	C1-C2	2.55	1.58	1.52
2	E	3	BMA	O5-C5	2.14	1.47	1.43

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	4	MAN	C1-C2-C3	4.60	116.34	109.64
2	E	3	BMA	C1-O5-C5	4.04	117.59	112.19
2	E	4	MAN	C1-O5-C5	3.99	117.54	112.19
2	E	4	MAN	O2-C2-C3	-3.82	102.24	110.15
2	C	2	NAG	C2-N2-C7	3.60	127.73	122.90

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	4	MAN	C4-C5-C6-O6
2	E	2	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	O5-C5-C6-O6
2	E	2	NAG	O5-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6


There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	3	BMA	2	0
2	C	1	NAG	1	0
2	C	3	BMA	1	0
2	E	4	MAN	2	0
2	C	2	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 39 ligands modelled in this entry, 21 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	A	801	1	11,11,12	2.07	3 (27%)	15,15,17	2.73	3 (20%)
3	MAN	B	603	1	11,11,12	0.67	0	15,15,17	1.28	3 (20%)
3	MAN	B	604	1	11,11,12	0.94	1 (9%)	15,15,17	1.23	1 (6%)
3	MAN	B	605	1	11,11,12	0.70	0	15,15,17	1.74	4 (26%)
3	MAN	A	804	1	11,11,12	0.90	1 (9%)	15,15,17	0.94	1 (6%)
7	EDO	A	820	-	3,3,3	0.42	0	2,2,2	0.36	0
7	EDO	A	819	-	3,3,3	0.41	0	2,2,2	0.37	0
6	SO4	A	817	-	4,4,4	0.23	0	6,6,6	0.07	0
7	EDO	B	619	-	3,3,3	0.70	0	2,2,2	2.05	1 (50%)
3	MAN	B	602	1	11,11,12	0.66	0	15,15,17	1.20	2 (13%)
3	MAN	A	802	1	11,11,12	0.84	0	15,15,17	1.48	2 (13%)
3	MAN	A	803	1	11,11,12	0.91	0	15,15,17	1.03	1 (6%)
4	NAG	A	805	1	14,14,15	1.08	2 (14%)	17,19,21	1.03	1 (5%)
7	EDO	B	618	-	3,3,3	0.41	0	2,2,2	0.34	0
4	NAG	B	601	1	14,14,15	0.52	0	17,19,21	0.67	1 (5%)
6	SO4	B	617	-	4,4,4	0.24	0	6,6,6	0.09	0
7	EDO	A	818	-	3,3,3	0.41	0	2,2,2	0.48	0
4	NAG	A	806	1	14,14,15	1.63	1 (7%)	17,19,21	1.63	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	A	801	1	-	0/2/19/22	0/1/1/1
3	MAN	B	603	1	-	0/2/19/22	1/1/1/1
3	MAN	B	604	1	-	1/2/19/22	0/1/1/1
3	MAN	B	605	1	-	0/2/19/22	0/1/1/1
3	MAN	A	804	1	-	1/2/19/22	0/1/1/1
7	EDO	A	820	-	-	0/1/1/1	-
7	EDO	A	819	-	-	1/1/1/1	-
7	EDO	B	619	-	-	0/1/1/1	-
3	MAN	B	602	1	-	0/2/19/22	0/1/1/1
3	MAN	A	802	1	-	0/2/19/22	1/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	A	803	1	-	0/2/19/22	0/1/1/1
4	NAG	A	805	1	-	1/6/23/26	0/1/1/1
7	EDO	B	618	-	-	1/1/1/1	-
4	NAG	B	601	1	-	2/6/23/26	0/1/1/1
7	EDO	A	818	-	-	1/1/1/1	-
4	NAG	A	806	1	-	5/6/23/26	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	806	NAG	O5-C1	5.84	1.53	1.43
3	A	801	MAN	O6-C6	5.60	1.65	1.42
4	A	805	NAG	O5-C1	3.14	1.49	1.43
3	B	604	MAN	O5-C5	2.54	1.48	1.43
4	A	805	NAG	C1-C2	2.48	1.55	1.52

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	801	MAN	C1-O5-C5	9.48	124.89	112.19
3	B	605	MAN	C1-O5-C5	4.69	118.48	112.19
3	A	802	MAN	C1-O5-C5	4.59	118.33	112.19
4	A	806	NAG	C2-N2-C7	3.85	128.06	122.90
3	B	604	MAN	C1-O5-C5	3.68	117.12	112.19

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	601	NAG	O5-C5-C6-O6
4	A	806	NAG	C8-C7-N2-C2
4	A	806	NAG	O7-C7-N2-C2
4	B	601	NAG	C4-C5-C6-O6
4	A	806	NAG	O5-C5-C6-O6

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	802	MAN	C1-C2-C3-C4-C5-O5
3	B	603	MAN	C1-C2-C3-C4-C5-O5

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	801	MAN	1	0
3	B	603	MAN	2	0
7	B	619	EDO	1	0
3	B	602	MAN	1	0
4	A	806	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	443/451 (98%)	0.55	48 (10%) 11 9	67, 98, 216, 241	0
1	B	438/451 (97%)	0.17	12 (2%) 56 53	66, 92, 136, 180	0
All	All	881/902 (97%)	0.36	60 (6%) 23 20	66, 95, 205, 241	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	441	PHE	4.6
1	A	528	SER	4.1
1	A	513	VAL	4.0
1	A	543	THR	3.9
1	A	442	ILE	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BMA	E	3	11/12	0.31	0.13	158,185,213,222	0
2	MAN	E	4	11/12	0.34	0.11	164,186,218,222	0
2	BMA	C	3	11/12	0.41	0.12	177,195,231,234	0
2	MAN	C	4	11/12	0.42	0.11	181,206,242,247	0
2	BMA	D	3	11/12	0.49	0.13	145,170,196,200	0

Continued on next page...

Continued from previous page...



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MAN	D	4	11/12	0.61	0.11	150,172,202,207	0
2	NAG	C	2	14/15	0.71	0.15	138,166,193,202	0
2	NAG	C	1	14/15	0.77	0.13	94,124,152,159	0
2	NAG	D	1	14/15	0.79	0.14	82,117,141,157	0
2	NAG	D	2	14/15	0.87	0.10	96,124,161,193	0
2	NAG	E	2	14/15	0.88	0.10	93,130,163,184	0
2	NAG	E	1	14/15	0.91	0.10	72,100,134,151	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	EDO	A	820	4/4	0.46	0.19	98,118,122,133	0
3	MAN	B	603	11/12	0.53	0.14	124,149,179,185	0
3	MAN	B	605	11/12	0.60	0.12	124,149,178,179	0
7	EDO	A	819	4/4	0.62	0.15	135,162,168,168	0
4	NAG	A	806	14/15	0.65	0.10	164,197,226,234	0
3	MAN	B	602	11/12	0.66	0.12	117,146,169,179	0
3	MAN	A	801	11/12	0.66	0.16	114,136,159,176	0
3	MAN	A	804	11/12	0.67	0.12	113,147,171,194	0
4	NAG	A	805	14/15	0.70	0.13	121,149,181,193	0
3	MAN	B	604	11/12	0.74	0.14	116,143,183,187	0
7	EDO	A	818	4/4	0.76	0.29	106,128,132,132	0
3	MAN	A	803	11/12	0.79	0.11	94,119,141,144	0
3	MAN	A	802	11/12	0.80	0.11	108,133,157,166	0
7	EDO	B	619	4/4	0.80	0.20	97,132,142,158	0
6	SO4	A	817	5/5	0.81	0.07	150,154,165,171	0
4	NAG	B	601	14/15	0.82	0.12	99,123,145,155	0
7	EDO	B	618	4/4	0.84	0.10	92,110,126,126	0
6	SO4	B	617	5/5	0.84	0.10	139,141,148,165	0
5	CA	B	613	1/1	0.85	0.21	217,217,217,217	0
5	CA	B	616	1/1	0.91	0.14	119,119,119,119	0
5	CA	A	816	1/1	0.95	0.11	123,123,123,123	0
5	CA	B	606	1/1	0.97	0.09	103,103,103,103	0
5	CA	B	608	1/1	0.97	0.05	91,91,91,91	0
5	CA	A	810	1/1	0.97	0.11	83,83,83,83	0
5	CA	B	615	1/1	0.97	0.16	102,102,102,102	0
5	CA	A	813	1/1	0.98	0.10	119,119,119,119	0
5	CA	B	611	1/1	0.98	0.07	77,77,77,77	0
5	CA	A	814	1/1	0.98	0.11	110,110,110,110	0
5	CA	A	811	1/1	0.99	0.07	72,72,72,72	0
5	CA	A	812	1/1	0.99	0.04	74,74,74,74	0
5	CA	B	607	1/1	0.99	0.04	81,81,81,81	0
5	CA	A	809	1/1	0.99	0.07	76,76,76,76	0
5	CA	B	610	1/1	0.99	0.05	72,72,72,72	0
5	CA	A	807	1/1	0.99	0.14	116,116,116,116	0
5	CA	B	612	1/1	0.99	0.07	92,92,92,92	0
5	CA	A	815	1/1	0.99	0.07	86,86,86,86	0
5	CA	B	614	1/1	0.99	0.08	76,76,76,76	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	CA	A	808	1/1	1.00	0.09	81,81,81,81	0
5	CA	B	609	1/1	1.00	0.02	75,75,75,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.