



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 05:36 AM UTC

PDB ID : 8ER0 / pdb_00008er0
Title : X-ray crystal structure of Tet(X6) bound to anhydrotetracycline
Authors : Kumar, H.; Tolia, N.H.
Deposited on : 2022-10-11
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	403	
1	B	403	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12213 atoms, of which 5905 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flavin-dependent monooxygenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	371	Total	C	H	N	O	S	0	0	0
			5872	1864	2920	510	566	12			
1	B	368	Total	C	H	N	O	S	0	0	0
			5810	1848	2885	503	562	12			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	MET	-	initiating methionine	UNP A0A316WTJ0
A	-2	THR	-	expression tag	UNP A0A316WTJ0
A	-1	SER	-	expression tag	UNP A0A316WTJ0
A	0	SER	-	expression tag	UNP A0A316WTJ0
A	1	SER	-	expression tag	UNP A0A316WTJ0
A	2	HIS	-	expression tag	UNP A0A316WTJ0
A	3	HIS	-	expression tag	UNP A0A316WTJ0
A	4	HIS	-	expression tag	UNP A0A316WTJ0
A	5	HIS	-	expression tag	UNP A0A316WTJ0
A	6	HIS	-	expression tag	UNP A0A316WTJ0
A	7	HIS	-	expression tag	UNP A0A316WTJ0
A	8	SER	-	expression tag	UNP A0A316WTJ0
A	9	SER	-	expression tag	UNP A0A316WTJ0
A	10	GLY	-	expression tag	UNP A0A316WTJ0
A	11	LYS	-	expression tag	UNP A0A316WTJ0
A	12	LEU	-	expression tag	UNP A0A316WTJ0
B	-3	MET	-	initiating methionine	UNP A0A316WTJ0
B	-2	THR	-	expression tag	UNP A0A316WTJ0
B	-1	SER	-	expression tag	UNP A0A316WTJ0
B	0	SER	-	expression tag	UNP A0A316WTJ0
B	1	SER	-	expression tag	UNP A0A316WTJ0
B	2	HIS	-	expression tag	UNP A0A316WTJ0
B	3	HIS	-	expression tag	UNP A0A316WTJ0
B	4	HIS	-	expression tag	UNP A0A316WTJ0
B	5	HIS	-	expression tag	UNP A0A316WTJ0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	6	HIS	-	expression tag	UNP A0A316WTJ0
B	7	HIS	-	expression tag	UNP A0A316WTJ0
B	8	SER	-	expression tag	UNP A0A316WTJ0
B	9	SER	-	expression tag	UNP A0A316WTJ0
B	10	GLY	-	expression tag	UNP A0A316WTJ0
B	11	LYS	-	expression tag	UNP A0A316WTJ0
B	12	LEU	-	expression tag	UNP A0A316WTJ0

- Molecule 2 is 5A,6-ANHYDROTETRACYCLINE (CCD ID: TDC) (formula: $C_{22}H_{22}N_2O_7$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			50	22	19	2	7		
2	B	1	Total	C	H	N	O	0	0
			50	22	19	2	7		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	0	0
			84	27	31	9	15	2		
3	B	1	Total	C	H	N	O	P	0	0
			84	27	31	9	15	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	138	Total	O	0	0
			138	138		
4	B	125	Total	O	0	0
			125	125		

- Molecule 1: Flavin-dependent monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.09Å 52.21Å 94.87Å 90.00° 95.44° 90.00°	Depositor
Resolution (Å)	19.84 – 2.20 19.84 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.6 (19.84-2.20) 98.5 (19.84-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 2.19Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.193 , 0.231 0.194 , 0.231	Depositor DCC
R_{free} test set	2006 reflections (4.60%)	wwPDB-VP
Wilson B-factor (Å ²)	25.6	Xtrriage
Anisotropy	0.590	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12213	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4696e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, TDC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/3008	0.32	0/4066
1	B	0.13	0/2980	0.31	0/4028
All	All	0.13	0/5988	0.32	0/8094

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2952	2920	2919	8	0
1	B	2925	2885	2885	7	0
2	A	31	19	19	0	0
2	B	31	19	19	1	0
3	A	53	31	31	1	0
3	B	53	31	31	1	0
4	A	138	0	0	0	0
4	B	125	0	0	0	0
All	All	6308	5905	5904	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:ILE:HD13	1:A:51:VAL:CG1	2.39	0.53
1:B:335:VAL:HG12	3:B:402:FAD:O2	2.11	0.50
1:B:325:HIS:NE2	1:B:341:ASP:OD2	2.42	0.47
1:B:30:ILE:HD12	1:B:30:ILE:N	2.30	0.47
1:B:206:ILE:HD12	1:B:288:LEU:HD12	1.95	0.47
1:B:325:HIS:O	1:B:326:LEU:C	2.59	0.45
1:A:325:HIS:NE2	1:A:341:ASP:OD2	2.47	0.44
1:A:377:ALA:O	1:A:381:ILE:HG12	2.17	0.44
1:A:206:ILE:HD12	1:A:288:LEU:HD12	2.00	0.43
1:A:152:THR:HB	1:A:163:THR:HB	1.99	0.43
1:A:325:HIS:O	1:A:326:LEU:C	2.61	0.43
1:B:235:PHE:CD1	2:B:401:TDC:H4	2.54	0.43
1:A:255:GLU:O	1:A:259:LYS:N	2.53	0.42
1:A:335:VAL:HG12	3:A:402:FAD:O2	2.21	0.41
1:B:97:PRO:HB2	1:B:115:VAL:HG21	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	369/403 (92%)	356 (96%)	13 (4%)	0	100	100
1	B	364/403 (90%)	351 (96%)	13 (4%)	0	100	100
All	All	733/806 (91%)	707 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/354 (92%)	324 (100%)	0	100	100
1	B	321/354 (91%)	321 (100%)	0	100	100
All	All	645/708 (91%)	645 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	HIS
1	A	366	GLN
1	A	367	GLN
1	B	201	ASN
1	B	203	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TDC	B	401	-	34,34,34	1.19	2 (5%)	43,55,55	0.96	4 (9%)
3	FAD	B	402	-	58,58,58	0.35	0	85,89,89	0.36	0
2	TDC	A	401	-	34,34,34	1.22	2 (5%)	43,55,55	0.95	4 (9%)
3	FAD	A	402	-	58,58,58	0.32	0	85,89,89	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TDC	B	401	-	-	0/8/50/50	0/4/4/4
3	FAD	B	402	-	-	1/34/50/50	0/6/6/6
2	TDC	A	401	-	-	0/8/50/50	0/4/4/4
3	FAD	A	402	-	-	0/34/50/50	0/6/6/6

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	TDC	C21-N21	5.08	1.48	1.33
2	B	401	TDC	C21-N21	4.91	1.47	1.33
2	B	401	TDC	C1C-C1	-2.43	1.51	1.55
2	A	401	TDC	C1C-C1	-2.39	1.51	1.55

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	TDC	O21-C21-N21	-2.49	117.31	122.89
2	B	401	TDC	C41-C1C-C12	2.48	113.89	111.05
2	A	401	TDC	O21-C21-N21	-2.45	117.41	122.89
2	B	401	TDC	C1C-C41-C4	2.42	114.94	111.64
2	A	401	TDC	C41-C1C-C12	2.35	113.75	111.05
2	A	401	TDC	C1C-C41-C4	2.24	114.71	111.64
2	A	401	TDC	C2-C21-N21	2.16	123.05	118.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	TDC	C2-C21-N21	2.05	122.82	118.64

There are no chirality outliers.

All (1) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	B	402	FAD	O4'-C4'-C5'-O5'



There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	TDC	1	0
3	B	402	FAD	1	0
3	A	402	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	371/403 (92%)	-0.03	10 (2%) 56 53	19, 32, 64, 109	0
1	B	368/403 (91%)	-0.08	4 (1%) 78 76	20, 32, 61, 89	0
All	All	739/806 (91%)	-0.05	14 (1%) 66 63	19, 32, 63, 109	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	260	THR	4.2
1	A	259	LYS	3.6
1	A	115	VAL	3.4
1	A	258	SER	3.3
1	A	112	THR	2.7
1	B	258	SER	2.7
1	A	121	PHE	2.7
1	A	390	ASP	2.7
1	B	115	VAL	2.6
1	A	113	LYS	2.5
1	B	263	ASP	2.5
1	B	76	ASP	2.4
1	A	263	ASP	2.3
1	A	114	ASN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TDC	A	401	31/31	0.76	0.15	32,61,71,78	0
2	TDC	B	401	31/31	0.77	0.14	27,63,76,82	0
3	FAD	B	402	53/53	0.96	0.08	17,26,48,64	0
3	FAD	A	402	53/53	0.97	0.07	17,26,46,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around TDC B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.