



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:53 AM UTC

PDB ID : 9ERC / pdb_00009erc
Title : Hydrogenase-2 Ni-Li state
Authors : Wong, K.L.; Carr, S.B.; Ash, P.A.; Vincent, K.A.
Deposited on : 2024-03-22
Resolution : 1.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1553 (1.30-1.30)
Clashscore	190562	1595 (1.30-1.30)
Ramachandran outliers	187476	1551 (1.30-1.30)
Sidechain outliers	187428	1551 (1.30-1.30)
RSRZ outliers	180081	1549 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	S	298	 2% 85% 5% 10%
1	T	298	 2% 86% 5% 10%
2	L	567	 % 92% 5% .
2	M	567	 4% 93% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	L	607	-	-	X	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 14333 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrogenase-2 small chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	S	268	Total 2063	C 1306	N 363	O 381	S 13	0	4	0
1	T	267	Total 2088	C 1323	N 365	O 387	S 13	0	8	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
S	294	HIS	-	expression tag	UNP P69741
S	295	HIS	-	expression tag	UNP P69741
S	296	HIS	-	expression tag	UNP P69741
S	297	HIS	-	expression tag	UNP P69741
S	298	HIS	-	expression tag	UNP P69741
S	299	HIS	-	expression tag	UNP P69741
T	294	HIS	-	expression tag	UNP P69741
T	295	HIS	-	expression tag	UNP P69741
T	296	HIS	-	expression tag	UNP P69741
T	297	HIS	-	expression tag	UNP P69741
T	298	HIS	-	expression tag	UNP P69741
T	299	HIS	-	expression tag	UNP P69741

- Molecule 2 is a protein called Hydrogenase-2 large chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	551	Total 4346	C 2761	N 748	O 819	S 18	0	9	0
2	M	550	Total 4332	C 2754	N 746	O 814	S 18	0	7	0

- Molecule 3 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	S	1	Total	Fe S	0	0
			8	4 4		
3	S	1	Total	Fe S	0	0
			8	4 4		
3	T	1	Total	Fe S	0	0
			8	4 4		
3	T	1	Total	Fe S	0	0
			8	4 4		

- Molecule 4 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe₃S₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	S	1	Total Fe S 7 3 4	0	0
4	T	1	Total Fe S 7 3 4	0	0

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	S	1	Total C O 6 3 3	0	0
5	S	1	Total C O 6 3 3	0	0
5	S	1	Total C O 6 3 3	0	0
5	L	1	Total C O 6 3 3	0	0
5	L	1	Total C O 6 3 3	0	0
5	L	1	Total C O 6 3 3	0	0
5	L	1	Total C O 6 3 3	0	0
5	T	1	Total C O 6 3 3	0	0
5	T	1	Total C O 6 3 3	0	0
5	M	1	Total C O 6 3 3	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	M	1	Total	C	O	0	0
			6	3	3		
5	M	1	Total	C	O	0	0
			6	3	3		
5	M	1	Total	C	O	0	0
			6	3	3		
5	M	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	S	1	Total	Mg	0	0
			1	1		
6	L	4	Total	Mg	0	0
			4	4		
6	M	3	Total	Mg	0	0
			3	3		

- Molecule 7 is CARBONMONOXIDE-(DICYANO) IRON (CCD ID: FCO) (formula: C₃FeN₂O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	L	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	M	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		

- Molecule 8 is NICKEL (II) ION (CCD ID: NI) (formula: Ni) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	L	1	Total	Ni	0	0
			1	1		
8	M	1	Total	Ni	0	0
			1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	S	220	Total	O	0	0
			220	220		
9	L	487	Total	O	0	0
			487	487		
9	T	201	Total	O	0	0
			201	201		
9	M	436	Total	O	0	0
			436	436		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hydrogenase-2 small chain




- Molecule 1: Hydrogenase-2 small chain



- Molecule 2: Hydrogenase-2 large chain



- Molecule 2: Hydrogenase-2 large chain



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	99.88Å 99.97Å 168.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	86.07 – 1.31 85.92 – 1.31	Depositor EDS
% Data completeness (in resolution range)	97.5 (86.07-1.31) 97.4 (85.92-1.31)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.98 (at 1.31Å)	Xtrriage
Refinement program	REFMAC 5.8.0405	Depositor
R, R_{free}	0.174 , 0.193 0.181 , 0.198	Depositor DCC
R_{free} test set	1979 reflections (0.49%)	wwPDB-VP
Wilson B-factor (Å ²)	14.8	Xtrriage
Anisotropy	0.133	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.014 for k,h,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	14333	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SF4, MG, NI, F3S, FCO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	S	0.63	0/2124	0.93	4/2892 (0.1%)
1	T	0.63	0/2154	0.93	4/2932 (0.1%)
2	L	0.62	0/4462	0.91	5/6081 (0.1%)
2	M	0.62	0/4445	0.92	3/6057 (0.0%)
All	All	0.63	0/13185	0.92	16/17962 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	S	0	1
1	T	0	1
2	L	0	1
All	All	0	3

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	S	66	HIS	CA-CB-CG	-8.60	105.20	113.80
2	M	196	GLU	CB-CG-CD	7.71	125.70	112.60
2	M	382	ASN	CA-CB-CG	6.90	119.50	112.60
2	L	99	HIS	CA-CB-CG	-6.81	106.99	113.80
2	L	382	ASN	CA-CB-CG	6.78	119.38	112.60
2	L	196	GLU	CB-CG-CD	6.74	124.06	112.60
1	S	212	ASP	CA-CB-CG	6.57	119.17	112.60
1	T	212	ASP	CA-CB-CG	6.25	118.85	112.60
2	M	99	HIS	CA-CB-CG	-6.14	107.66	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	T	81	ASP	CA-CB-CG	5.79	118.39	112.60
1	S	89	ASN	CB-CA-C	-5.70	102.51	111.51
1	S	81	ASP	CA-CB-CG	5.67	118.28	112.60
1	T	34	HIS	CA-CB-CG	5.65	119.45	113.80
2	L	126	ASP	CA-CB-CG	5.44	118.04	112.60
1	T	66	HIS	CA-CB-CG	-5.22	108.58	113.80
2	L	107	HIS	CA-CB-CG	-5.03	108.77	113.80

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	L	59	ARG	Sidechain
1	S	198	ARG	Sidechain
1	T	198	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	2063	0	1984	7	0
1	T	2088	0	2012	3	0
2	L	4346	0	4274	16	0
2	M	4332	0	4262	13	0
3	S	16	0	0	0	0
3	T	16	0	0	0	0
4	S	7	0	0	0	0
4	T	7	0	0	0	0
5	L	24	0	32	6	0
5	M	36	0	47	3	0
5	S	18	0	24	2	0
5	T	12	0	16	0	0
6	L	4	0	0	0	0
6	M	3	0	0	0	0
6	S	1	0	0	0	0
7	L	7	0	0	0	0
7	M	7	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	L	1	0	0	0	0
8	M	1	0	0	0	0
9	L	487	0	0	7	0
9	M	436	0	0	4	0
9	S	220	0	0	3	0
9	T	201	0	0	0	0
All	All	14333	0	12651	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:L:607:GOL:H12	9:L:855:HOH:O	1.87	0.74
2:M:481:MET:HE3	9:M:724:HOH:O	1.87	0.74
2:M:455:HIS:HD2	9:M:1068:HOH:O	1.82	0.63
2:M:313:ARG:HD3	2:M:322:TYR:CE2	2.36	0.61
1:S:140:GLN:HG2	9:S:611:HOH:O	2.03	0.57
2:L:481:MET:HE3	9:L:716:HOH:O	2.05	0.57
2:L:455:HIS:HD2	9:L:1098:HOH:O	1.88	0.56
2:L:6[A]:THR:HG21	9:L:820:HOH:O	2.06	0.55
5:L:606:GOL:H2	9:L:976:HOH:O	2.07	0.55
2:L:268:PHE:CZ	5:L:606:GOL:H31	2.42	0.54
5:M:607:GOL:H11	9:M:825:HOH:O	2.08	0.54
2:L:81:ASN:HA	5:L:607:GOL:H2	1.90	0.54
5:L:607:GOL:C1	9:L:855:HOH:O	2.52	0.52
1:S:202:ASP:OD2	1:T:202:ASP:OD2	2.28	0.50
2:M:156:LYS:NZ	9:M:703:HOH:O	2.26	0.50
1:S:62:GLU:HG2	9:S:584:HOH:O	2.12	0.50
2:L:313:ARG:HD3	2:L:322:TYR:CE2	2.48	0.48
2:L:374[B]:THR:HG21	2:L:516:GLY:HA2	1.95	0.48
2:M:115:ASP:HB3	2:M:538:ARG:HG2	1.96	0.48
2:L:81:ASN:C	5:L:607:GOL:H32	2.39	0.47
2:M:500:VAL:CG1	2:M:501:PRO:HD2	2.45	0.46
2:L:155:ILE:HD11	2:L:188:VAL:HG11	1.97	0.45
2:L:82:ILE:HD11	2:L:461:PRO:HB3	1.99	0.45
1:S:228:GLY:N	1:S:229:PRO:CD	2.81	0.44
5:S:306:GOL:H11	1:T:205:ARG:CZ	2.46	0.44
2:L:123:LEU:HD22	2:L:156:LYS:HG2	2.00	0.43
2:M:155:ILE:HD11	2:M:188:VAL:HG11	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:M:203:ARG:CZ	5:M:609:GOL:H32	2.49	0.43
1:T:228:GLY:N	1:T:229:PRO:CD	2.82	0.43
2:L:356:TRP:CD1	2:L:356:TRP:C	2.97	0.43
2:M:320:ASP:OD1	2:M:322:TYR:HB3	2.19	0.42
2:M:42:GLU:HB3	5:M:607:GOL:H32	2.01	0.42
2:M:159:VAL:HG22	2:M:164:LEU:HD11	2.01	0.42
2:L:159:VAL:HG22	2:L:164:LEU:HD11	2.02	0.42
2:M:356:TRP:CD1	2:M:356:TRP:C	2.98	0.42
2:L:115:ASP:HB3	2:L:538:ARG:HG2	2.02	0.42
2:L:500:VAL:CG1	2:L:501:PRO:HD2	2.50	0.41
1:S:271:GLN:OE1	5:S:305:GOL:H2	2.20	0.41
1:S:247:TRP:CZ2	1:S:249:VAL:HB	2.55	0.41
1:S:208:LYS:HE2	9:S:593:HOH:O	2.22	0.40
2:M:500:VAL:HG11	2:M:549:CYS:HB3	2.03	0.40
2:L:292[A]:SER:OG	9:L:701:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	270/298 (91%)	260 (96%)	10 (4%)	0	100	100
1	T	273/298 (92%)	264 (97%)	9 (3%)	0	100	100
2	L	557/567 (98%)	536 (96%)	20 (4%)	1 (0%)	43	17
2	M	554/567 (98%)	533 (96%)	20 (4%)	1 (0%)	43	17
All	All	1654/1730 (96%)	1593 (96%)	59 (4%)	2 (0%)	48	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	211	LYS
2	M	211	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	S	217/239 (91%)	215 (99%)	2 (1%)	70	42
1	T	220/239 (92%)	218 (99%)	2 (1%)	70	42
2	L	474/479 (99%)	471 (99%)	3 (1%)	78	54
2	M	471/479 (98%)	467 (99%)	4 (1%)	73	45
All	All	1382/1436 (96%)	1371 (99%)	11 (1%)	73	45

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	S	142	VAL
1	S	251	ILE
2	L	102	HIS
2	L	337	LYS
2	L	382	ASN
1	T	142	VAL
1	T	251	ILE
2	M	102	HIS
2	M	156	LYS
2	M	312	TYR
2	M	382	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	S	59	HIS
1	S	60	GLN
1	S	75	GLN
1	S	89	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	3	GLN
2	L	411	ASN
2	L	436	GLN
2	L	449	ASN
2	L	520	GLN
2	L	531	ASN
1	T	60	GLN
1	T	274	ASN
2	M	411	ASN
2	M	449	ASN
2	M	455	HIS
2	M	512	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 33 ligands modelled in this entry, 10 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	M	608	-	5,5,5	0.15	0	5,5,5	0.35	0
4	F3S	T	303	1	0,9,9	-	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SF4	S	301	1	0,12,12	-	-	-		
3	SF4	T	304	1	0,12,12	-	-	-		
5	GOL	L	607	-	5,5,5	0.26	0	5,5,5	0.91	0
7	FCO	L	602	2	0,6,6	-	-	-		
5	GOL	S	304	-	5,5,5	0.15	0	5,5,5	0.56	0
5	GOL	L	605	-	5,5,5	0.15	0	5,5,5	0.31	0
5	GOL	M	606	-	5,5,5	0.18	0	5,5,5	0.32	0
4	F3S	S	302	1	0,9,9	-	-	-		
5	GOL	M	609	6	5,5,5	0.20	0	5,5,5	0.34	0
5	GOL	L	601	-	5,5,5	0.25	0	5,5,5	0.60	0
5	GOL	T	301	-	5,5,5	0.19	0	5,5,5	0.51	0
5	GOL	M	607	-	5,5,5	0.05	0	5,5,5	0.26	0
5	GOL	T	305	-	5,5,5	0.21	0	5,5,5	0.47	0
5	GOL	S	306	-	5,5,5	0.26	0	5,5,5	0.75	0
5	GOL	L	606	-	5,5,5	0.11	0	5,5,5	0.32	0
5	GOL	M	605	-	5,5,5	0.14	0	5,5,5	0.22	0
5	GOL	S	305	-	5,5,5	0.21	0	5,5,5	0.27	0
5	GOL	M	601	-	5,5,5	0.39	0	5,5,5	1.08	0
3	SF4	S	303	1	0,12,12	-	-	-		
7	FCO	M	602	2	0,6,6	-	-	-		
3	SF4	T	302	1	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	M	608	-	-	0/4/4/4	-
4	F3S	T	303	1	-	-	0/3/3/3
3	SF4	S	301	1	-	-	0/6/5/5
3	SF4	T	304	1	-	-	0/6/5/5
5	GOL	L	607	-	-	4/4/4/4	-
5	GOL	S	304	-	-	0/4/4/4	-
5	GOL	L	605	-	-	0/4/4/4	-
5	GOL	M	606	-	-	0/4/4/4	-
4	F3S	S	302	1	-	-	0/3/3/3
5	GOL	M	609	6	-	0/4/4/4	-
5	GOL	L	601	-	-	0/4/4/4	-
5	GOL	T	301	-	-	4/4/4/4	-
5	GOL	M	607	-	-	4/4/4/4	-
5	GOL	T	305	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	S	306	-	-	4/4/4/4	-
5	GOL	L	606	-	-	3/4/4/4	-
5	GOL	M	605	-	-	0/4/4/4	-
5	GOL	S	305	-	-	1/4/4/4	-
5	GOL	M	601	-	-	3/4/4/4	-
3	SF4	S	303	1	-	-	0/6/5/5
3	SF4	T	302	1	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (23) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	S	306	GOL	O1-C1-C2-C3
5	S	306	GOL	C1-C2-C3-O3
5	L	606	GOL	O1-C1-C2-C3
5	L	607	GOL	C1-C2-C3-O3
5	T	301	GOL	O1-C1-C2-C3
5	T	301	GOL	C1-C2-C3-O3
5	M	601	GOL	C1-C2-C3-O3
5	M	607	GOL	O1-C1-C2-C3
5	M	607	GOL	C1-C2-C3-O3
5	L	607	GOL	O2-C2-C3-O3
5	S	306	GOL	O1-C1-C2-O2
5	S	306	GOL	O2-C2-C3-O3
5	L	606	GOL	O1-C1-C2-O2
5	T	301	GOL	O1-C1-C2-O2
5	T	301	GOL	O2-C2-C3-O3
5	M	601	GOL	O2-C2-C3-O3
5	M	607	GOL	O1-C1-C2-O2
5	M	607	GOL	O2-C2-C3-O3
5	L	607	GOL	O1-C1-C2-O2
5	S	305	GOL	O1-C1-C2-O2
5	L	606	GOL	O2-C2-C3-O3
5	L	607	GOL	O1-C1-C2-C3
5	M	601	GOL	O1-C1-C2-O2


There are no ring outliers.

6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	L	607	GOL	4	0
5	M	609	GOL	1	0
5	M	607	GOL	2	0
5	S	306	GOL	1	0
5	L	606	GOL	2	0
5	S	305	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	S	268/298 (89%)	0.10	5 (1%) 66 70	6, 15, 30, 53	4 (1%)
1	T	267/298 (89%)	0.29	6 (2%) 62 67	7, 17, 32, 50	8 (2%)
2	L	551/567 (97%)	-0.04	6 (1%) 78 81	5, 15, 27, 38	9 (1%)
2	M	550/567 (97%)	0.24	21 (3%) 44 49	6, 17, 31, 51	7 (1%)
All	All	1636/1730 (94%)	0.13	38 (2%) 61 66	5, 16, 30, 53	28 (1%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	M	27	VAL	4.5
2	M	137	VAL	4.1
2	M	515	VAL	3.8
1	S	9	PRO	3.8
2	L	137	VAL	3.5
1	T	66	HIS	3.3
1	S	276	GLU	3.1
2	M	23	ILE	3.0
2	M	413	LEU	3.0
1	T	145	GLY	2.9
1	T	276	GLU	2.9
2	M	160	ALA	2.8
2	L	515	VAL	2.7
2	M	528	ALA	2.6
1	S	275	VAL	2.6
1	S	66	HIS	2.4
1	T	176[A]	LYS	2.4
1	T	275	VAL	2.4
2	L	166	ILE	2.3
2	M	32	TRP	2.3
2	M	167	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	T	142	VAL	2.3
2	L	136	GLY	2.2
2	M	119	ILE	2.2
2	M	407	LYS	2.2
2	M	516	GLY	2.2
2	M	5	ILE	2.2
1	S	145	GLY	2.2
2	M	25	ASN	2.2
2	M	415	VAL	2.1
2	M	166	ILE	2.1
2	M	159	VAL	2.1
2	M	28	VAL	2.1
2	L	160	ALA	2.0
2	M	414	GLU	2.0
2	M	172	TRP	2.0
2	L	415	VAL	2.0
2	M	410	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	L	607	6/6	0.77	0.18	33,39,43,44	0
5	GOL	M	605	6/6	0.83	0.14	43,44,46,49	0
5	GOL	M	609	6/6	0.83	0.15	27,33,35,42	0
6	MG	M	611	1/1	0.83	0.14	41,41,41,41	0
5	GOL	L	605	6/6	0.85	0.14	31,37,38,43	0
5	GOL	M	607	6/6	0.86	0.15	32,44,49,59	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	S	306	6/6	0.87	0.15	30,37,39,41	0
6	MG	L	609	1/1	0.87	0.13	51,51,51,51	0
5	GOL	L	601	6/6	0.87	0.14	22,34,40,48	0
5	GOL	T	301	6/6	0.88	0.15	30,40,41,46	0
5	GOL	S	305	6/6	0.88	0.12	35,36,37,38	0
5	GOL	M	608	6/6	0.90	0.12	29,36,40,41	0
5	GOL	M	601	6/6	0.90	0.13	19,27,35,38	0
5	GOL	L	606	6/6	0.91	0.11	40,45,51,53	0
6	MG	L	610	1/1	0.94	0.10	37,37,37,37	0
5	GOL	M	606	6/6	0.94	0.10	18,34,36,42	0
6	MG	L	608	1/1	0.96	0.09	24,24,24,24	0
5	GOL	S	304	6/6	0.97	0.06	19,19,20,20	0
5	GOL	T	305	6/6	0.97	0.06	17,21,22,23	0
6	MG	M	610	1/1	0.98	0.10	21,21,21,21	0
8	NI	M	603	1/1	0.99	0.03	15,15,15,15	0
6	MG	S	307	1/1	1.00	0.02	12,12,12,12	0
6	MG	L	604	1/1	1.00	0.04	7,7,7,7	0
3	SF4	T	304	8/8	1.00	0.02	11,13,13,14	0
4	F3S	S	302	7/7	1.00	0.02	10,11,11,11	0
4	F3S	T	303	7/7	1.00	0.02	11,12,12,12	0
6	MG	M	604	1/1	1.00	0.04	9,9,9,9	0
3	SF4	S	301	8/8	1.00	0.02	12,12,13,14	0
3	SF4	S	303	8/8	1.00	0.02	10,11,11,12	0
7	FCO	L	602	7/7	1.00	0.02	10,10,11,12	0
7	FCO	M	602	7/7	1.00	0.03	10,11,13,13	0
8	NI	L	603	1/1	1.00	0.01	13,13,13,13	0
3	SF4	T	302	8/8	1.00	0.02	13,13,13,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NI M 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FCO L 602:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FCO M 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.