



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 08:02 PM UTC

PDB ID : 1EUZ / pdb_00001euz
Title : GLUTAMATE DEHYDROGENASE FROM THERMOCOCCUS PROFUNDUS IN THE UNLIGATED STATE
Authors : Nakasako, M.
Deposited on : 2000-04-19
Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	419	75% 22% ..
1	B	419	75% 22% ..
1	C	419	81% 16% ..
1	D	419	74% 24% ..
1	E	419	78% 18% ..
1	F	419	71% 24% ..

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 19422 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTAMATE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	416	Total	C	N	O	S	0	0	0
			3226	2062	546	604	14			
1	B	416	Total	C	N	O	S	0	0	0
			3205	2050	543	598	14			
1	C	416	Total	C	N	O	S	0	0	0
			3225	2063	550	598	14			
1	D	416	Total	C	N	O	S	0	0	0
			3215	2054	551	596	14			
1	E	416	Total	C	N	O	S	0	0	0
			3258	2080	556	608	14			
1	F	417	Total	C	N	O	S	0	0	0
			3208	2047	542	605	14			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: GLUTAMATE DEHYDROGENASE




• Molecule 1: GLUTAMATE DEHYDROGENASE




• Molecule 1: GLUTAMATE DEHYDROGENASE






• Molecule 1: GLUTAMATE DEHYDROGENASE

Chain D: 74% 24% ..




• Molecule 1: GLUTAMATE DEHYDROGENASE

Chain E: 78% 18% ..



• Molecule 1: GLUTAMATE DEHYDROGENASE

Chain F: 71% 24% ..



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	112.48Å 163.12Å 132.19Å 90.00° 113.20° 90.00°	Depositor
Resolution (Å)	8.00 – 2.25	Depositor
% Data completeness (in resolution range)	88.8 (8.00-2.25)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.187 , 0.250	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	19422	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.67	0/3299	0.89	4/4479 (0.1%)
1	B	0.63	1/3278 (0.0%)	0.88	3/4453 (0.1%)
1	C	0.70	2/3298 (0.1%)	0.89	3/4476 (0.1%)
1	D	0.64	2/3288 (0.1%)	0.89	4/4465 (0.1%)
1	E	0.67	0/3331	0.89	1/4514 (0.0%)
1	F	0.67	1/3280 (0.0%)	0.89	7/4459 (0.2%)
All	All	0.67	6/19774 (0.0%)	0.89	22/26846 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	3
1	D	0	3
1	E	0	4
1	F	0	6
All	All	0	19

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	35	MET	SD-CE	-11.74	1.50	1.79
1	C	9	MET	SD-CE	8.36	2.00	1.79
1	D	35	MET	SD-CE	-8.27	1.58	1.79
1	C	35	MET	SD-CE	-7.65	1.60	1.79
1	D	9	MET	SD-CE	7.39	1.98	1.79

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	406	ARG	NE-CZ-NH2	-7.48	112.47	119.20
1	F	273	PHE	CA-C-N	6.20	127.58	119.84
1	F	273	PHE	C-N-CA	6.20	127.58	119.84
1	A	406	ARG	CG-CD-NE	-6.16	98.44	112.00
1	E	40	VAL	N-CA-C	6.07	117.23	108.48

There are no chirality outliers.

5 of 19 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	101	TYR	Sidechain
1	A	341	PHE	Sidechain
1	A	363	TYR	Sidechain
1	B	160	TYR	Sidechain
1	B	363	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3226	0	3192	59	0
1	B	3205	0	3160	67	0
1	C	3225	0	3204	43	0
1	D	3215	0	3178	69	0
1	E	3258	0	3255	54	0
1	F	3208	0	3141	82	0
2	A	25	0	0	1	0
2	B	15	0	0	1	0
2	C	10	0	0	0	0
2	D	5	0	0	0	0
2	E	20	0	0	2	0
2	F	10	0	0	0	0
All	All	19422	0	19130	363	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 363 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:5:ASP:HA	1:F:8:GLU:HB2	1.33	1.09
1:C:35:MET:HE3	1:C:62:ASN:HA	1.48	0.95
1:D:35:MET:HE3	1:D:62:ASN:HA	1.50	0.94
1:F:4:ILE:HA	1:F:81:SER:OG	1.71	0.89
1:F:35:MET:HE3	1:F:62:ASN:HA	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	414/419 (99%)	395 (95%)	18 (4%)	1 (0%)	43	50
1	B	414/419 (99%)	395 (95%)	19 (5%)	0	100	100
1	C	414/419 (99%)	397 (96%)	17 (4%)	0	100	100
1	D	414/419 (99%)	394 (95%)	19 (5%)	1 (0%)	43	50
1	E	414/419 (99%)	395 (95%)	19 (5%)	0	100	100
1	F	415/419 (99%)	377 (91%)	30 (7%)	8 (2%)	6	3
All	All	2485/2514 (99%)	2353 (95%)	122 (5%)	10 (0%)	30	31

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	4	ILE
1	F	5	ASP
1	F	297	ILE
1	F	232	LYS
1	F	274	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/342 (96%)	310 (94%)	20 (6%)	17	17
1	B	325/342 (95%)	306 (94%)	19 (6%)	18	19
1	C	329/342 (96%)	308 (94%)	21 (6%)	16	15
1	D	326/342 (95%)	316 (97%)	10 (3%)	35	44
1	E	337/342 (98%)	319 (95%)	18 (5%)	20	22
1	F	325/342 (95%)	296 (91%)	29 (9%)	9	7
All	All	1972/2052 (96%)	1855 (94%)	117 (6%)	18	18

5 of 117 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	379	LYS
1	F	332	GLU
1	E	28	LEU
1	F	323	THR
1	F	249	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	75	HIS
1	F	120	GLN
1	F	344	ASN
1	C	75	HIS
1	C	62	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	601	-	4,4,4	0.39	0	6,6,6	1.13	1 (16%)
2	SO4	E	613	-	4,4,4	0.38	0	6,6,6	1.02	0
2	SO4	F	615	-	4,4,4	0.37	0	6,6,6	0.37	0
2	SO4	E	611	-	4,4,4	0.35	0	6,6,6	0.90	0
2	SO4	B	608	-	4,4,4	0.26	0	6,6,6	0.86	0
2	SO4	B	607	-	4,4,4	0.13	0	6,6,6	0.38	0
2	SO4	E	612	-	4,4,4	0.46	0	6,6,6	0.96	0
2	SO4	C	604	-	4,4,4	0.35	0	6,6,6	1.83	2 (33%)
2	SO4	A	602	-	4,4,4	0.27	0	6,6,6	0.62	0
2	SO4	A	603	-	4,4,4	0.26	0	6,6,6	0.54	0
2	SO4	C	609	-	4,4,4	0.38	0	6,6,6	1.16	0
2	SO4	A	605	-	4,4,4	0.11	0	6,6,6	0.18	0
2	SO4	E	614	-	4,4,4	0.41	0	6,6,6	0.87	0
2	SO4	D	610	-	4,4,4	0.34	0	6,6,6	0.78	0
2	SO4	B	606	-	4,4,4	0.14	0	6,6,6	0.74	0
2	SO4	A	600	-	4,4,4	0.55	0	6,6,6	1.00	0
2	SO4	F	616	-	4,4,4	0.36	0	6,6,6	0.57	0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	604	SO4	O4-S-O2	3.01	125.31	109.56
2	C	604	SO4	O4-S-O1	-2.34	97.32	109.56
2	A	601	SO4	O3-S-O2	2.27	121.42	109.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	613	SO4	1	0
2	E	612	SO4	1	0
2	A	605	SO4	1	0
2	B	606	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.