



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 02:24 PM UTC

PDB ID : 7EVU / pdb_00007evu
Title : Co-crystal Structure of Toxoplasma gondii Prolyl tRNA Synthetase (TgPRS)
in complex with HFG and JE6
Authors : Mishra, S.; Malhotra, N.; Manickam, Y.; Sharma, A.
Deposited on : 2021-05-22
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	500	13% 90% 7%
1	B	500	13% 89% 6%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8557 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Prolyl-tRNA synthetase (ProRS).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	484	Total	C	N	O	S	0	0	0
			3948	2535	676	714	23			
1	B	480	Total	C	N	O	S	0	1	0
			3938	2526	676	714	22			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	331	GLY	-	expression tag	UNP A0A7J6JUK2
A	332	ALA	-	expression tag	UNP A0A7J6JUK2
A	333	MET	-	expression tag	UNP A0A7J6JUK2
B	331	GLY	-	expression tag	UNP A0A7J6JUK2
B	332	ALA	-	expression tag	UNP A0A7J6JUK2
B	333	MET	-	expression tag	UNP A0A7J6JUK2

- Molecule 2 is 7-bromo-6-chloro-3-{3-[(2R,3S)-3-hydroxypiperidin-2-yl]-2-oxopropyl}quinazolin-4(3H)-one (CCD ID: HFG) (formula: C₁₆H₁₇BrClN₃O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	Br	C	Cl	N	O	0	0
			24	1	16	1	3	3		
2	B	1	Total	Br	C	Cl	N	O	0	0
			24	1	16	1	3	3		

- Molecule 3 is {N}-[4-[(3 {S})-3-cyano-3-cyclopropyl-2-oxidanylidene-pyrrolidin-1-yl]-6-methyl-pyridin-2-yl]-2-phenyl-ethanamide (CCD ID: JE6) (formula: C₂₂H₂₂N₄O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			28	22	4	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			28	22	4	2		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	277	Total	O	0	0
			277	277		
5	B	278	Total	O	0	0
			278	278		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Prolyl-tRNA synthetase (ProRS)



• Molecule 1: Prolyl-tRNA synthetase (ProRS)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	70.75Å 70.83Å 76.04Å 92.16° 101.44° 115.61°	Depositor
Resolution (Å)	54.24 – 1.70 54.24 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.8 (54.24-1.70) 96.9 (54.24-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.15rc2_3428	Depositor
R, R_{free}	0.195 , 0.227 0.203 , 0.231	Depositor DCC
R_{free} test set	6806 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.264	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8557	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.32 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.9190e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: JE6, MES, HFG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	2/4053 (0.0%)	0.74	2/5481 (0.0%)
1	B	0.53	0/4043	0.76	6/5465 (0.1%)
All	All	0.54	2/8096 (0.0%)	0.75	8/10946 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	590	GLY	N-CA	5.67	1.51	1.45
1	A	591	CYS	CA-C	-5.09	1.46	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	659	SER	N-CA-C	7.54	122.30	113.18
1	A	437	ARG	N-CA-C	6.41	117.46	109.57
1	B	659	SER	CA-C-N	5.43	132.21	122.38
1	B	659	SER	C-N-CA	5.43	132.21	122.38
1	B	487	TRP	CA-C-N	-5.41	112.82	122.07
1	B	487	TRP	C-N-CA	-5.41	112.82	122.07
1	B	658	ARG	O-C-N	-5.38	116.28	122.95
1	A	828	ARG	N-CA-C	-5.32	103.00	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3948	0	3915	22	0
1	B	3938	0	3904	24	0
2	A	24	0	17	1	0
2	B	24	0	17	0	0
3	A	28	0	0	0	0
3	B	28	0	0	0	0
4	B	12	0	13	2	0
5	A	277	0	0	2	0
5	B	278	0	0	4	0
All	All	8557	0	7866	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:487:TRP:CD1	1:B:487:TRP:O	1.87	1.28
1:B:487:TRP:O	1:B:487:TRP:HD1	1.18	1.11
1:B:487:TRP:CD1	1:B:487:TRP:C	2.53	0.86
1:A:632:THR:HG23	1:A:634:GLU:H	1.51	0.74
1:A:367:MET:HE1	1:A:593:THR:HG21	1.73	0.70
1:A:632:THR:HG21	1:A:634:GLU:OE1	1.95	0.67
1:B:367:MET:HE1	1:B:593:THR:HG21	1.80	0.63
1:B:627:PHE:O	1:B:630:GLU:HG3	2.01	0.60
1:A:628:LYS:HG2	1:A:659:SER:O	2.04	0.58
1:A:578:GLU:O	1:A:578:GLU:HG3	2.04	0.57
1:A:628:LYS:HE3	1:A:660:ASN:HA	1.86	0.56
1:A:750:ASN:CG	1:A:780:GLN:HE22	2.13	0.55
1:B:515:TYR:CZ	1:B:556:ALA:HB1	2.42	0.55
1:A:515:TYR:CZ	1:A:556:ALA:HB1	2.43	0.53
1:B:402:ARG:O	1:B:406:GLU:HG3	2.10	0.52
1:B:487:TRP:HB2	1:B:592:THR:HG22	1.91	0.51
1:A:623:ILE:HD12	1:A:678:ARG:HD2	1.93	0.51
1:B:404:LYS:NZ	5:B:1017:HOH:O	2.44	0.50
1:B:333:MET:O	1:B:352:ARG:NH1	2.34	0.50
1:B:466:CYS:HA	4:B:903:MES:H32	1.94	0.50
1:B:625:ILE:HD11	1:B:680:GLU:HB3	1.94	0.49
1:B:776:LEU:O	1:B:780:GLN:HG3	2.11	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:903:MES:H82	5:B:1212:HOH:O	2.12	0.48
1:B:684:LYS:O	1:B:688:LYS:HG3	2.13	0.48
1:A:638:LYS:HD3	1:A:641:GLU:OE1	2.14	0.48
1:A:453:ARG:HD3	1:A:453:ARG:HA	1.62	0.46
1:A:677:LEU:HD13	1:A:713:LEU:HD22	1.96	0.46
1:A:455:HIS:HE1	5:A:1240:HOH:O	1.99	0.46
1:B:638:LYS:HD3	1:B:638:LYS:HA	1.48	0.45
1:A:685:ASP:HB3	1:A:690:THR:O	2.17	0.44
1:A:805:PRO:HG2	1:A:823:TRP:CD1	2.52	0.44
1:B:620:VAL:HG22	1:B:677:LEU:HB2	1.99	0.44
1:A:373:HIS:CE1	5:A:1082:HOH:O	2.70	0.44
1:B:685:ASP:HB3	1:B:690:THR:O	2.18	0.44
1:B:487:TRP:CB	1:B:592:THR:HG22	2.49	0.43
1:B:453:ARG:HG2	1:B:457:ASP:OD2	2.18	0.43
1:B:684:LYS:HD3	1:B:688:LYS:HE2	2.00	0.43
1:A:638:LYS:HD3	1:A:638:LYS:HA	1.85	0.43
1:A:619:GLN:HB3	1:A:717:MET:SD	2.59	0.42
1:A:419:VAL:HG13	2:A:901:HFG:BR1	2.75	0.41
1:B:559:SER:HA	1:B:589:TRP:HB3	2.01	0.41
1:B:609:LEU:HD13	1:B:611:ILE:HD11	2.03	0.41
1:A:373:HIS:HB3	1:A:617:SER:HB3	2.02	0.41
1:B:539:LYS:NZ	5:B:1012:HOH:O	2.40	0.41
1:A:636:LEU:HD23	1:A:636:LEU:HA	1.87	0.41
1:B:685:ASP:OD2	5:B:1001:HOH:O	2.22	0.40
1:A:374:ILE:HD13	1:A:599:MET:HE1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	478/500 (96%)	471 (98%)	7 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	475/500 (95%)	468 (98%)	7 (2%)	0	100	100
All	All	953/1000 (95%)	939 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	422/436 (97%)	422 (100%)	0	100	100
1	B	422/436 (97%)	417 (99%)	5 (1%)	63	51
All	All	844/872 (97%)	839 (99%)	5 (1%)	81	72

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	453	ARG
1	B	487	TRP
1	B	638	LYS
1	B	654[A]	ARG
1	B	654[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	780	GLN
1	B	555	GLN
1	B	669	HIS
1	B	780	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HFG	A	901	-	26,26,26	0.55	1 (3%)	29,37,37	0.79	1 (3%)
3	JE6	A	902	-	29,31,31	0.35	0	42,45,45	0.69	1 (2%)
4	MES	B	903	-	12,12,12	2.33	1 (8%)	15,16,16	1.17	1 (6%)
3	JE6	B	902	-	29,31,31	0.38	0	42,45,45	0.74	1 (2%)
2	HFG	B	901	-	26,26,26	0.67	1 (3%)	29,37,37	0.86	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HFG	A	901	-	-	0/8/19/19	0/3/3/3
3	JE6	A	902	-	-	0/16/39/39	0/4/4/4
4	MES	B	903	-	-	5/6/14/14	0/1/1/1
3	JE6	B	902	-	-	0/16/39/39	0/4/4/4
2	HFG	B	901	-	-	0/8/19/19	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	903	MES	C8-S	-7.92	1.66	1.77
2	B	901	HFG	C1'-C21	3.07	1.55	1.51
2	A	901	HFG	C1'-C21	2.51	1.54	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	901	HFG	C6'-N1'-C2'	-3.32	109.55	111.62
3	A	902	JE6	C15-C16-C17	2.78	115.77	111.33
3	B	902	JE6	C15-C16-C17	2.68	115.61	111.33
2	A	901	HFG	C5'-C4'-C39	-2.65	106.64	111.56
4	B	903	MES	O2S-S-C8	2.41	110.37	106.73

There are no chirality outliers.

All (5) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	B	903	MES	C7-C8-S-O1S
4	B	903	MES	C7-C8-S-O3S
4	B	903	MES	C8-C7-N4-C3
4	B	903	MES	C7-C8-S-O2S
4	B	903	MES	C8-C7-N4-C5


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	HFG	1	0
4	B	903	MES	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	484/500 (96%)	0.84	65 (13%) 7 6	20, 35, 61, 87	0
1	B	480/500 (96%)	0.90	65 (13%) 7 6	21, 36, 66, 86	1 (0%)
All	All	964/1000 (96%)	0.87	130 (13%) 7 6	20, 35, 64, 87	1 (0%)

All (130) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	473	PHE	8.1
1	B	473	PHE	6.8
1	B	781	ALA	6.2
1	A	332	ALA	5.3
1	A	792	THR	4.2
1	B	333	MET	4.1
1	B	792	THR	4.1
1	A	707	ALA	4.0
1	B	646	LEU	3.9
1	A	626	LEU	3.8
1	A	628	LYS	3.8
1	A	333	MET	3.8
1	A	646	LEU	3.7
1	B	537	GLY	3.7
1	A	417	PRO	3.7
1	B	417	PRO	3.6
1	A	651	ILE	3.5
1	A	819	PRO	3.4
1	B	627	PHE	3.4
1	B	708	ASP	3.4
1	B	626	LEU	3.4
1	B	334	VAL	3.4
1	B	709	LEU	3.3
1	A	649	ALA	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	707	ALA	3.2
1	B	740	SER	3.2
1	A	629	ASP	3.2
1	A	630	GLU	3.2
1	B	741	THR	3.1
1	B	487	TRP	3.1
1	A	431	PRO	3.0
1	A	776	LEU	3.0
1	B	681	LEU	3.0
1	A	416	SER	3.0
1	B	743	ASP	3.0
1	B	632	THR	3.0
1	B	649	ALA	3.0
1	A	627	PHE	2.9
1	A	733	HIS	2.9
1	A	782	ILE	2.9
1	B	654[A]	ARG	2.8
1	A	334	VAL	2.8
1	A	781	ALA	2.8
1	B	418	GLU	2.8
1	A	747	PRO	2.8
1	A	373	HIS	2.8
1	A	631	ASN	2.8
1	B	416	SER	2.8
1	A	711	PRO	2.7
1	B	704	ILE	2.7
1	B	668	ASN	2.7
1	A	578	GLU	2.6
1	A	808	PRO	2.6
1	B	536	GLY	2.6
1	B	733	HIS	2.6
1	A	740	SER	2.6
1	B	335	THR	2.6
1	B	714	LEU	2.6
1	A	565	ASN	2.6
1	B	660	ASN	2.6
1	A	810	GLY	2.6
1	A	425	TYR	2.6
1	A	335	THR	2.5
1	A	360	ASP	2.5
1	A	537	GLY	2.5
1	A	741	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	711	PRO	2.5
1	B	808	PRO	2.5
1	B	686	LEU	2.5
1	A	645	MET	2.5
1	B	568	LYS	2.4
1	A	577	GLU	2.4
1	A	687	ALA	2.4
1	A	734	GLU	2.4
1	B	811	THR	2.4
1	A	704	ILE	2.4
1	A	791	MET	2.4
1	A	405	LEU	2.4
1	A	818	LYS	2.4
1	A	632	THR	2.3
1	B	629	ASP	2.3
1	B	501	TRP	2.3
1	A	472	GLU	2.3
1	B	715	GLU	2.3
1	A	803	ASP	2.3
1	A	806	PRO	2.3
1	B	713	LEU	2.3
1	B	645	MET	2.3
1	A	811	THR	2.3
1	B	405	LEU	2.3
1	A	633	GLY	2.3
1	B	806	PRO	2.2
1	A	648	LYS	2.2
1	A	637	GLY	2.2
1	B	419	VAL	2.2
1	B	747	PRO	2.2
1	B	644	THR	2.2
1	A	471	TRP	2.2
1	A	501	TRP	2.2
1	B	538	LYS	2.2
1	B	636	LEU	2.2
1	B	814	PHE	2.2
1	B	687	ALA	2.2
1	B	471	TRP	2.2
1	A	418	GLU	2.2
1	A	742	PHE	2.2
1	A	661	TYR	2.2
1	B	651	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	713	LEU	2.1
1	A	714	LEU	2.1
1	A	659	SER	2.1
1	A	743	ASP	2.1
1	B	429	PRO	2.1
1	B	431	PRO	2.1
1	B	819	PRO	2.1
1	A	830	TYR	2.1
1	B	625	ILE	2.1
1	B	738	LYS	2.1
1	B	631	ASN	2.1
1	B	803	ASP	2.1
1	B	640	ARG	2.1
1	B	736	ILE	2.1
1	B	344	GLU	2.1
1	A	650	ASP	2.1
1	A	536	GLY	2.0
1	A	820	ALA	2.0
1	B	755	VAL	2.0
1	B	565	ASN	2.0
1	B	361	ILE	2.0
1	B	779	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	HFG	B	901	24/24	0.81	0.14	19,25,33,71	24

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MES	B	903	12/12	0.82	0.16	25,30,46,47	12
2	HFG	A	901	24/24	0.87	0.12	17,23,33,78	24
3	JE6	B	902	28/28	0.92	0.11	19,27,37,42	0
3	JE6	A	902	28/28	0.93	0.10	21,28,38,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around HFG A 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around JE6 B 902:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.