



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 24, 2026 – 01:13 PM UTC

PDB ID : 8EVV / pdb_00008evv
Title : Apo form of DdlA from Pseudomonas aeruginosa PAO1
Authors : Pederick, J.L.; Woolman, J.C.; Bruning, J.B.
Deposited on : 2022-10-21
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	
1	B	347	
1	C	347	
1	D	347	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-alanine–D-alanine ligase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	338	Total 2467	C 1551	N 435	O 472	S 9	0	0	0
1	B	307	Total 2219	C 1399	N 391	O 420	S 9	0	0	0
1	C	314	Total 2294	C 1442	N 405	O 438	S 9	0	1	0
1	D	312	Total 2274	C 1433	N 402	O 430	S 9	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q9HWI0
B	0	GLY	-	expression tag	UNP Q9HWI0
C	0	GLY	-	expression tag	UNP Q9HWI0
D	0	GLY	-	expression tag	UNP Q9HWI0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	342	Total 342	O 342	0	0
2	B	151	Total 151	O 151	0	0
2	C	209	Total 209	O 209	0	0
2	D	214	Total 214	O 214	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: D-alanine–D-alanine ligase A




- Molecule 1: D-alanine–D-alanine ligase A




- Molecule 1: D-alanine–D-alanine ligase A



- Molecule 1: D-alanine–D-alanine ligase A





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.37Å 80.33Å 271.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.73 – 2.05 39.73 – 2.05	Depositor EDS
% Data completeness (in resolution range)	98.1 (39.73-2.05) 98.1 (39.73-2.05)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.05Å)	Xtrriage
Refinement program	PHENIX 1.20.1-4487	Depositor
R, R_{free}	0.211 , 0.252 0.211 , 0.251	Depositor DCC
R_{free} test set	5048 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtrriage
Anisotropy	0.238	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 30.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10170	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/2503	0.33	0/3391
1	B	0.10	0/2246	0.26	0/3040
1	C	0.11	0/2325	0.29	0/3145
1	D	0.11	0/2305	0.28	0/3117
All	All	0.11	0/9379	0.29	0/12693

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2467	0	2449	23	0
1	B	2219	0	2219	13	0
1	C	2294	0	2273	20	0
1	D	2274	0	2265	9	0
2	A	342	0	0	2	0
2	B	151	0	0	0	0
2	C	209	0	0	1	0
2	D	214	0	0	0	0
All	All	10170	0	9206	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:THR:HA	1:A:75:ILE:HD13	1.71	0.72
1:A:325:GLU:OE2	1:A:329:ARG:NH1	2.28	0.68
1:A:180:VAL:HG11	1:A:198:ALA:HA	1.78	0.65
1:D:127:ALA:HB1	1:D:279:LEU:HD23	1.79	0.64
1:A:71:ASP:OD2	1:A:77:ARG:NE	2.26	0.64
1:C:127:ALA:HB1	1:C:279:LEU:HD23	1.83	0.61
1:B:127:ALA:HB1	1:B:279:LEU:HD23	1.83	0.60
1:C:5:LYS:HE3	1:C:37:GLY:HA3	1.84	0.59
1:C:159:GLU:OE1	1:C:162:ARG:NH1	2.36	0.58
1:D:53:GLU:HG3	1:D:57:ARG:HH12	1.67	0.57
1:C:147:GLU:OE2	1:C:294[B]:ARG:NH2	2.38	0.56
1:C:178:SER:O	1:C:180:VAL:N	2.38	0.53
1:A:127:ALA:HB1	1:A:279:LEU:HD23	1.92	0.51
1:C:14:GLY:H	1:C:45:THR:HG1	1.57	0.51
1:A:185:VAL:HG13	1:A:190:GLU:HB2	1.91	0.51
1:B:171:VAL:HG22	1:B:207:LEU:HD22	1.93	0.50
1:B:72:SER:OG	1:B:73:LEU:N	2.43	0.50
1:C:337:ARG:NH1	2:C:411:HOH:O	2.44	0.50
1:A:76:ILE:HD12	1:A:77:ARG:N	2.27	0.49
1:B:292:GLN:OE1	1:B:294:ARG:NH2	2.41	0.49
1:B:76:ILE:HG22	1:B:109:LEU:HD13	1.95	0.49
1:A:75:ILE:HB	1:A:84:LEU:CD1	2.42	0.49
1:A:75:ILE:HB	1:A:84:LEU:HD11	1.93	0.49
1:B:19:ARG:NH1	1:B:44:ASP:OD2	2.36	0.49
1:A:138:LYS:NZ	1:A:148:THR:OG1	2.37	0.48
1:C:53:GLU:OE2	1:C:56:ARG:NH1	2.45	0.48
1:D:158:GLU:HB2	1:D:188:GLN:NE2	2.28	0.47
1:D:171:VAL:HG22	1:D:207:LEU:HD22	1.97	0.47
1:C:233:LEU:O	1:C:235:GLY:N	2.47	0.47
1:A:240:ASP:HB3	1:A:243:HIS:HB2	1.95	0.47
1:C:233:LEU:HD23	1:C:240:ASP:O	2.15	0.47
1:B:107:GLY:O	1:B:125:HIS:HB2	2.15	0.46
1:B:259:PRO:HG2	1:B:262:ILE:HG13	1.96	0.46
1:B:138:LYS:HE3	1:B:151:TRP:CD2	2.51	0.46
1:D:109:LEU:O	1:D:113:LEU:HG	2.16	0.46
1:A:337:ARG:NH2	2:A:417:HOH:O	2.48	0.45
1:C:233:LEU:HD22	1:C:244:LYS:HA	1.98	0.45
1:B:44:ASP:OD1	1:B:45:THR:N	2.50	0.45
1:C:89:LEU:HD13	1:C:95:PHE:HZ	1.82	0.45
1:D:15:SER:O	1:D:19:ARG:HG3	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:LYS:NZ	1:A:90:ALA:O	2.41	0.45
1:C:34:ARG:HH22	1:C:55:GLU:CD	2.25	0.44
1:C:335:ILE:O	1:C:339:LYS:HG2	2.16	0.44
1:C:292:GLN:HB2	1:C:294[A]:ARG:HH12	1.83	0.44
1:A:53:GLU:OE2	1:A:57:ARG:NE	2.50	0.43
1:A:329:ARG:NH2	2:A:404:HOH:O	2.36	0.43
1:D:99:LEU:HD13	1:D:103:THR:HG22	2.01	0.43
1:C:23:ILE:HG21	1:C:63:VAL:HA	1.99	0.43
1:A:41:LEU:HD11	1:A:88:GLU:HG2	2.01	0.42
1:A:231:ILE:HD13	1:A:239:PHE:CE2	2.54	0.42
1:C:294[A]:ARG:HB2	1:C:294[A]:ARG:NH1	2.35	0.42
1:A:77:ARG:HH22	1:C:115:ALA:C	2.27	0.42
1:B:14:GLY:HA2	1:B:103:THR:HG21	2.01	0.42
1:A:171:VAL:HG22	1:A:207:LEU:HD22	2.02	0.42
1:A:243:HIS:HB3	1:A:245:TYR:CE1	2.54	0.41
1:D:165:LEU:O	1:D:209:ARG:HG3	2.19	0.41
1:C:15:SER:O	1:C:19:ARG:HG3	2.19	0.41
1:D:312:LEU:HB3	1:D:313:PRO:HD3	2.03	0.41
1:A:71:ASP:CG	1:A:77:ARG:HE	2.19	0.41
1:C:271:LEU:HB3	1:C:275:ARG:NH1	2.36	0.41
1:A:77:ARG:HH21	1:A:77:ARG:HD3	1.77	0.40
1:B:138:LYS:HA	1:B:141:PHE:CD2	2.56	0.40
1:B:230:GLU:HG2	1:B:232:LEU:HD22	2.03	0.40
1:A:135:ASP:O	1:A:139:ARG:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	334/347 (96%)	325 (97%)	8 (2%)	1 (0%)	36 30

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	297/347 (86%)	286 (96%)	10 (3%)	1 (0%)	36	30
1	C	309/347 (89%)	300 (97%)	8 (3%)	1 (0%)	36	30
1	D	304/347 (88%)	295 (97%)	9 (3%)	0	100	100
All	All	1244/1388 (90%)	1206 (97%)	35 (3%)	3 (0%)	43	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	234	GLY
1	B	201	TYR
1	A	245	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/261 (94%)	242 (99%)	2 (1%)	73	77
1	B	218/261 (84%)	217 (100%)	1 (0%)	81	85
1	C	225/261 (86%)	222 (99%)	3 (1%)	61	63
1	D	224/261 (86%)	224 (100%)	0	100	100
All	All	911/1044 (87%)	905 (99%)	6 (1%)	81	80

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	88	GLU
1	A	277	LEU
1	B	125	HIS
1	C	294[A]	ARG
1	C	294[B]	ARG
1	C	295	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	100	HIS
1	B	224	GLN
1	C	292	GLN
1	C	314	GLN
1	D	188	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	338/347 (97%)	0.12	19 (5%) 30 30	21, 31, 62, 88	0
1	B	307/347 (88%)	0.65	27 (8%) 15 16	26, 44, 69, 79	0
1	C	314/347 (90%)	0.36	25 (7%) 18 18	20, 37, 65, 80	1 (0%)
1	D	312/347 (89%)	0.44	24 (7%) 19 20	24, 38, 69, 88	0
All	All	1271/1388 (91%)	0.38	95 (7%) 20 20	20, 37, 67, 88	1 (0%)

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	235	GLY	5.5
1	D	245	TYR	5.0
1	D	340	GLY	4.8
1	D	85	SER	4.7
1	D	250	VAL	4.2
1	B	76	ILE	4.1
1	C	177	GLY	4.0
1	D	180	VAL	3.8
1	A	177	GLY	3.7
1	D	201	TYR	3.7
1	A	78	SER	3.6
1	D	63	VAL	3.5
1	C	233	LEU	3.5
1	B	125	HIS	3.5
1	C	234	GLY	3.5
1	A	82	SER	3.4
1	B	75	ILE	3.4
1	D	67	PRO	3.3
1	C	89	LEU	3.3
1	B	182	LEU	3.3
1	C	62	LYS	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	236	GLN	3.2
1	C	179	THR	3.2
1	C	66	VAL	3.2
1	B	73	LEU	3.2
1	D	197	LEU	3.1
1	B	181	GLY	3.1
1	C	87	GLY	3.1
1	B	72	SER	3.0
1	C	63	VAL	3.0
1	A	247	ALA	3.0
1	A	343	LYS	3.0
1	B	46	ALA	2.9
1	C	238	VAL	2.9
1	B	341	ALA	2.9
1	D	4	ASP	2.9
1	C	239	PHE	2.7
1	B	84	LEU	2.7
1	A	76	ILE	2.7
1	C	251	ARG	2.7
1	A	3	GLN	2.7
1	D	118	PHE	2.7
1	D	66	VAL	2.7
1	B	14	GLY	2.6
1	C	2	GLY	2.6
1	A	238	VAL	2.6
1	C	178	SER	2.6
1	C	241	TYR	2.5
1	D	241	TYR	2.5
1	A	243	HIS	2.5
1	B	251	ARG	2.5
1	A	68	PRO	2.5
1	C	118	PHE	2.5
1	B	201	TYR	2.5
1	A	246	GLN	2.5
1	C	88	GLU	2.5
1	D	88	GLU	2.5
1	B	66	VAL	2.4
1	C	232	LEU	2.4
1	A	245	TYR	2.4
1	D	125	HIS	2.4
1	D	90	ALA	2.4
1	D	181	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	75	ILE	2.3
1	C	176	GLN	2.3
1	B	63	VAL	2.3
1	B	19	ARG	2.3
1	A	244	LYS	2.3
1	B	86	ALA	2.3
1	D	89	LEU	2.3
1	B	175	SER	2.2
1	D	288	ARG	2.2
1	A	85	SER	2.2
1	D	233	LEU	2.2
1	C	236	GLN	2.2
1	C	201	TYR	2.2
1	A	203	ASP	2.2
1	D	182	LEU	2.2
1	C	294[A]	ARG	2.2
1	D	203	ASP	2.2
1	D	200	ARG	2.1
1	B	4	ASP	2.1
1	B	3	GLN	2.1
1	A	234	GLY	2.1
1	B	340	GLY	2.1
1	C	180	VAL	2.1
1	B	112	LEU	2.1
1	B	89	LEU	2.1
1	B	232	LEU	2.1
1	B	106	ASP	2.1
1	D	115	ALA	2.1
1	A	179	THR	2.0
1	B	21	VAL	2.0
1	C	237	GLU	2.0
1	B	102	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.