



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 02:36 AM UTC

PDB ID : 3EWM / pdb_00003ewm
Title : Crystal structure of an uncharacterized sugar kinase PH1459 from pyrococcus horikoshii
Authors : Kumar, G.; Eswaramoorthy, S.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2008-10-15
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	
1	B	313	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5025 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized sugar kinase PH1459.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	304	Total	C	N	O	S	Se	0	0	0
			2342	1499	390	447	1	5			
1	B	291	Total	C	N	O	S	Se	0	1	0
			2242	1439	370	427	1	5			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP O59128
A	2	SER	-	expression tag	UNP O59128
A	3	LEU	-	expression tag	UNP O59128
A	306	GLU	-	expression tag	UNP O59128
A	307	GLY	-	expression tag	UNP O59128
A	308	HIS	-	expression tag	UNP O59128
A	309	HIS	-	expression tag	UNP O59128
A	310	HIS	-	expression tag	UNP O59128
A	311	HIS	-	expression tag	UNP O59128
A	312	HIS	-	expression tag	UNP O59128
A	313	HIS	-	expression tag	UNP O59128
B	1	MSE	-	expression tag	UNP O59128
B	2	SER	-	expression tag	UNP O59128
B	3	LEU	-	expression tag	UNP O59128
B	306	GLU	-	expression tag	UNP O59128
B	307	GLY	-	expression tag	UNP O59128
B	308	HIS	-	expression tag	UNP O59128
B	309	HIS	-	expression tag	UNP O59128
B	310	HIS	-	expression tag	UNP O59128
B	311	HIS	-	expression tag	UNP O59128
B	312	HIS	-	expression tag	UNP O59128
B	313	HIS	-	expression tag	UNP O59128


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	269	Total 269	O 269	0	0
2	B	172	Total 172	O 172	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized sugar kinase PH1459



• Molecule 1: Uncharacterized sugar kinase PH1459



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.76Å 79.37Å 82.99Å 90.00° 94.18° 90.00°	Depositor
Resolution (Å)	57.26 – 1.90 57.26 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (57.26-1.90) 99.1 (57.26-1.90)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.190 , 0.239 0.189 , 0.240	Depositor DCC
R_{free} test set	2259 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5025	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/2368	0.82	2/3188 (0.1%)
1	B	0.58	0/2267	0.84	0/3047
All	All	0.62	0/4635	0.83	2/6235 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	105	ASP	CA-C-N	-5.08	114.54	122.16
1	A	105	ASP	C-N-CA	-5.08	114.54	122.16

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2342	0	2442	45	0
1	B	2242	0	2340	47	0
2	A	269	0	0	10	0
2	B	172	0	0	2	0
All	All	5025	0	4782	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:163:LEU:CD1	1:B:175:MSE:HE2	1.78	1.13
1:B:135:LEU:O	1:B:141:ARG:HD3	1.56	1.05
1:B:168:TRP:CD1	1:B:175:MSE:HE3	1.94	1.03
1:B:163:LEU:HD11	1:B:175:MSE:HE2	1.00	0.97
1:B:163:LEU:HD11	1:B:175:MSE:CE	1.96	0.91
1:A:135:LEU:O	1:A:141:ARG:HD3	1.73	0.87
1:A:89:ILE:CD1	1:B:89:ILE:CD1	2.59	0.80
1:A:210:MSE:HG3	1:A:263:LEU:HD21	1.63	0.80
1:A:89:ILE:HD11	1:B:89:ILE:CD1	2.13	0.78
1:A:210:MSE:CG	1:A:263:LEU:HD21	2.17	0.74
1:B:45:ARG:HD3	1:B:72:GLU:OE1	1.89	0.71
1:A:210:MSE:HG3	1:A:263:LEU:CD2	2.21	0.71
1:B:82:ASP:OD1	1:B:84:LYS:HG3	1.90	0.70
1:A:86:HIS:HD2	2:A:576:HOH:O	1.75	0.69
1:B:196:VAL:HG21	1:B:214:ILE:HD12	1.74	0.68
1:A:89:ILE:HD11	1:B:89:ILE:HD11	1.76	0.68
1:B:163:LEU:CD1	1:B:175:MSE:CE	2.68	0.64
1:A:141:ARG:HD2	1:A:178:VAL:HG11	1.80	0.63
1:B:152:LYS:HD2	1:B:157:ILE:HD12	1.80	0.63
1:A:86:HIS:CD2	2:A:576:HOH:O	2.51	0.62
1:B:174:GLU:O	1:B:178:VAL:HG13	2.00	0.61
1:A:210:MSE:HE3	2:A:521:HOH:O	1.99	0.61
1:A:21:ASP:HB3	2:A:346:HOH:O	2.00	0.60
1:B:291:LYS:HG3	1:B:304:LEU:HD13	1.83	0.60
1:B:141:ARG:HD2	1:B:178:VAL:HG11	1.86	0.57
1:B:298:LYS:O	1:B:302:GLU:HG2	2.04	0.57
1:A:152:LYS:HD2	1:A:157:ILE:HD12	1.86	0.57
1:A:128:VAL:HG21	1:A:151:ILE:HD13	1.86	0.57
1:A:11:LEU:HD13	1:A:90:VAL:HG12	1.87	0.56
1:A:89:ILE:HD11	1:B:89:ILE:HD13	1.86	0.56
1:B:82:ASP:OD1	1:B:84:LYS:CG	2.54	0.56
1:A:89:ILE:CD1	1:B:89:ILE:HD13	2.34	0.56
1:B:197:LEU:HD23	2:B:445:HOH:O	2.06	0.55
1:B:32:PRO:HD3	1:B:64:TYR:OH	2.09	0.53
1:B:11:LEU:HD13	1:B:90:VAL:HG12	1.90	0.52
1:B:152:LYS:HD2	1:B:157:ILE:CD1	2.41	0.51
1:B:10:LEU:HG	1:B:65:LEU:HD13	1.92	0.50
1:B:196:VAL:CG2	1:B:214:ILE:HD12	2.40	0.50
1:B:119:TRP:HA	1:B:122:VAL:HG22	1.94	0.50
1:B:15:ILE:HD12	1:B:31:HIS:CD2	2.46	0.50
1:A:296:LYS:HG3	2:A:518:HOH:O	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:LYS:HG3	2:A:520:HOH:O	2.12	0.49
1:B:165:LEU:HD23	1:B:175:MSE:HE1	1.94	0.49
1:A:161:VAL:O	1:A:162:ASN:C	2.56	0.49
1:A:15:ILE:HA	1:A:92:VAL:HG22	1.95	0.48
1:A:32:PRO:HB2	1:A:286:TRP:CZ2	2.49	0.48
1:A:89:ILE:CD1	1:B:89:ILE:HD11	2.36	0.48
1:A:152:LYS:HD2	1:A:157:ILE:CD1	2.44	0.48
1:A:89:ILE:HD13	1:B:89:ILE:CD1	2.39	0.48
1:A:119:TRP:CZ3	1:A:150:LYS:HD3	2.49	0.48
1:A:138:ASN:HD21	1:A:142:GLU:HG3	1.79	0.47
1:B:225:LYS:NZ	1:B:263:LEU:HB3	2.30	0.46
1:B:89:ILE:HG13	1:B:104:TYR:HB2	1.97	0.46
1:A:130:PHE:O	1:A:159:PHE:HA	2.15	0.46
1:A:118:ASN:O	1:A:122:VAL:HG13	2.15	0.46
1:B:149:LYS:HE3	1:B:181:GLU:OE1	2.16	0.46
1:B:226:ASN:HB3	1:B:227:GLU:H	1.62	0.46
1:A:15:ILE:HG12	1:A:92:VAL:CG2	2.46	0.46
1:A:222:ARG:HG3	1:A:229:VAL:HG13	1.97	0.45
1:A:196:VAL:HG21	1:A:214:ILE:HD12	1.98	0.45
1:B:41:VAL:HG21	1:B:286:TRP:CE2	2.52	0.45
1:B:271:PHE:HB2	1:B:303:VAL:HG21	1.98	0.45
1:A:168:TRP:CE3	1:A:175:MSE:HB2	2.51	0.45
1:A:59:ASP:HB2	1:A:60:PRO:CD	2.46	0.44
1:A:166:ASP:HB2	2:A:500:HOH:O	2.17	0.44
1:B:45:ARG:NH2	1:B:286:TRP:O	2.48	0.44
1:B:163:LEU:CD1	1:B:179:LEU:HD11	2.48	0.44
1:A:196:VAL:CG2	1:A:214:ILE:HD12	2.48	0.44
1:B:131:GLY:N	2:B:427:HOH:O	2.40	0.43
1:A:174:GLU:HB2	2:A:364:HOH:O	2.19	0.43
1:B:149:LYS:CE	1:B:181:GLU:OE1	2.67	0.43
1:B:161:VAL:O	1:B:162:ASN:C	2.61	0.43
1:B:3:LEU:HD23	1:B:125:ALA:HA	2.01	0.43
1:A:243:THR:HG21	2:A:578:HOH:O	2.19	0.42
1:A:168:TRP:CH2	1:A:178:VAL:HG21	2.55	0.42
1:A:225:LYS:HE2	1:A:263:LEU:HD23	2.01	0.42
1:A:267:LYS:HZ2	1:A:302:GLU:CD	2.28	0.42
1:A:191:ALA:O	1:A:214:ILE:HA	2.20	0.42
1:B:163:LEU:HD12	1:B:163:LEU:HA	1.87	0.41
1:A:131:GLY:N	2:A:541:HOH:O	2.35	0.41
1:A:138:ASN:ND2	1:A:142:GLU:H	2.17	0.41
1:B:130:PHE:O	1:B:159:PHE:HA	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:84:LYS:HE2	1:B:85:LYS:HE2	2.02	0.40
1:A:18:GLU:OE1	1:A:26:ARG:NH2	2.40	0.40
1:B:41:VAL:HG21	1:B:286:TRP:CD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	302/313 (96%)	298 (99%)	4 (1%)	0	100	100
1	B	286/313 (91%)	278 (97%)	8 (3%)	0	100	100
All	All	588/626 (94%)	576 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/260 (99%)	250 (97%)	8 (3%)	35	29
1	B	246/260 (95%)	237 (96%)	9 (4%)	30	22
All	All	504/520 (97%)	487 (97%)	17 (3%)	32	25

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	VAL
1	A	27	LEU
1	A	122	VAL
1	A	227	GLU
1	A	231	ASP
1	A	263	LEU
1	A	295	LEU
1	A	298	LYS
1	B	3	LEU
1	B	51	SER
1	B	142	GLU
1	B	178	VAL
1	B	201	ASN
1	B	205	GLU
1	B	261	LYS
1	B	263	LEU
1	B	295	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	HIS
1	A	115	ASN
1	A	129	ASN
1	A	138	ASN
1	B	115	ASN
1	B	129	ASN
1	B	138	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/313 (95%)	-0.08	3 (1%) 79 82	7, 14, 28, 36	0
1	B	286/313 (91%)	0.22	10 (3%) 47 50	10, 19, 34, 41	1 (0%)
All	All	585/626 (93%)	0.06	13 (2%) 62 66	7, 17, 32, 41	1 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	218	PRO	5.1
1	B	280	THR	4.3
1	B	197	LEU	3.8
1	A	235	TYR	3.5
1	B	97	ALA	3.2
1	A	2	SER	3.1
1	B	201	ASN	3.0
1	B	218	PRO	2.7
1	B	292	ASP	2.6
1	B	306	GLU	2.5
1	B	305	ALA	2.3
1	B	295	LEU	2.3
1	B	84	LYS	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.