



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 10:12 AM UTC

PDB ID : 5EWU / pdb_00005ewu
Title : Crystal structure of the Arabidopsis thaliana C-terminal Chlh at 1.25Å
Authors : Chen, Z.; Zhang, X.; Liu, Y.; Jiang, L.
Deposited on : 2015-11-21
Resolution : 1.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1693 (1.28-1.24)
Clashscore	190562	1730 (1.28-1.24)
Ramachandran outliers	187476	1695 (1.28-1.24)
Sidechain outliers	187428	1694 (1.28-1.24)
RSRZ outliers	180081	1693 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	5% 91% 8% .
1	B	388	14% 86% 10% ..

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 7148 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Magnesium-chelatase subunit ChlH, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	383	Total	C	N	O	S	0	9	0
			3004	1896	504	588	16			
1	B	377	Total	C	N	O	S	0	2	0
			2893	1823	490	566	14			

- Molecule 2 is BENZOIC ACID (CCD ID: BEZ) (formula: $C_7H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			9	7	2		
2	B	1	Total	C	O	0	0
			9	7	2		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	Mg 2	0	0
3	B	3	Total 3	Mg 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	634	Total 634	O 634	0	0
4	B	594	Total 594	O 594	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Magnesium-chelatase subunit ChlH, chloroplastic



- Molecule 1: Magnesium-chelatase subunit ChlH, chloroplastic



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.10Å 72.78Å 72.91Å 90.00° 109.76° 90.00°	Depositor
Resolution (Å)	50.00 – 1.25 50.00 – 1.25	Depositor EDS
% Data completeness (in resolution range)	89.8 (50.00-1.25) 89.8 (50.00-1.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 1.25Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.169 , 0.184 (Not available) , 0.187	Depositor DCC
R_{free} test set	8309 reflections (4.42%)	wwPDB-VP
Wilson B-factor (Å ²)	11.6	Xtriage
Anisotropy	0.209	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 62.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7148	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BEZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/3080	0.70	0/4170
1	B	0.48	0/2947	0.78	4/3993 (0.1%)
All	All	0.47	0/6027	0.74	4/8163 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	1324	ASP	N-CA-C	-9.39	92.08	108.23
1	B	1323	GLN	N-CA-C	7.07	121.48	112.86
1	B	1199	MET	CG-SD-CE	-6.55	86.50	100.90
1	B	1325	GLU	N-CA-C	6.04	118.64	111.33

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	1323	GLN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3004	0	2925	23	1
1	B	2893	0	2786	43	1
2	A	9	0	5	1	0
2	B	9	0	5	0	0
3	A	2	0	0	0	0
3	B	3	0	0	0	0
4	A	634	0	0	6	1
4	B	594	0	0	16	0
All	All	7148	0	5721	66	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1327:MET:CE	4:B:1953:HOH:O	2.11	0.98
1:B:1327:MET:HE1	4:B:1953:HOH:O	1.61	0.97
1:A:1082:VAL:HB	4:A:1784:HOH:O	1.65	0.93
1:A:1311:ASN:HD21	1:A:1356:ASP:H	1.09	0.93
1:B:1311:ASN:HD21	1:B:1356:ASP:H	1.11	0.92
1:B:1327:MET:HE3	4:B:1804:HOH:O	1.71	0.89
1:B:1366:LYS:HA	4:B:1617:HOH:O	1.71	0.89
2:A:1401:BEZ:H2	4:A:1506:HOH:O	1.73	0.88
1:B:1164:ASN:H	1:B:1168:GLN:NE2	1.72	0.87
1:A:1186:PRO:HA	4:A:1505:HOH:O	1.77	0.83
1:B:1324:ASP:HB2	4:B:1503:HOH:O	1.85	0.75
1:A:1134:ASP:OD1	4:A:1501:HOH:O	2.04	0.74
1:B:1164:ASN:H	1:B:1168:GLN:HE22	1.35	0.73
1:A:1188:ALA:HB3	4:A:1887:HOH:O	1.89	0.71
1:A:1127:GLN:HG2	1:A:1199:MET:HE2	1.74	0.70
1:B:1311:ASN:ND2	1:B:1356:ASP:H	1.89	0.70
1:B:1322:ILE:HD13	4:B:1506:HOH:O	1.93	0.67
1:B:1327:MET:HE3	4:B:1953:HOH:O	1.87	0.66
1:A:1144[A]:SER:OG	1:A:1190:MET:HE2	1.97	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1343:LEU:HD13	1:B:1369:TYR:HA	1.78	0.65
1:B:1324:ASP:CB	4:B:1503:HOH:O	2.44	0.63
1:A:1311:ASN:ND2	1:A:1356:ASP:H	1.89	0.63
1:A:1033:TRP:HE1	1:A:1209:GLN:HE21	1.49	0.60
1:B:1311:ASN:HD21	1:B:1356:ASP:N	1.92	0.59
1:B:1033:TRP:HE1	1:B:1209:GLN:HE21	1.48	0.58
1:B:1369:TYR:HB2	4:B:1617:HOH:O	2.06	0.56
1:B:1281:MET:O	1:B:1288:GLY:HA3	2.06	0.56
1:B:1324:ASP:O	1:B:1327:MET:HG2	2.08	0.54
1:B:1163:TRP:HA	1:B:1168:GLN:HE22	1.73	0.54
1:B:1278:TYR:O	1:B:1282:MET:HG2	2.09	0.52
1:B:995:ALA:N	4:B:1505:HOH:O	2.43	0.51
1:A:1311:ASN:HD21	1:A:1356:ASP:N	1.93	0.51
1:A:1281:MET:HE2	1:A:1281:MET:HA	1.94	0.49
1:B:1275:PRO:HA	1:B:1278:TYR:CZ	2.47	0.49
1:A:1033:TRP:HE1	1:A:1209:GLN:NE2	2.11	0.48
1:B:1033:TRP:HE1	1:B:1209:GLN:NE2	2.12	0.48
1:B:1315:GLU:HG3	4:B:1501:HOH:O	2.14	0.47
1:A:1027:THR:HB	1:A:1204:ALA:HA	1.96	0.47
1:B:1282:MET:HG3	4:B:1953:HOH:O	2.14	0.47
1:A:1312:TRP:CE2	1:B:1021:GLU:HG3	2.50	0.46
1:A:1017:LYS:HG2	1:A:1022:GLY:HA2	1.96	0.46
1:A:1281:MET:O	1:A:1288:GLY:HA3	2.16	0.46
1:B:1324:ASP:HB3	1:B:1327:MET:HG3	1.99	0.45
1:B:1172[B]:MET:HE2	1:B:1172[B]:MET:HB2	1.72	0.43
1:A:1021:GLU:O	1:A:1021:GLU:HG3	2.19	0.42
1:B:1152:ALA:O	1:B:1172[B]:MET:HG2	2.19	0.42
1:B:1318:ASN:ND2	4:B:1502:HOH:O	2.39	0.42
1:B:1278:TYR:CE2	1:B:1320:THR:HG21	2.54	0.42
1:A:1030:LEU:HG	4:A:1784:HOH:O	2.19	0.42
1:B:1137:GLU:CD	4:B:1579:HOH:O	2.63	0.42
1:A:996:ILE:HA	1:A:997:PRO:C	2.45	0.41
1:B:1172[A]:MET:HE2	4:B:2052:HOH:O	2.20	0.41
1:B:1221:VAL:HG21	1:B:1223:HIS:CE1	2.54	0.41
1:B:1228:ASP:OD2	1:B:1306:SER:HB3	2.20	0.41
1:B:1265:ARG:HB2	1:B:1312:TRP:HZ2	1.84	0.41
1:B:1369:TYR:CB	4:B:1617:HOH:O	2.64	0.41
1:B:1097:GLN:HE21	1:B:1097:GLN:H	1.68	0.41
1:B:1107[B]:MET:HE3	1:B:1107[B]:MET:HB3	1.95	0.41
1:B:1026:GLU:HA	1:B:1119:PHE:CD2	2.56	0.41
1:A:1017:LYS:HG3	1:A:1022:GLY:C	2.46	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1082:VAL:O	1:A:1141:ARG:HD2	2.20	0.41
1:B:1099:ASN:HD22	1:B:1181:PHE:HE1	1.67	0.41
1:A:1206:VAL:HA	1:A:1244:SER:O	2.21	0.41
1:B:1027:THR:HB	1:B:1204:ALA:HA	2.03	0.41
1:A:1228:ASP:OD2	1:A:1306:SER:HB3	2.21	0.40
1:B:1343:LEU:HD13	1:B:1369:TYR:CA	2.48	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1060:THR:O	1:B:1168:GLN:NE2[2_656]	2.08	0.12
1:A:1367:GLU:OE2	4:A:1501:HOH:O[1_455]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/388 (100%)	384 (98%)	6 (2%)	0	100	100
1	B	375/388 (97%)	365 (97%)	10 (3%)	0	100	100
All	All	765/776 (99%)	749 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/337 (77%)	258 (100%)	1 (0%)	84	63
1	B	300/337 (89%)	298 (99%)	2 (1%)	76	47
All	All	559/674 (83%)	556 (100%)	3 (0%)	81	56

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1011	ARG
1	B	1097	GLN
1	B	1265	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1209	GLN
1	A	1231	ASN
1	A	1298	ASN
1	A	1311	ASN
1	A	1323	GLN
1	A	1361	ASN
1	A	1371	GLN
1	B	1020	ASN
1	B	1085	ASN
1	B	1097	GLN
1	B	1164	ASN
1	B	1168	GLN
1	B	1209	GLN
1	B	1256	GLN
1	B	1311	ASN
1	B	1318	ASN
1	B	1361	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BEZ	A	1401	-	9,9,9	0.99	1 (11%)	11,11,11	1.49	2 (18%)
2	BEZ	B	1401	-	9,9,9	0.78	0	11,11,11	1.11	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEZ	A	1401	-	-	0/4/4/4	0/1/1/1
2	BEZ	B	1401	-	-	0/4/4/4	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1401	BEZ	O2-C	-2.26	1.23	1.30

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1401	BEZ	O2-C-C1	3.29	123.29	114.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	1401	BEZ	O2-C-O1	-3.10	116.69	123.35
2	B	1401	BEZ	O2-C-C1	2.49	121.22	114.84
2	B	1401	BEZ	O2-C-O1	-2.30	118.41	123.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1401	BEZ	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/388 (98%)	0.01	21 (5%) 30 32	5, 12, 24, 34	9 (2%)
1	B	377/388 (97%)	0.49	54 (14%) 6 7	6, 14, 29, 37	12 (3%)
All	All	760/776 (97%)	0.25	75 (9%) 13 14	5, 13, 26, 37	21 (2%)

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1322	ILE	6.8
1	A	1187	GLY	5.2
1	B	1372	VAL	5.2
1	B	1188	ALA	5.2
1	B	1369	TYR	5.1
1	B	1060	THR	5.1
1	B	1190	MET	4.7
1	A	1186	PRO	4.2
1	B	1286	TYR	4.0
1	A	1188	ALA	4.0
1	B	1370	SER	4.0
1	A	1255	ALA	3.9
1	B	1359	ALA	3.9
1	B	1061	PHE	3.8
1	A	1252	THR	3.8
1	B	1329	ASN	3.8
1	B	1328	LEU	3.7
1	B	1362	ILE	3.7
1	B	1185	ALA	3.7
1	B	1317	ALA	3.7
1	A	1376	ILE	3.6
1	A	1251	THR	3.5
1	A	1253	ALA	3.5
1	B	1187	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1339	PHE	3.4
1	B	1371	GLN	3.4
1	A	1185	ALA	3.4
1	B	1318	ASN	3.3
1	B	1189	GLY	3.3
1	A	1254	ASN	3.2
1	B	1327	MET	3.2
1	B	1284	SER	3.2
1	B	1275	PRO	3.2
1	B	1366	LYS	3.2
1	B	1365	LEU	3.1
1	B	1357	THR	3.0
1	B	1018	LEU	3.0
1	B	1363	GLU	2.9
1	B	1278	TYR	2.9
1	B	1323	GLN	2.9
1	A	995	ALA	2.9
1	A	1060	THR	2.8
1	A	1018	LEU	2.8
1	A	1061	PHE	2.8
1	B	1336	PRO	2.8
1	B	1162	SER	2.7
1	B	1331	LEU	2.7
1	B	1367	GLU	2.6
1	B	1253	ALA	2.6
1	B	1282	MET	2.6
1	A	1041	TYR	2.5
1	B	1321	PHE	2.5
1	B	1319	SER	2.5
1	B	1326	GLU	2.5
1	B	995	ALA	2.4
1	A	994	GLN	2.4
1	B	1181	PHE	2.4
1	B	1347	LEU	2.3
1	A	1373	GLU	2.3
1	B	1273	LEU	2.3
1	B	1191	ALA	2.2
1	B	1280	GLY	2.2
1	A	1372	VAL	2.2
1	B	1062	GLY	2.2
1	B	1358	SER	2.2
1	B	1368	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	996	ILE	2.1
1	B	1324	ASP	2.1
1	B	1283	SER	2.1
1	B	1252	THR	2.1
1	A	1374	ASP	2.1
1	B	1355	TRP	2.1
1	B	1343	LEU	2.0
1	A	1369	TYR	2.0
1	A	1189	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BEZ	A	1401	9/9	0.91	0.08	14,14,16,16	0
2	BEZ	B	1401	9/9	0.95	0.07	11,14,15,16	0
3	MG	B	1402	1/1	0.97	0.07	24,24,24,24	0
3	MG	B	1404	1/1	0.98	0.07	22,22,22,22	0
3	MG	A	1402	1/1	0.99	0.10	23,23,23,23	0
3	MG	A	1403	1/1	0.99	0.03	11,11,11,11	0
3	MG	B	1403	1/1	1.00	0.05	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.