



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:18 PM UTC

PDB ID : 8EWS / pdb_00008ews
Title : Crystal structure of CYP3A4 bound to an inhibitor
Authors : Sevrioukova, I.F.
Deposited on : 2022-10-24
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	487	12% 76% 14% 8%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3780 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 3A4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	0	2	0
			3609	2356	589	639	25			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LEU	deletion	UNP P08684
A	?	-	ILE	deletion	UNP P08684
A	?	-	PRO	deletion	UNP P08684
A	?	-	ASP	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	ALA	deletion	UNP P08684
A	?	-	MET	deletion	UNP P08684
A	?	-	GLU	deletion	UNP P08684
A	?	-	THR	deletion	UNP P08684
A	?	-	TRP	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	ALA	deletion	UNP P08684
A	?	-	VAL	deletion	UNP P08684
A	?	-	SER	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	VAL	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	504	HIS	-	expression tag	UNP P08684
A	505	HIS	-	expression tag	UNP P08684
A	506	HIS	-	expression tag	UNP P08684
A	507	HIS	-	expression tag	UNP P08684

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is {N-[(2,2'-bipyridin]-5-yl-kappa 2 N 1 ,N 1 ')methyl]-3-(pyridin-4-yl)propanamide}bis[2-(quinolin-2-yl-kappaN)-1-benzothiophen-3-yl-kappaC 3]iridium(1+) (CCD ID: X2Q) (formula: C₅₃H₃₈IrN₆OS₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Ir	N	O	S	0	0
			63	53	1	6	1	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	65	Total	O	0	0
			65	65		

- Molecule 1: Cytochrome P450 3A4



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	78.39Å 100.32Å 129.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	79.29 – 2.15 79.28 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.0 (79.29-2.15) 97.9 (79.28-2.15)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.14Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.238 , 0.275 0.238 , 0.273	Depositor DCC
R_{free} test set	1399 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	56.8	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3780	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, X2Q

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.18	0/3702	0.35	1/5003 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	29	SER	CA-C-O	6.24	126.97	119.61

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3609	0	3702	41	0
2	A	43	0	30	4	0
3	A	63	0	0	2	0
4	A	65	0	0	0	0
All	All	3780	0	3732	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:THR:HG21	1:A:46:PHE:H	1.23	1.00
1:A:27:THR:CG2	1:A:46:PHE:H	1.94	0.79
1:A:27:THR:HG21	1:A:46:PHE:N	2.04	0.71
1:A:27:THR:HG22	1:A:30:HIS:CE1	2.27	0.69
1:A:181:MET:HG2	1:A:208:LYS:HD3	1.81	0.62
1:A:275:MET:HE3	1:A:295:LEU:HG	1.82	0.62
2:A:601:HEM:HBB2	2:A:601:HEM:HMB2	1.87	0.57
1:A:172:LEU:HD11	1:A:491:LEU:HD12	1.86	0.57
1:A:102:PHE:HB3	1:A:375:ARG:HB3	1.87	0.56
1:A:120:ILE:HD11	3:A:602:X2Q:C13	2.36	0.55
1:A:55:LYS:NZ	1:A:63:GLU:OE1	2.30	0.55
2:A:601:HEM:HBC2	2:A:601:HEM:HMC1	1.89	0.55
1:A:305:ALA:O	1:A:309:THR:OG1	2.20	0.54
1:A:27:THR:HG21	1:A:45:PRO:HA	1.91	0.52
1:A:468:CYS:SG	1:A:469:LYS:N	2.83	0.52
1:A:32:LEU:HA	1:A:35:LYS:HE2	1.93	0.50
1:A:421:LYS:HD2	1:A:424:LYS:HE3	1.95	0.49
1:A:87:PRO:HG3	1:A:431:ILE:HD11	1.95	0.49
1:A:290:LEU:HD23	1:A:295:LEU:HD12	1.95	0.48
1:A:70:LYS:HG2	1:A:71:VAL:HG23	1.95	0.48
1:A:101:VAL:HG21	1:A:381:VAL:HG11	1.95	0.46
1:A:28:HIS:C	1:A:30:HIS:N	2.73	0.46
1:A:117:ALA:HB1	1:A:301:ILE:HG13	1.97	0.46
1:A:163:GLU:HA	1:A:166:THR:HB	1.98	0.46
1:A:58:CYS:HB3	1:A:399:TYR:CD1	2.51	0.45
1:A:128:ARG:NE	1:A:288:LYS:HB3	2.31	0.45
1:A:30:HIS:HA	1:A:75:TYR:CZ	2.51	0.45
1:A:208:LYS:O	1:A:212:ARG:HG3	2.17	0.45
1:A:188:SER:HA	1:A:271:PHE:HB3	1.99	0.44
1:A:27:THR:HA	1:A:78:GLN:HE22	1.83	0.43
1:A:287:HIS:HB3	1:A:288:LYS:H	1.60	0.43
1:A:339:LEU:HD21	1:A:349:THR:HG21	1.99	0.43
1:A:246:THR:O	1:A:250:ARG:N	2.45	0.43
1:A:436:GLY:HA3	2:A:601:HEM:HBA1	2.00	0.43
1:A:257:LYS:HA	1:A:257:LYS:HD2	1.52	0.43
1:A:128:ARG:HE	1:A:288:LYS:HB3	1.84	0.42
1:A:442:CYS:HB2	2:A:601:HEM:NA	2.34	0.42
3:A:602:X2Q:C10	3:A:602:X2Q:N03	2.83	0.42
1:A:409:THR:HG22	1:A:410:GLU:HG3	2.00	0.42
1:A:101:VAL:HG11	1:A:381:VAL:HG21	2.02	0.42
1:A:28:HIS:C	1:A:30:HIS:H	2.28	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:GLU:HG2	1:A:170:VAL:HG22	2.02	0.41
1:A:215:PHE:CD2	1:A:218:PRO:HD3	2.56	0.41
1:A:127:LYS:HB2	1:A:127:LYS:HE3	1.83	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	441/487 (91%)	420 (95%)	20 (4%)	1 (0%)	43 44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	190	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	410/443 (93%)	384 (94%)	26 (6%)	16 11

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	THR
1	A	28	HIS
1	A	35	LYS
1	A	50	ILE
1	A	66	LYS
1	A	78	GLN
1	A	125	GLU
1	A	131	SER
1	A	160	LEU
1	A	161	ARG
1	A	166	THR
1	A	175	VAL
1	A	187	THR
1	A	196	LEU
1	A	197	ASN
1	A	205	GLU
1	A	208	LYS
1	A	209	LYS
1	A	211	LEU
1	A	239	CYS
1	A	257	LYS
1	A	295	LEU
1	A	351	LEU
1	A	373	LEU
1	A	406	LYS
1	A	459	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	198	ASN
1	A	287	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	601	1,3	50,50,50	1.43	9 (18%)	67,82,82	1.12	4 (5%)
3	X2Q	A	602	2	68,76,76	1.83	10 (14%)	73,122,122	3.66	14 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	601	1,3	-	0/14/54/54	-
3	X2Q	A	602	2	-	4/9/154/154	0/14/14/14

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	X2Q	C54-N53	5.85	1.47	1.33
3	A	602	X2Q	IR50-N03	-5.66	2.01	2.15
3	A	602	X2Q	IR50-N01	-5.54	2.02	2.15
3	A	602	X2Q	IR50-N04	-5.49	2.02	2.15
2	A	601	HEM	FE-NB	3.57	2.05	1.94
2	A	601	HEM	FE-ND	3.36	2.05	1.94
3	A	602	X2Q	C27-C28	3.21	1.40	1.34
3	A	602	X2Q	C40-C39	-3.16	1.38	1.45
2	A	601	HEM	CAC-C3C	3.15	1.55	1.47
2	A	601	HEM	CAB-C3B	3.09	1.55	1.47
2	A	601	HEM	FE-NA	2.94	2.04	1.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	HEM	FE-NC	2.88	2.04	1.95
3	A	602	X2Q	C22-C25	2.58	1.41	1.37
3	A	602	X2Q	C16-C15	-2.51	1.41	1.45
3	A	602	X2Q	C23-C22	-2.27	1.41	1.45
2	A	601	HEM	CMC-C2C	2.12	1.55	1.50
2	A	601	HEM	CMB-C2B	2.11	1.55	1.50
3	A	602	X2Q	C29-C28	-2.02	1.40	1.44
2	A	601	HEM	CMA-C3A	2.00	1.54	1.50

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	X2Q	C26-C25-C22	27.60	120.75	107.74
3	A	602	X2Q	C26-C25-S52	-5.35	128.28	136.58
3	A	602	X2Q	C47-C48-N03	-4.74	119.99	124.62
3	A	602	X2Q	C21-C16-C15	-4.65	124.55	129.67
3	A	602	X2Q	C05-C14-C15	4.16	109.70	107.74
3	A	602	X2Q	C22-C25-S52	-4.16	110.97	115.21
3	A	602	X2Q	C25-C26-N02	3.58	115.08	107.59
3	A	602	X2Q	C23-C22-C25	2.84	113.08	110.09
3	A	602	X2Q	C55-C56-C58	2.79	122.50	112.70
3	A	602	X2Q	C07-C06-C05	-2.75	118.64	122.59
2	A	601	HEM	C4D-ND-C1D	2.53	108.20	105.21
3	A	602	X2Q	C48-C47-C46	2.33	120.08	118.02
2	A	601	HEM	C1B-NB-C4B	2.29	107.92	105.21
2	A	601	HEM	C3D-C4D-ND	-2.29	107.66	110.17
2	A	601	HEM	C2A-C1A-NA	-2.25	107.65	110.15
3	A	602	X2Q	C42-C43-C44	-2.10	118.36	121.20
3	A	602	X2Q	C29-C28-C27	-2.04	118.68	121.52
3	A	602	X2Q	C62-N61-C60	2.01	121.46	116.86

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	602	X2Q	C55-C56-C58-C63
3	A	602	X2Q	C55-C56-C58-C59
3	A	602	X2Q	O57-C54-C55-C56
3	A	602	X2Q	N53-C54-C55-C56


There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	HEM	4	0
3	A	602	X2Q	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	447/487 (91%)	1.02	57 (12%) 7 8	39, 74, 129, 157	2 (0%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	211	LEU	6.1
1	A	226	PHE	4.8
1	A	271	PHE	4.6
1	A	108	PHE	4.3
1	A	241	PHE	4.0
1	A	274	LEU	3.9
1	A	46	PHE	3.7
1	A	189	PHE	3.7
1	A	216	LEU	3.7
1	A	57	PHE	3.6
1	A	304	PHE	3.5
1	A	381	VAL	3.5
1	A	109	GLY	3.3
1	A	407	TYR	3.3
1	A	275	MET	3.2
1	A	242	PRO	3.2
1	A	160	LEU	3.1
1	A	29	SER	3.1
1	A	289	ALA	3.1
1	A	213	PHE	3.0
1	A	295	LEU	3.0
1	A	258	GLU	3.0
1	A	240	VAL	2.9
1	A	222	SER	2.8
1	A	203	PHE	2.8
1	A	475	LEU	2.8
1	A	181	MET	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	215	PHE	2.7
1	A	198	ASN	2.7
1	A	290	LEU	2.6
1	A	276	ILE	2.6
1	A	257	LYS	2.6
1	A	204	VAL	2.6
1	A	272	LEU	2.4
1	A	302	PHE	2.4
1	A	305	ALA	2.4
1	A	210	LEU	2.3
1	A	444	GLY	2.3
1	A	491	LEU	2.3
1	A	149	ILE	2.2
1	A	479	LEU	2.2
1	A	63	GLU	2.2
1	A	319	TYR	2.2
1	A	383	ILE	2.2
1	A	253	VAL	2.2
1	A	322	ALA	2.2
1	A	367	PHE	2.2
1	A	489	VAL	2.2
1	A	50	ILE	2.1
1	A	465	PHE	2.1
1	A	197	ASN	2.1
1	A	225	VAL	2.1
1	A	473	ILE	2.1
1	A	278	SER	2.1
1	A	54	HIS	2.1
1	A	166	THR	2.0
1	A	53	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HEM	A	601	43/43	0.97	0.09	40,49,55,62	0
3	X2Q	A	602	63/63	0.98	0.11	46,67,79,88	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.