



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:09 AM UTC

PDB ID : 3EXS / pdb_00003exs
Title : Crystal structure of KGPDC from Streptococcus mutans in complex with D-R5P
Authors : Li, G.L.; Liu, X.; Wang, K.T.; Li, L.F.; Su, X.D.
Deposited on : 2008-10-17
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	% 83% 10% 6%
1	B	221	85% 10% 5%
1	C	221	74% 17% • 6%
1	D	221	% 82% 14% ••

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6664 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RmpD (Hexulose-6-phosphate synthase).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	208	Total	C	N	O	S	0	0	0
			1563	985	269	302	7			
1	B	211	Total	C	N	O	S	0	0	0
			1573	996	268	302	7			
1	C	208	Total	C	N	O	S	0	0	0
			1563	985	269	302	7			
1	D	218	Total	C	N	O	S	0	1	0
			1640	1034	281	318	7			

- Molecule 2 is RIBULOSE-5-PHOSPHATE (CCD ID: 5RP) (formula: C₅H₁₁O₈P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			14	5	8	1		
2	D	1	Total	C	O	P	0	0
			13	5	7	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	64	Total 64	O 64	0	0
3	B	83	Total 83	O 83	0	0
3	C	79	Total 79	O 79	0	0
3	D	72	Total 72	O 72	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RmpD (Hexulose-6-phosphate synthase)




- Molecule 1: RmpD (Hexulose-6-phosphate synthase)



- Molecule 1: RmpD (Hexulose-6-phosphate synthase)



- Molecule 1: RmpD (Hexulose-6-phosphate synthase)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.15Å 124.25Å 130.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.74 – 2.50 19.74 – 2.50	Depositor EDS
% Data completeness (in resolution range)	90.9 (19.74-2.50) 90.7 (19.74-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.50Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.184 , 0.244 0.182 , 0.242	Depositor DCC
R_{free} test set	1548 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.220	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.015 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6664	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5RP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	1/1585 (0.1%)	0.93	0/2147
1	B	0.75	1/1597 (0.1%)	0.93	0/2167
1	C	0.77	0/1585	0.92	0/2147
1	D	0.79	2/1672 (0.1%)	1.00	3/2268 (0.1%)
All	All	0.77	4/6439 (0.1%)	0.95	3/8729 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	3
1	D	0	1
All	All	0	4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	94	ILE	CA-CB	5.83	1.57	1.54
1	B	216	ILE	CA-CB	5.33	1.61	1.54
1	D	144[A]	ARG	CB-CG	-5.00	1.37	1.52
1	D	144[B]	ARG	CB-CG	-5.00	1.37	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	144[A]	ARG	CA-CB-CG	6.87	127.83	114.10
1	D	144[B]	ARG	CA-CB-CG	6.87	127.83	114.10
1	D	121	ASP	N-CA-C	6.34	119.13	109.62

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	200	THR	Peptide
1	C	202	ALA	Peptide
1	C	203	LYS	Peptide
1	D	141	HIS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1563	0	1565	12	0
1	B	1573	0	1562	18	0
1	C	1563	0	1565	34	0
1	D	1640	0	1624	26	0
2	A	14	0	9	0	0
2	D	13	0	6	0	0
3	A	64	0	0	5	0
3	B	83	0	0	4	0
3	C	79	0	0	5	0
3	D	72	0	0	4	0
All	All	6664	0	6331	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:152:THR:CG2	3:B:256:HOH:O	1.96	1.10
1:B:152:THR:HG23	3:B:256:HOH:O	1.55	1.03
1:D:218:ARG:HG2	1:D:218:ARG:HH11	1.23	1.02
1:D:210:ARG:HH11	1:D:210:ARG:HG3	1.24	1.02
1:C:203:LYS:HA	1:C:205:PRO:HD3	1.41	1.01
1:D:210:ARG:HH11	1:D:210:ARG:CG	1.82	0.91
1:A:215:GLU:OE1	1:A:218:ARG:HD2	1.71	0.89

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:197:ARG:H	1:C:197:ARG:HD3	1.36	0.89
1:C:203:LYS:CA	1:C:205:PRO:HD3	2.07	0.85
1:C:144:ARG:HG2	3:C:269:HOH:O	1.80	0.79
1:B:203:LYS:O	3:B:244:HOH:O	1.98	0.79
1:D:178:SER:H	1:D:181:THR:HG23	1.49	0.76
1:D:218:ARG:HG2	1:D:218:ARG:NH1	1.91	0.76
1:C:197:ARG:H	1:C:197:ARG:CD	1.99	0.75
1:B:152:THR:N	3:B:284:HOH:O	2.19	0.74
1:D:210:ARG:HG3	1:D:210:ARG:NH1	1.90	0.73
1:C:202:ALA:HB3	1:C:203:LYS:HA	1.70	0.72
1:D:142:GLN:HG2	1:D:147:LEU:HD21	1.74	0.70
1:D:178:SER:H	1:D:181:THR:CG2	2.07	0.67
1:C:144:ARG:O	3:C:282:HOH:O	2.15	0.65
1:A:58:ASP:OD2	3:A:258:HOH:O	2.14	0.65
1:D:186:GLU:HG3	1:D:219:ILE:HG23	1.79	0.64
1:A:127:ALA:HA	1:A:130:TRP:CE3	2.33	0.62
1:D:117:GLU:OE1	1:D:141:HIS:NE2	2.31	0.62
1:B:127:ALA:HA	1:B:130:TRP:CE3	2.34	0.62
1:C:202:ALA:CB	1:C:203:LYS:HA	2.29	0.62
1:D:101:ARG:NH1	3:D:231:HOH:O	2.34	0.61
1:B:93:THR:HB	1:B:95:PRO:HD2	1.83	0.60
1:C:117:GLU:OE1	1:C:141:HIS:NE2	2.32	0.60
1:D:17:LEU:HG	1:D:21:ILE:HD12	1.84	0.60
1:C:17:LEU:HG	1:C:21:ILE:HD12	1.84	0.59
1:C:202:ALA:HB3	1:C:205:PRO:HG3	1.84	0.59
1:B:6:PRO:HD2	1:B:216:ILE:HD11	1.83	0.59
1:D:4:GLN:CB	1:D:5:LEU:HA	2.32	0.58
1:D:101:ARG:NH2	3:D:231:HOH:O	2.36	0.58
1:C:197:ARG:HD3	1:C:197:ARG:N	2.16	0.57
1:D:146:ALA:HB1	1:D:151:GLU:HG3	1.88	0.55
1:C:56:PHE:HB3	1:C:59:LYS:HB2	1.89	0.55
1:A:30:GLU:OE1	1:A:210:ARG:HG3	2.08	0.54
1:B:94:ILE:N	1:B:95:PRO:CD	2.71	0.53
1:C:127:ALA:HA	1:C:130:TRP:CE3	2.42	0.53
1:D:127:ALA:HA	1:D:130:TRP:CE3	2.44	0.53
1:B:212:PHE:O	1:B:216:ILE:HG22	2.09	0.52
1:C:154:GLY:N	3:C:243:HOH:O	2.42	0.52
1:B:5:LEU:HD21	1:B:221:GLY:HA2	1.92	0.52
1:A:174:THR:HB	1:A:194:ILE:HB	1.92	0.51
1:D:218:ARG:HH11	1:D:218:ARG:CG	2.05	0.50
1:B:50:GLU:OE1	1:B:81:ARG:NH1	2.37	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:GLU:O	1:A:218:ARG:HG2	2.12	0.50
1:B:216:ILE:C	1:B:216:ILE:HD13	2.37	0.49
1:C:202:ALA:CB	1:C:203:LYS:CA	2.90	0.49
1:A:53:ARG:NH2	1:A:84:ASP:OD2	2.44	0.49
1:C:183:LYS:HA	1:C:219:ILE:HD13	1.94	0.48
1:C:204:ASN:O	1:C:207:GLY:N	2.47	0.48
1:D:101:ARG:CZ	3:D:231:HOH:O	2.61	0.47
1:B:93:THR:CB	1:B:95:PRO:HD2	2.45	0.47
1:C:144:ARG:CG	3:C:269:HOH:O	2.51	0.47
1:C:203:LYS:C	1:C:205:PRO:HD3	2.41	0.46
1:A:5:LEU:N	3:A:264:HOH:O	2.49	0.46
1:C:17:LEU:HG	1:C:21:ILE:CD1	2.45	0.46
1:D:94:ILE:HB	1:D:95:PRO:HD3	1.98	0.46
1:D:97:MET:HB3	1:D:135:ILE:CD1	2.47	0.45
1:C:53:ARG:CZ	1:C:61:ILE:HD12	2.47	0.45
1:C:197:ARG:O	1:C:201:GLU:HG2	2.18	0.45
1:D:87:THR:HB	1:D:117:GLU:HG3	1.98	0.45
1:C:219:ILE:HG22	1:C:219:ILE:O	2.15	0.44
1:D:93:THR:HB	1:D:95:PRO:HD2	1.99	0.44
1:D:142:GLN:CG	1:D:147:LEU:HD21	2.46	0.44
1:A:185:PHE:O	1:A:188:VAL:HB	2.17	0.44
1:B:94:ILE:N	1:B:95:PRO:HD2	2.32	0.43
1:B:56:PHE:HB3	1:B:59:LYS:HB2	1.99	0.43
1:C:182:LEU:HD12	1:C:216:ILE:HG13	2.00	0.43
1:A:5:LEU:N	3:A:249:HOH:O	2.52	0.43
1:C:197:ARG:HA	1:C:200:THR:OG1	2.18	0.43
1:B:17:LEU:HG	1:B:21:ILE:HD12	2.00	0.43
1:D:146:ALA:O	1:D:151:GLU:HB2	2.18	0.42
1:C:161:VAL:HG13	1:C:171:VAL:HG11	2.01	0.42
1:D:56:PHE:HB3	1:D:59:LYS:HB2	2.02	0.42
1:C:7:ASN:ND2	3:C:266:HOH:O	2.53	0.41
1:C:208:ALA:O	1:C:211:ALA:HB3	2.20	0.41
1:C:31:VAL:O	1:C:59:LYS:HE3	2.21	0.41
1:C:97:MET:HB3	1:C:135:ILE:CD1	2.51	0.41
1:C:30:GLU:HG3	1:C:206:ALA:O	2.21	0.41
1:C:27:VAL:HG11	1:C:199:ILE:CG2	2.51	0.41
1:D:178:SER:N	1:D:181:THR:HG23	2.26	0.41
1:A:167:MET:HE3	3:A:251:HOH:O	2.21	0.41
1:B:31:VAL:O	1:B:59:LYS:HE3	2.21	0.41
1:C:102:LYS:HE3	1:C:102:LYS:HA	2.02	0.40
1:A:58:ASP:CG	3:A:258:HOH:O	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	204/221 (92%)	200 (98%)	4 (2%)	0	100	100
1	B	207/221 (94%)	204 (99%)	2 (1%)	1 (0%)	24	43
1	C	204/221 (92%)	197 (97%)	6 (3%)	1 (0%)	24	43
1	D	217/221 (98%)	215 (99%)	2 (1%)	0	100	100
All	All	832/884 (94%)	816 (98%)	14 (2%)	2 (0%)	43	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	153	TRP
1	C	176	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/175 (93%)	154 (94%)	9 (6%)	19	40
1	B	161/175 (92%)	157 (98%)	4 (2%)	42	69
1	C	163/175 (93%)	153 (94%)	10 (6%)	17	35
1	D	170/175 (97%)	158 (93%)	12 (7%)	13	29
All	All	657/700 (94%)	622 (95%)	35 (5%)	20	42

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	VAL
1	A	91	SER
1	A	142	GLN
1	A	144	ARG
1	A	178	SER
1	A	179	VAL
1	A	180	ASP
1	A	203	LYS
1	A	218	ARG
1	B	39	VAL
1	B	152	THR
1	B	180	ASP
1	B	216	ILE
1	C	91	SER
1	C	102	LYS
1	C	142	GLN
1	C	160	LYS
1	C	173	VAL
1	C	178	SER
1	C	182	LEU
1	C	197	ARG
1	C	201	GLU
1	C	217	LYS
1	D	5	LEU
1	D	86	MET
1	D	91	SER
1	D	126	GLN
1	D	158	LEU
1	D	173	VAL
1	D	179	VAL
1	D	181	THR
1	D	183	LYS
1	D	210	ARG
1	D	217	LYS
1	D	218	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	129	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	159	ASN
1	B	7	ASN
1	B	78	ASN
1	B	126	GLN
1	B	128	GLN
1	B	159	ASN
1	C	7	ASN
1	C	78	ASN
1	C	115	GLN
1	C	128	GLN
1	C	129	GLN
1	C	159	ASN
1	D	7	ASN
1	D	78	ASN
1	D	129	GLN
1	D	142	GLN
1	D	159	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	5RP	D	502	-	12,12,13	0.97	0	14,17,18	1.44	2 (14%)
2	5RP	A	501	-	13,13,13	0.75	0	12,18,18	1.61	4 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5RP	D	502	-	-	3/14/14/16	-
2	5RP	A	501	-	-	10/16/16/16	-

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	502	5RP	C2-C3-C5	3.12	121.41	118.13
2	D	502	5RP	O13-C5-C3	-2.74	106.78	111.31
2	A	501	5RP	O8-C7-C6	-2.71	102.14	109.36
2	A	501	5RP	O14-C6-C5	-2.63	105.85	109.67
2	A	501	5RP	O10-P9-O11	2.47	117.06	107.80
2	A	501	5RP	O10-P9-O8	-2.19	100.95	106.67

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	5RP	O1-C2-C3-C5
2	A	501	5RP	O1-C2-C3-O4
2	A	501	5RP	O4-C3-C5-O13
2	A	501	5RP	O13-C5-C6-O14
2	A	501	5RP	C5-C6-C7-O8
2	A	501	5RP	O14-C6-C7-O8
2	A	501	5RP	C7-O8-P9-O11
2	D	502	5RP	O4-C3-C5-O13
2	A	501	5RP	C2-C3-C5-C6
2	A	501	5RP	O4-C3-C5-C6
2	D	502	5RP	C6-C7-O8-P9
2	D	502	5RP	C2-C3-C5-O13
2	A	501	5RP	O13-C5-C6-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	208/221 (94%)	-0.41	2 (0%) 79 76	9, 21, 44, 54	0
1	B	211/221 (95%)	-0.56	0 100 100	8, 20, 38, 42	0
1	C	208/221 (94%)	-0.41	1 (0%) 87 85	7, 18, 49, 56	0
1	D	218/221 (98%)	-0.55	2 (0%) 81 78	8, 18, 34, 40	1 (0%)
All	All	845/884 (95%)	-0.48	5 (0%) 85 83	7, 19, 43, 56	1 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	221	GLY	2.9
1	C	202	ALA	2.6
1	D	144[A]	ARG	2.4
1	A	183	LYS	2.2
1	A	184	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	5RP	A	501	14/14	0.92	0.10	35,43,45,46	0
2	5RP	D	502	13/14	0.98	0.07	22,26,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.