



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:26 AM UTC

PDB ID : 6EXE / pdb_00006exe
Title : Crystal structure of DotM cytoplasmic domain (residues 153-380),R217E
Authors : Meir, A.; Waksman, G.
Deposited on : 2017-11-08
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	6% 77% 14% • 7%
1	B	235	10% 80% 14% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	B	403	-	X	X	-
2	GOL	B	405	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3721 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IcmP (DotM).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1756	1120	307	316	13			
1	B	223	Total	C	N	O	S	0	1	0
			1789	1140	313	323	13			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	146	GLY	-	expression tag	UNP Q5ZYC7
A	147	PRO	-	expression tag	UNP Q5ZYC7
A	148	SER	-	expression tag	UNP Q5ZYC7
A	149	GLY	-	expression tag	UNP Q5ZYC7
A	150	GLY	-	expression tag	UNP Q5ZYC7
A	151	GLY	-	expression tag	UNP Q5ZYC7
A	152	ALA	-	expression tag	UNP Q5ZYC7
A	217	GLU	ARG	engineered mutation	UNP Q5ZYC7
B	146	GLY	-	expression tag	UNP Q5ZYC7
B	147	PRO	-	expression tag	UNP Q5ZYC7
B	148	SER	-	expression tag	UNP Q5ZYC7
B	149	GLY	-	expression tag	UNP Q5ZYC7
B	150	GLY	-	expression tag	UNP Q5ZYC7
B	151	GLY	-	expression tag	UNP Q5ZYC7
B	152	ALA	-	expression tag	UNP Q5ZYC7
B	217	GLU	ARG	engineered mutation	UNP Q5ZYC7

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	68	Total	O	0	0
			68	68		
4	B	66	Total	O	0	0
			66	66		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: IcmP (DotM)



- Molecule 1: IcmP (DotM)



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	118.28Å 118.28Å 66.57Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	53.45 – 2.00 53.45 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (53.45-2.00) 99.5 (53.45-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.223 , 0.264 0.229 , 0.270	Depositor DCC
R_{free} test set	1749 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	33.9	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.052 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3721	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	2/1794 (0.1%)	1.14	5/2424 (0.2%)
1	B	1.06	1/1832 (0.1%)	1.12	2/2479 (0.1%)
All	All	1.07	3/3626 (0.1%)	1.13	7/4903 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	300	VAL	CA-C	5.48	1.59	1.52
1	A	306	PHE	CA-CB	5.08	1.58	1.53
1	B	159	TRP	CG-CD2	-5.02	1.34	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	159	TRP	N-CA-C	7.24	121.81	112.26
1	A	312	VAL	N-CA-CB	6.50	118.16	110.55
1	B	252	VAL	CB-CA-C	-6.06	104.21	111.97
1	A	254	GLY	N-CA-C	5.51	119.16	112.49
1	A	304	SER	N-CA-C	5.47	119.05	112.38
1	A	236	ASP	CA-C-N	5.24	127.55	120.38
1	A	236	ASP	C-N-CA	5.24	127.55	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1756	0	1768	21	0
1	B	1789	0	1801	28	0
2	A	6	0	8	3	0
2	B	24	0	30	11	0
3	B	12	0	13	0	0
4	A	68	0	0	0	0
4	B	66	0	0	1	0
All	All	3721	0	3620	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:159:TRP:HH2	2:B:403:GOL:HO1	0.94	0.87
1:A:246:VAL:O	1:A:250:THR:OG1	1.92	0.87
1:B:230:MET:HG2	1:B:274:VAL:HG21	1.69	0.74
1:B:159:TRP:CZ3	2:B:403:GOL:H12	2.25	0.71
1:B:159:TRP:HH2	2:B:403:GOL:O1	1.73	0.67
1:A:164:THR:HG23	1:A:167:GLU:H	1.63	0.64
1:B:159:TRP:CH2	2:B:403:GOL:H12	2.35	0.61
1:A:166:MET:HE3	1:A:370:LEU:HD21	1.83	0.61
1:B:159:TRP:CH2	2:B:403:GOL:C1	2.86	0.59
1:A:257:ASP:O	1:A:260:VAL:HG12	2.03	0.57
1:A:159:TRP:CZ3	1:B:256:PRO:HD3	2.40	0.57
1:B:300:VAL:HA	2:B:404:GOL:H31	1.88	0.56
1:A:257:ASP:OD1	1:A:259:SER:OG	2.23	0.55
1:B:159:TRP:HZ3	2:B:403:GOL:C3	2.21	0.53
1:A:301:VAL:N	2:A:401:GOL:O1	2.37	0.53
1:A:300:VAL:HA	2:A:401:GOL:O1	2.10	0.52
1:B:255[B]:LYS:HD2	1:B:256:PRO:CD	2.40	0.52
1:B:159:TRP:CH2	1:B:311:PRO:HG3	2.44	0.52
1:A:164:THR:HG22	1:A:167:GLU:CD	2.36	0.50
1:B:195:ILE:HD11	1:B:365:VAL:HG21	1.94	0.50
1:A:159:TRP:CZ3	1:B:251:PHE:HA	2.48	0.49
1:A:195:ILE:HD11	1:A:365:VAL:HG21	1.96	0.48
1:A:340:LYS:O	1:A:344:GLU:HG3	2.14	0.48
1:A:250:THR:HB	1:A:256:PRO:HA	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:159:TRP:CH2	2:B:403:GOL:O1	2.57	0.47
1:B:176:ARG:NH2	2:B:402:GOL:O1	2.48	0.46
1:B:159:TRP:CZ3	2:B:403:GOL:C1	2.96	0.45
1:A:193:ALA:HB2	1:A:378:LEU:HD21	1.98	0.45
1:B:172:TYR:O	1:B:174:LEU:HD13	2.17	0.44
1:A:243:ILE:HD11	1:A:264:VAL:HG21	1.99	0.44
1:A:215:PHE:HB3	1:A:265:MET:HE3	2.00	0.44
1:A:164:THR:HG22	1:A:167:GLU:CG	2.48	0.44
1:A:251:PHE:CD2	1:B:307:LEU:HD21	2.53	0.43
1:B:354:ILE:HD12	1:B:354:ILE:C	2.44	0.43
1:A:296:ARG:NE	2:A:401:GOL:H11	2.34	0.43
1:A:159:TRP:CH2	1:B:256:PRO:HG3	2.53	0.42
1:B:164:THR:HG22	1:B:167:GLU:CG	2.49	0.42
1:B:159:TRP:HZ3	2:B:403:GOL:H32	1.83	0.42
1:B:164:THR:HG21	4:B:554:HOH:O	2.19	0.42
1:A:300:VAL:HG22	1:A:302:PRO:HD3	2.02	0.41
1:B:225:LEU:CD1	1:B:309:LEU:HD22	2.51	0.41
1:B:203:VAL:HA	1:B:206:MET:HE3	2.03	0.41
1:B:202:ARG:HG2	1:B:206:MET:HE2	2.02	0.40
1:B:370:LEU:HB3	1:B:374:GLN:HB2	2.03	0.40
1:B:255[B]:LYS:HD2	1:B:256:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/235 (92%)	205 (95%)	10 (5%)	0	100	100
1	B	222/235 (94%)	212 (96%)	10 (4%)	0	100	100
All	All	437/470 (93%)	417 (95%)	20 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/195 (95%)	175 (95%)	10 (5%)	20	17
1	B	189/195 (97%)	173 (92%)	16 (8%)	10	7
All	All	374/390 (96%)	348 (93%)	26 (7%)	15	10

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	181	LEU
1	A	208	LEU
1	A	255	LYS
1	A	262	ARG
1	A	297	GLU
1	A	310	LYS
1	A	312	VAL
1	A	340	LYS
1	A	376	GLU
1	A	378	LEU
1	B	159	TRP
1	B	161	MET
1	B	174	LEU
1	B	182	LEU
1	B	190	GLU
1	B	208	LEU
1	B	216	GLU
1	B	253	ASP
1	B	255[A]	LYS
1	B	255[B]	LYS
1	B	276	GLU
1	B	307	LEU
1	B	321	ASN
1	B	327	THR
1	B	343	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	356	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	173	ASN
1	A	234	ASN
1	A	269	GLN
1	A	326	GLN
1	B	269	GLN
1	B	321	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	405	-	5,5,5	2.99	1 (20%)	5,5,5	2.37	3 (60%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	B	404	-	5,5,5	0.34	0	5,5,5	0.65	0
2	GOL	A	401	-	5,5,5	0.90	0	5,5,5	2.06	2 (40%)
2	GOL	B	402	-	5,5,5	0.73	0	5,5,5	0.52	0
2	GOL	B	403	-	5,5,5	2.75	1 (20%)	5,5,5	4.59	4 (80%)
3	MES	B	401	-	12,12,12	2.12	1 (8%)	15,16,16	1.72	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	405	-	-	3/4/4/4	-
2	GOL	B	404	-	-	4/4/4/4	-
2	GOL	A	401	-	-	2/4/4/4	-
2	GOL	B	402	-	-	2/4/4/4	-
2	GOL	B	403	-	-	3/4/4/4	-
3	MES	B	401	-	-	6/6/14/14	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	MES	C8-S	-6.76	1.68	1.77
2	B	405	GOL	O2-C2	-6.18	1.25	1.43
2	B	403	GOL	O2-C2	-5.90	1.26	1.43

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	403	GOL	O2-C2-C3	-7.25	79.17	109.18
2	B	403	GOL	C3-C2-C1	5.28	131.16	111.80
3	B	401	MES	O1S-S-C8	4.59	113.67	106.73
2	B	403	GOL	O2-C2-C1	-3.76	93.60	109.18
2	A	401	GOL	O2-C2-C1	-3.64	94.11	109.18
2	B	405	GOL	O2-C2-C1	-3.47	94.81	109.18
2	B	403	GOL	O3-C3-C2	-3.26	95.70	110.38
3	B	401	MES	O2S-S-C8	2.83	111.00	106.73
3	B	401	MES	O2S-S-O1S	-2.77	104.81	113.82
2	B	405	GOL	C3-C2-C1	2.65	121.53	111.80
2	A	401	GOL	O2-C2-C3	2.30	118.69	109.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	405	GOL	O2-C2-C3	-2.24	99.90	109.18

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	GOL	C1-C2-C3-O3
2	B	402	GOL	C1-C2-C3-O3
2	B	405	GOL	C1-C2-C3-O3
2	B	405	GOL	O2-C2-C3-O3
3	B	401	MES	C8-C7-N4-C3
3	B	401	MES	C8-C7-N4-C5
3	B	401	MES	N4-C7-C8-S
3	B	401	MES	C7-C8-S-O1S
2	B	402	GOL	O2-C2-C3-O3
2	B	403	GOL	O1-C1-C2-C3
2	B	404	GOL	O1-C1-C2-C3
2	B	404	GOL	C1-C2-C3-O3
2	B	405	GOL	O1-C1-C2-C3
3	B	401	MES	C7-C8-S-O3S
2	A	401	GOL	O2-C2-C3-O3
2	B	403	GOL	O1-C1-C2-O2
2	B	404	GOL	O2-C2-C3-O3
2	B	404	GOL	O1-C1-C2-O2
3	B	401	MES	C7-C8-S-O2S
2	B	403	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	404	GOL	1	0
2	A	401	GOL	3	0
2	B	402	GOL	1	0
2	B	403	GOL	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	219/235 (93%)	0.45	15 (6%)	23	22	20, 41, 70, 108	0
1	B	223/235 (94%)	0.57	24 (10%)	11	10	21, 43, 82, 99	1 (0%)
All	All	442/470 (94%)	0.51	39 (8%)	15	14	20, 42, 77, 108	1 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	256	PRO	5.7
1	A	180	ALA	5.0
1	B	300	VAL	4.9
1	A	182	LEU	4.6
1	B	182	LEU	4.3
1	A	159	TRP	4.0
1	A	181	LEU	3.9
1	B	255[A]	LYS	3.6
1	B	157	GLY	3.5
1	B	257	ASP	3.4
1	B	253	ASP	3.3
1	A	300	VAL	3.3
1	B	250	THR	3.2
1	A	254	GLY	3.2
1	B	158	PRO	3.0
1	A	160	ALA	3.0
1	B	186	VAL	3.0
1	B	187	PRO	3.0
1	B	307	LEU	2.9
1	A	158	PRO	2.9
1	B	159	TRP	2.8
1	B	254	GLY	2.7
1	A	256	PRO	2.7
1	A	179	ASP	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	249	LYS	2.7
1	B	181	LEU	2.7
1	B	258	PHE	2.6
1	B	251	PHE	2.6
1	B	160	ALA	2.5
1	A	373	ARG	2.4
1	B	244	LEU	2.4
1	B	243	ILE	2.4
1	B	180	ALA	2.3
1	A	299	GLY	2.3
1	A	304	SER	2.2
1	B	239	ALA	2.2
1	A	307	LEU	2.1
1	A	252	VAL	2.1
1	B	304	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	B	404	6/6	0.75	0.20	53,59,71,83	0
2	GOL	B	402	6/6	0.76	0.16	39,43,49,54	0
3	MES	B	401	12/12	0.78	0.19	86,90,105,107	0
2	GOL	A	401	6/6	0.83	0.15	46,50,56,63	0
2	GOL	B	405	6/6	0.88	0.17	14,32,37,37	0
2	GOL	B	403	6/6	0.92	0.18	19,37,49,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.