



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 02:31 PM UTC

PDB ID : 7EXJ / pdb_00007exj
Title : Crystal structure of alkaline alpha-galctosidase D383A mutant from Arabidopsis thaliana complexed with Raffinose
Authors : Chuankhayan, P.; Guan, H.H.; Lin, C.C.; Chen, N.C.; Huang, Y.C.; Yoshimura, M.; Nakagawa, A.; Lee, R.H.; Chen, C.J.
Deposited on : 2021-05-27
Resolution : 2.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	749	 81% 13% . .
1	B	749	 82% 13% . .
2	C	3	 67% 33%
2	D	3	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FRU	C	1	X	-	-	-
2	GLC	C	2	X	-	-	-
2	GLA	C	3	X	-	-	-
2	FRU	D	1	X	-	-	-
2	GLC	D	2	X	-	-	-
2	GLA	D	3	X	-	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 11324 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable galactinol–sucrose galactosyltransferase 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	718	Total	C	N	O	S	0	0	0
			5620	3573	969	1049	29			
1	A	718	Total	C	N	O	S	0	0	0
			5620	3573	969	1049	29			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	302	ARG	LYS	conflict	UNP Q8RX87
B	383	ALA	ASP	engineered mutation	UNP Q8RX87
A	302	ARG	LYS	conflict	UNP Q8RX87
A	383	ALA	ASP	engineered mutation	UNP Q8RX87

- Molecule 2 is an oligosaccharide called alpha-D-galactopyranose-(1-6)-alpha-D-glucopyranoside-(1-2)-beta-D-fructofuranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	3	Total	C	O	0	0	0
			34	18	16			
2	D	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	9	Total	O	0	0
			9	9		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	7	Total	O	0	0
			7	7		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable galactinol–sucrose galactosyltransferase 6



- Molecule 1: Probable galactinol–sucrose galactosyltransferase 6



- Molecule 2: alpha-D-galactopyranose-(1-6)-alpha-D-glucopyranose-(1-2)-beta-D-fructofuranose





- Molecule 2: alpha-D-galactopyranose-(1-6)-alpha-D-glucopyranose-(1-2)-beta-D-fructofuranose

Chain D:

100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	92.60Å 103.30Å 181.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.47 30.00 – 2.47	Depositor EDS
% Data completeness (in resolution range)	99.3 (30.00-2.47) 99.3 (30.00-2.47)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.191 , 0.249 0.199 , 0.193	Depositor DCC
R_{free} test set	3138 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	62.5	Xtriage
Anisotropy	0.156	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 26.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11324	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLA, GLC, FRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.84	0/5754	0.99	7/7793 (0.1%)
1	B	0.98	0/5754	1.06	15/7793 (0.2%)
All	All	0.91	0/11508	1.03	22/15586 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	37	ALA	N-CA-C	8.50	121.68	111.82
1	A	387	VAL	CB-CA-C	-7.87	101.73	112.04
1	A	124	VAL	CB-CA-C	-7.49	100.52	111.68
1	B	387	VAL	CB-CA-C	-7.08	102.59	112.14
1	B	387	VAL	N-CA-CB	6.60	119.52	110.54
1	A	398	ARG	N-CA-C	6.21	118.05	111.28
1	A	387	VAL	N-CA-CB	6.18	118.20	110.47
1	B	124	VAL	CB-CA-C	-6.07	101.59	110.81
1	B	41	GLU	N-CA-C	5.97	118.57	111.71
1	B	475	GLN	CA-C-N	-5.89	113.90	119.85
1	B	475	GLN	C-N-CA	-5.89	113.90	119.85
1	B	708	LYS	N-CA-C	-5.70	104.44	111.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	738	GLU	CA-C-N	5.59	131.76	121.70
1	A	738	GLU	C-N-CA	5.59	131.76	121.70
1	B	738	GLU	CA-C-N	5.49	131.59	121.70
1	B	738	GLU	C-N-CA	5.49	131.59	121.70
1	B	121	VAL	N-CA-C	5.46	115.81	108.17
1	B	612	THR	CB-CA-C	-5.42	101.41	110.56
1	B	211	TRP	N-CA-C	5.36	117.54	111.11
1	B	505	SER	N-CA-CB	-5.31	103.11	111.39
1	A	674	GLY	N-CA-C	5.20	120.39	114.67
1	B	35	SER	N-CA-C	-5.05	102.68	109.95

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	11	ASP	Peptide
1	B	34	ALA	Peptide
1	B	740	LYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5620	0	5539	37	0
1	B	5620	0	5539	43	0
2	C	34	0	28	1	0
2	D	34	0	29	1	0
3	A	7	0	0	0	0
3	B	9	0	0	0	0
All	All	11324	0	11135	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:95:GLN:HE22	1:B:429:ASN:HA	1.49	0.76
1:A:558:ASN:HD21	1:A:672:ASN:HD21	1.33	0.75
1:B:558:ASN:HD21	1:B:672:ASN:HD21	1.37	0.72
1:B:627:LEU:HD21	1:B:702:TYR:HB2	1.76	0.67
1:B:738:GLU:HB3	1:B:739:ASN:C	2.23	0.64
1:B:95:GLN:HE21	1:B:134:ARG:HH21	1.47	0.62
1:B:492:ALA:HB1	1:B:519:LEU:HD11	1.84	0.60
1:A:737:ILE:O	1:A:738:GLU:HB2	2.02	0.58
1:A:7:VAL:HG13	1:A:16:ILE:CD1	2.33	0.58
1:A:95:GLN:HE21	1:A:134:ARG:HH21	1.50	0.58
1:A:442:ILE:HD12	1:A:474:MET:HE3	1.86	0.57
1:B:463:VAL:HG13	1:B:478:TRP:CE2	2.40	0.56
1:A:307:TRP:HE1	2:C:3:GLA:H4	1.71	0.55
1:B:693:VAL:HG21	1:B:745:ILE:HD12	1.90	0.54
1:B:495:ARG:HB3	1:B:502:LEU:HD22	1.90	0.54
1:B:632:TYR:O	1:B:633:ASN:HB2	2.08	0.53
1:A:738:GLU:HB3	1:A:740:LYS:N	2.24	0.53
1:A:95:GLN:HE22	1:A:429:ASN:HA	1.74	0.53
1:A:495:ARG:HB3	1:A:502:LEU:HD22	1.89	0.52
1:A:384:VAL:O	1:A:387:VAL:HG22	2.10	0.51
2:D:1:FRU:H61	2:D:2:GLC:H62	1.92	0.51
1:A:233:GLY:HA3	1:A:513:PHE:CE1	2.45	0.51
1:B:77:TRP:CD2	1:B:426:MET:HE2	2.46	0.50
1:B:496:ALA:HB2	1:B:519:LEU:CD2	2.41	0.50
1:A:266:ILE:HD12	1:A:347:VAL:HG12	1.92	0.50
1:B:502:LEU:HD21	1:B:519:LEU:HD22	1.94	0.50
1:A:95:GLN:HE21	1:A:134:ARG:NH2	2.11	0.49
1:B:95:GLN:HE21	1:B:134:ARG:NH2	2.10	0.49
1:B:7:VAL:HG22	1:B:16:ILE:HD12	1.94	0.49
1:B:693:VAL:CG2	1:B:745:ILE:HD12	2.43	0.49
1:A:77:TRP:HB2	1:A:78:TRP:CD1	2.49	0.48
1:A:358:PRO:HB3	1:A:404:GLN:OE1	2.14	0.48
1:B:384:VAL:HG12	1:B:387:VAL:HG22	1.96	0.47
1:A:96:PHE:HB3	1:A:469:PHE:CE2	2.50	0.47
1:A:558:ASN:ND2	1:A:672:ASN:HD21	2.08	0.47
1:A:40:VAL:HG21	1:A:183:THR:CG2	2.46	0.46
1:A:83:MET:HE2	1:A:543:ASP:HA	1.97	0.46
1:B:172:ASP:C	1:B:172:ASP:OD1	2.59	0.46
1:A:651:SER:OG	1:A:664:PRO:O	2.23	0.45
1:B:124:VAL:O	1:B:167:ILE:HA	2.17	0.45
1:B:266:ILE:HD12	1:B:347:VAL:HG12	1.98	0.45
1:B:572:ALA:O	1:B:573:ALA:HB2	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:560:TYR:C	1:B:606:ALA:HB1	2.42	0.44
1:A:489:GLU:OE1	1:A:671:TYR:OH	2.36	0.44
1:B:7:VAL:HG13	1:B:16:ILE:CD1	2.47	0.44
1:B:737:ILE:O	1:B:738:GLU:HB2	2.17	0.44
1:A:96:PHE:HB3	1:A:469:PHE:CD2	2.52	0.44
1:B:68:PHE:CD1	1:B:68:PHE:C	2.95	0.43
1:B:126:LEU:C	1:B:165:LEU:HD12	2.44	0.43
1:A:194:ARG:HD3	1:A:471:GLY:O	2.19	0.43
1:B:77:TRP:HB2	1:B:78:TRP:CD1	2.54	0.43
1:B:515:LEU:C	1:B:515:LEU:HD23	2.43	0.43
1:A:134:ARG:NH1	1:A:136:CYS:SG	2.92	0.43
1:B:332:MET:HB3	1:B:337:VAL:HG21	2.01	0.43
1:A:286:VAL:HG13	1:A:289:LYS:HE2	2.01	0.43
1:A:521:LEU:HD13	1:A:695:GLY:HA2	2.01	0.43
1:B:37:ALA:N	1:B:38:GLY:HA2	2.34	0.43
1:B:190:SER:O	1:B:435:CYS:HA	2.18	0.43
1:A:378:ASP:O	1:A:417:PHE:HB3	2.19	0.42
1:B:738:GLU:HG3	1:B:739:ASN:HB2	2.02	0.42
1:A:738:GLU:HB3	1:A:739:ASN:C	2.44	0.42
1:B:491:HIS:O	1:B:495:ARG:HG2	2.20	0.42
1:B:491:HIS:HD1	1:B:495:ARG:HH12	1.67	0.42
1:A:40:VAL:HG21	1:A:183:THR:HG23	2.01	0.42
1:B:9:ILE:HD13	1:B:25:VAL:HG11	2.02	0.42
1:A:135:SER:HB3	1:A:147:LEU:HD11	2.00	0.42
1:A:510:LYS:HA	1:A:510:LYS:HE2	2.02	0.42
1:B:627:LEU:HD22	1:B:727:LEU:HD13	2.00	0.42
1:A:463:VAL:HG22	1:A:478:TRP:CD1	2.55	0.41
1:B:208:TRP:O	1:B:240:VAL:HA	2.21	0.41
1:A:166:TYR:CZ	1:A:183:THR:HG21	2.56	0.41
1:A:281:LYS:O	1:A:282:ASP:CB	2.68	0.41
1:B:185:LYS:NZ	1:B:193:GLN:HE21	2.19	0.41
1:B:92:TYR:CE1	1:B:338:GLU:HG2	2.56	0.41
1:B:92:TYR:CD1	1:B:338:GLU:HG2	2.56	0.40
1:B:208:TRP:CH2	1:B:226:GLY:HA3	2.56	0.40
1:B:249:VAL:HG12	1:B:269:LEU:HA	2.02	0.40
1:A:207:GLY:HA3	1:A:239:PHE:CE2	2.56	0.40
1:A:281:LYS:O	1:A:282:ASP:HB2	2.22	0.40
1:B:166:TYR:CE1	1:B:168:HIS:HB2	2.57	0.40
1:A:717:ILE:O	1:A:717:ILE:HD12	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	712/749 (95%)	660 (93%)	43 (6%)	9 (1%)	9	16
1	B	712/749 (95%)	669 (94%)	37 (5%)	6 (1%)	16	28
All	All	1424/1498 (95%)	1329 (93%)	80 (6%)	15 (1%)	11	20

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	6	ALA
1	A	393	GLY
1	A	6	ALA
1	A	282	ASP
1	B	282	ASP
1	A	738	GLU
1	B	446	ASP
1	A	11	ASP
1	A	36	GLU
1	A	38	GLY
1	B	347	VAL
1	B	603	ILE
1	B	38	GLY
1	A	657	VAL
1	A	600	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	614/640 (96%)	565 (92%)	49 (8%)	11	21
1	B	614/640 (96%)	579 (94%)	35 (6%)	18	36
All	All	1228/1280 (96%)	1144 (93%)	84 (7%)	14	28

All (84) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	40	VAL
1	B	43	VAL
1	B	69	MET
1	B	124	VAL
1	B	149	LEU
1	B	164	SER
1	B	186	LEU
1	B	198	LYS
1	B	293	LYS
1	B	301	LEU
1	B	311	THR
1	B	344	LYS
1	B	347	VAL
1	B	359	LYS
1	B	384	VAL
1	B	387	VAL
1	B	388	LEU
1	B	411	SER
1	B	460	ILE
1	B	463	VAL
1	B	489	GLU
1	B	502	LEU
1	B	549	VAL
1	B	561	THR
1	B	585	GLN
1	B	586	THR
1	B	587	LYS
1	B	588	THR
1	B	622	GLN
1	B	636	LEU
1	B	661	SER
1	B	729	THR
1	B	732	LEU
1	B	734	LYS
1	B	741	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	32	THR
1	A	33	SER
1	A	35	SER
1	A	58	VAL
1	A	67	ARG
1	A	69	MET
1	A	70	SER
1	A	120	LYS
1	A	124	VAL
1	A	148	CYS
1	A	150	GLU
1	A	186	LEU
1	A	198	LYS
1	A	217	GLU
1	A	243	ASP
1	A	301	LEU
1	A	321	GLU
1	A	359	LYS
1	A	384	VAL
1	A	387	VAL
1	A	388	LEU
1	A	452	ASP
1	A	460	ILE
1	A	463	VAL
1	A	502	LEU
1	A	515	LEU
1	A	550	SER
1	A	559	LYS
1	A	576	SER
1	A	587	LYS
1	A	588	THR
1	A	589	ASP
1	A	590	SER
1	A	595	ILE
1	A	622	GLN
1	A	633	ASN
1	A	639	SER
1	A	658	ASP
1	A	694	LYS
1	A	708	LYS
1	A	716	GLU
1	A	723	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	729	THR
1	A	732	LEU
1	A	733	ASP
1	A	734	LYS
1	A	737	ILE
1	A	738	GLU
1	A	746	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	65	ASN
1	B	95	GLN
1	B	168	HIS
1	B	187	HIS
1	B	193	GLN
1	B	331	ASN
1	B	385	GLN
1	B	406	HIS
1	B	456	HIS
1	B	482	HIS
1	B	558	ASN
1	B	645	HIS
1	A	28	ASN
1	A	95	GLN
1	A	142	ASN
1	A	175	GLN
1	A	195	HIS
1	A	247	GLN
1	A	331	ASN
1	A	351	GLN
1	A	385	GLN
1	A	456	HIS
1	A	482	HIS
1	A	558	ASN
1	A	584	HIS
1	A	739	ASN
1	A	743	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FRU	C	1	2	11,12,12	3.94	8 (72%)	10,18,18	1.00	0
2	GLC	C	2	2	11,11,12	3.01	6 (54%)	15,15,17	1.84	3 (20%)
2	GLA	C	3	2	11,11,12	2.28	2 (18%)	15,15,17	3.02	8 (53%)
2	FRU	D	1	2	11,12,12	4.10	8 (72%)	10,18,18	1.24	2 (20%)
2	GLC	D	2	2	11,11,12	3.18	6 (54%)	15,15,17	2.22	5 (33%)
2	GLA	D	3	2	11,11,12	2.05	2 (18%)	15,15,17	2.03	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRU	C	1	2	1/1/4/4	3/5/24/24	0/1/1/1
2	GLC	C	2	2	1/1/4/5	1/2/19/22	0/1/1/1
2	GLA	C	3	2	3/3/4/5	0/2/19/22	0/1/1/1
2	FRU	D	1	2	1/1/4/4	1/5/24/24	0/1/1/1
2	GLC	D	2	2	1/1/4/5	2/2/19/22	0/1/1/1
2	GLA	D	3	2	3/3/4/5	0/2/19/22	0/1/1/1

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	GLC	C2-C3	-6.62	1.42	1.52
2	D	1	FRU	C4-C5	6.34	1.69	1.53
2	C	1	FRU	C4-C3	-6.33	1.27	1.53
2	D	1	FRU	C4-C3	-6.23	1.28	1.53
2	C	1	FRU	C4-C5	5.74	1.67	1.53
2	C	2	GLC	O5-C5	5.67	1.54	1.43
2	D	1	FRU	O5-C2	5.54	1.52	1.43
2	C	2	GLC	C2-C3	-5.48	1.44	1.52
2	D	2	GLC	O5-C5	5.20	1.53	1.43
2	C	1	FRU	O5-C2	5.15	1.51	1.43
2	D	3	GLA	C2-C3	-4.90	1.45	1.52
2	C	3	GLA	O5-C1	4.89	1.51	1.43
2	C	3	GLA	C2-C3	-4.71	1.45	1.52
2	D	1	FRU	O5-C5	-4.63	1.33	1.43
2	D	1	FRU	C1-C2	-4.44	1.43	1.52
2	C	1	FRU	O5-C5	-4.13	1.34	1.43
2	C	1	FRU	O2-C2	4.13	1.47	1.40
2	C	1	FRU	C1-C2	-4.01	1.44	1.52
2	C	2	GLC	O3-C3	3.71	1.52	1.43
2	D	3	GLA	O5-C1	3.70	1.49	1.43
2	D	2	GLC	O5-C1	3.54	1.49	1.43
2	C	1	FRU	O3-C3	3.50	1.49	1.42
2	D	1	FRU	O3-C3	3.32	1.49	1.42
2	D	1	FRU	O4-C4	3.27	1.51	1.43
2	D	2	GLC	O3-C3	3.17	1.50	1.43
2	D	1	FRU	O2-C2	3.15	1.46	1.40
2	D	2	GLC	O2-C2	3.08	1.49	1.43
2	C	2	GLC	C4-C5	2.99	1.59	1.53
2	C	2	GLC	O2-C2	2.90	1.49	1.43
2	C	1	FRU	O4-C4	2.75	1.49	1.43
2	D	2	GLC	C4-C5	2.55	1.58	1.53
2	C	2	GLC	O5-C1	2.35	1.47	1.43

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	GLA	C1-C2-C3	8.00	121.29	109.64
2	D	2	GLC	C1-O5-C5	5.52	119.59	112.19
2	C	2	GLC	C1-O5-C5	4.12	117.71	112.19
2	D	3	GLA	O3-C3-C2	-4.11	101.67	110.05
2	C	2	GLC	C3-C4-C5	3.96	117.40	110.23
2	C	3	GLA	O2-C2-C1	3.94	118.23	109.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	GLA	O3-C3-C2	-3.92	102.06	110.05
2	D	2	GLC	C3-C4-C5	3.80	117.13	110.23
2	C	3	GLA	O2-C2-C3	-3.54	102.82	110.15
2	D	3	GLA	C1-C2-C3	3.28	114.42	109.64
2	D	2	GLC	O5-C1-C2	2.99	117.92	110.79
2	D	3	GLA	C1-O5-C5	2.85	116.01	112.19
2	C	3	GLA	C2-C3-C4	2.82	115.82	110.86
2	D	2	GLC	O5-C5-C6	2.65	112.81	107.66
2	D	3	GLA	O5-C5-C4	-2.50	104.74	110.83
2	C	3	GLA	O4-C4-C3	-2.48	104.53	110.38
2	D	3	GLA	C2-C3-C4	2.37	115.03	110.86
2	C	2	GLC	C6-C5-C4	2.30	118.67	113.02
2	D	1	FRU	O4-C4-C5	2.19	117.38	111.08
2	D	2	GLC	O6-C6-C5	-2.13	104.07	111.33
2	D	1	FRU	C5-C4-C3	2.07	108.51	102.07
2	C	3	GLA	O4-C4-C5	2.05	114.38	109.32
2	C	3	GLA	C1-O5-C5	2.01	114.89	112.19

All (10) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	1	FRU	C4
2	C	2	GLC	C5
2	C	3	GLA	C4
2	C	3	GLA	C2
2	C	3	GLA	C3
2	D	1	FRU	C4
2	D	2	GLC	C5
2	D	3	GLA	C4
2	D	3	GLA	C2
2	D	3	GLA	C3

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	FRU	O1-C1-C2-C3
2	C	1	FRU	O1-C1-C2-O2
2	C	1	FRU	O1-C1-C2-O5
2	D	2	GLC	O5-C5-C6-O6
2	C	2	GLC	O5-C5-C6-O6
2	D	1	FRU	C4-C5-C6-O6
2	D	2	GLC	C4-C5-C6-O6


There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2	GLC	1	0
2	D	1	FRU	1	0
2	C	3	GLA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	718/749 (95%)	-0.15	3 (0%) 88 87	48, 73, 111, 143	0
1	B	718/749 (95%)	-0.27	2 (0%) 90 88	40, 62, 94, 132	0
All	All	1436/1498 (95%)	-0.21	5 (0%) 90 88	40, 68, 106, 143	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	253	ALA	2.5
1	A	6	ALA	2.3
1	A	253	ALA	2.2
1	B	37	ALA	2.1
1	A	740	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FRU	C	1	12/12	0.85	0.11	79,99,111,114	0
2	GLC	C	2	11/12	0.89	0.09	76,86,104,110	0
2	FRU	D	1	12/12	0.95	0.07	53,69,73,75	0
2	GLA	D	3	11/12	0.95	0.09	51,56,63,85	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	D	2	11/12	0.96	0.06	51,61,65,67	0
2	GLA	C	3	11/12	0.97	0.06	52,59,63,66	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.