



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:40 AM UTC

PDB ID : 8EXU / pdb_00008exu
Title : Crystal structure of PI3K-alpha in complex with compound 30
Authors : Kiefer, J.R.; Eigenbrot, C.; Staben, S.T.; Hanan, E.J.; Wallweber, H.J.A.;
Ultsch, M.; Braun, M.G.; Friedman, L.S.; Purkey, H.E.
Deposited on : 2022-10-25
Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1080	5% 74% 9% 17%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7495 atoms, of which 2 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit alpha isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	898	Total	C	N	O	S	0	3	0
			7371	4704	1268	1336	63			

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-27	MET	-	initiating methionine	UNP P42336
A	-26	SER	-	expression tag	UNP P42336
A	-25	TYR	-	expression tag	UNP P42336
A	-24	TYR	-	expression tag	UNP P42336
A	-23	HIS	-	expression tag	UNP P42336
A	-22	HIS	-	expression tag	UNP P42336
A	-21	HIS	-	expression tag	UNP P42336
A	-20	HIS	-	expression tag	UNP P42336
A	-19	HIS	-	expression tag	UNP P42336
A	-18	HIS	-	expression tag	UNP P42336
A	-17	ASP	-	expression tag	UNP P42336
A	-16	TYR	-	expression tag	UNP P42336
A	-15	ASP	-	expression tag	UNP P42336
A	-14	ILE	-	expression tag	UNP P42336
A	-13	PRO	-	expression tag	UNP P42336
A	-12	THR	-	expression tag	UNP P42336
A	-11	THR	-	expression tag	UNP P42336
A	-10	GLU	-	expression tag	UNP P42336
A	-9	ASN	-	expression tag	UNP P42336
A	-8	LEU	-	expression tag	UNP P42336
A	-7	TYR	-	expression tag	UNP P42336
A	-6	PHE	-	expression tag	UNP P42336
A	-5	GLN	-	expression tag	UNP P42336
A	-4	GLY	-	expression tag	UNP P42336
A	-3	ALA	-	expression tag	UNP P42336
A	-2	MET	-	expression tag	UNP P42336

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P42336
A	99	GLU	-	insertion	UNP P42336
A	100	ASN	-	insertion	UNP P42336
A	101	LEU	-	insertion	UNP P42336
A	102	TYR	-	insertion	UNP P42336
A	103	PHE	-	insertion	UNP P42336
A	104	GLN	-	insertion	UNP P42336
A	105	GLY	-	insertion	UNP P42336

- Molecule 2 is (2S)-2-cyclopropyl-2-({(4S,11aM)-2-[(4S)-2-oxo-4-(trifluoromethyl)-1,3-oxazolidin-3-yl]-5,6-dihydroimidazo[1,2-d][1,4]benzoxazepin-9-yl}amino)acetamide (CCD ID: X3R) (formula: C₂₀H₂₀F₃N₅O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	H	N	O	
			34	20	3	2	5	4	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	90	Total	O		
			90	90	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.64Å 133.72Å 140.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.46 – 2.68 29.46 – 2.68	Depositor EDS
% Data completeness (in resolution range)	92.6 (29.46-2.68) 92.9 (29.46-2.68)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.68Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.196 , 0.241 0.196 , 0.251	Depositor DCC
R_{free} test set	731 reflections (2.47%)	wwPDB-VP
Wilson B-factor (Å ²)	56.2	Xtriage
Anisotropy	0.285	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7495	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: X3R

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/7538	1.28	11/10181 (0.1%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	261	PHE	CA-CB-CG	6.56	120.36	113.80
1	A	603	ASP	CA-CB-CG	6.42	119.02	112.60
1	A	905	CYS	CA-C-N	5.71	127.86	120.56
1	A	905	CYS	C-N-CA	5.71	127.86	120.56
1	A	124	PRO	CA-C-N	5.49	128.26	120.53
1	A	124	PRO	C-N-CA	5.49	128.26	120.53
1	A	506	PHE	CA-CB-CG	5.29	119.09	113.80
1	A	937	PHE	CA-CB-CG	5.27	119.07	113.80
1	A	926	ASP	CA-CB-CG	5.10	117.70	112.60
1	A	843	ASP	CA-CB-CG	5.04	117.64	112.60
1	A	133	ASP	CA-CB-CG	5.02	117.62	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7371	0	7345	39	0
2	A	32	2	0	0	0
3	A	90	0	0	0	0
All	All	7493	2	7345	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:ASN:HD21	1:A:651:ARG:HH12	1.22	0.86
1:A:808:ARG:HB3	1:A:1010:MET:HE2	1.78	0.66
1:A:199:SER:HB3	1:A:203:ASP:HB2	1.80	0.63
1:A:145:ASN:HD21	1:A:651:ARG:NH1	1.94	0.62
1:A:809:GLN:HE22	1:A:1012:GLU:HG3	1.65	0.60
1:A:220:VAL:HG21	1:A:267:LEU:HD13	1.85	0.56
1:A:337:LYS:HB3	1:A:476:GLU:HB3	1.87	0.56
1:A:640:LYS:HE2	1:A:680:VAL:HG11	1.88	0.56
1:A:194:ILE:HD11	1:A:220:VAL:HG12	1.86	0.56
1:A:807:LEU:HD12	1:A:846:GLY:HA3	1.91	0.53
1:A:712:LEU:HG	1:A:748:LEU:HD21	1.91	0.52
1:A:328:TRP:HA	1:A:394:PRO:HB3	1.92	0.51
1:A:679:THR:HG22	1:A:680:VAL:HG13	1.93	0.50
1:A:288:MET:HE2	1:A:292:SER:HB3	1.94	0.50
1:A:298:PRO:HG2	1:A:697:MET:HE2	1.94	0.50
1:A:221:ILE:HG23	1:A:287:LEU:HD11	1.94	0.49
1:A:269:GLN:HA	1:A:274:ARG:HH21	1.77	0.49
1:A:294:TYR:HA	1:A:297:LEU:HD12	1.94	0.49
1:A:676:HIS:CG	1:A:843:ASP:HB2	2.48	0.49
1:A:408:SER:HB3	1:A:422:LEU:HD11	1.94	0.49
1:A:427:ILE:HD11	1:A:443:LEU:HD22	1.94	0.48
1:A:341:ALA:HB3	1:A:381:PRO:HB2	1.96	0.48
1:A:336:ILE:HD13	1:A:402:LEU:HD22	1.96	0.47
1:A:705:GLN:O	1:A:709:MET:HG2	2.15	0.46
1:A:117:ILE:HG12	1:A:692:CYS:HB3	1.98	0.46
1:A:916:ARG:HB3	1:A:921:ILE:HD11	1.98	0.45
1:A:334:LEU:HD13	1:A:396:LEU:HD11	2.00	0.44
1:A:592:PRO:HB2	1:A:622:TYR:HE2	1.83	0.44
1:A:392:TYR:CG	1:A:394:PRO:HD2	2.53	0.43
1:A:229:THR:HG23	1:A:232:MET:HE2	2.00	0.43
1:A:881:LEU:HD21	1:A:927:GLY:HA2	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:916:ARG:HD2	1:A:931:HIS:ND1	2.34	0.42
1:A:697:MET:SD	1:A:700:LYS:HD2	2.58	0.42
1:A:360:ILE:HG22	1:A:367:LEU:HD12	2.02	0.42
1:A:145:ASN:ND2	1:A:651:ARG:HH12	2.03	0.41
1:A:540:LEU:HB2	1:A:1023:ARG:HD2	2.02	0.41
1:A:985:TYR:HB3	1:A:1036:LEU:HD21	2.01	0.41
1:A:759:HIS:HB3	1:A:797:ASN:HD21	1.85	0.41
1:A:640:LYS:HG2	1:A:680:VAL:HG11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	887/1080 (82%)	857 (97%)	30 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	825/983 (84%)	798 (97%)	27 (3%)	33	60

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	VAL
1	A	145	ASN
1	A	198	VAL
1	A	251	ILE
1	A	329	VAL
1	A	335	ARG
1	A	338	ILE
1	A	353	LYS
1	A	376	VAL
1	A	409	VAL
1	A	474	GLU
1	A	475	LEU
1	A	545	GLU
1	A	561	ILE
1	A	573	LYS
1	A	578	ASP
1	A	679	THR
1	A	790	SER
1	A	822	ASN
1	A	834	LEU
1	A	845	VAL
1	A	913	ILE
1	A	937	PHE
1	A	942	LYS
1	A	971	CYS
1	A	1029	ASP
1	A	1036	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	145	ASN
1	A	160	HIS
1	A	189	GLN
1	A	370	ASN
1	A	384	ASN
1	A	467	ASN
1	A	556	HIS
1	A	597	GLN
1	A	661	GLN
1	A	676	HIS
1	A	682	GLN
1	A	731	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	795	GLN
1	A	797	ASN
1	A	918	ASN
1	A	940	HIS
1	A	993	GLN
1	A	994	HIS
1	A	1014	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	X3R	A	1101	-	35,36,36	1.41	6 (17%)	40,55,55	2.46	8 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	X3R	A	1101	-	-	7/19/47/47	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	X3R	C20-N19	3.65	1.39	1.36
2	A	1101	X3R	F32-C31	2.92	1.40	1.33
2	A	1101	X3R	F33-C31	2.59	1.39	1.33
2	A	1101	X3R	O16-C4	-2.42	1.34	1.38
2	A	1101	X3R	F34-C31	2.20	1.38	1.33
2	A	1101	X3R	C29-N24	-2.09	1.36	1.39

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	X3R	C23-C22-N21	-9.62	106.46	112.77
2	A	1101	X3R	O28-C29-N24	5.67	113.44	108.78
2	A	1101	X3R	C22-C23-N19	5.06	108.43	104.39
2	A	1101	X3R	C22-N24-C29	4.39	128.30	122.95
2	A	1101	X3R	C17-O16-C4	4.16	123.04	116.13
2	A	1101	X3R	N19-C20-N21	-3.88	106.71	111.67
2	A	1101	X3R	O30-C29-N24	-3.45	124.75	128.62
2	A	1101	X3R	C25-N24-C29	-2.79	107.71	110.46

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1101	X3R	C27-C25-C31-F32
2	A	1101	X3R	C27-C25-C31-F33
2	A	1101	X3R	C27-C25-C31-F34
2	A	1101	X3R	C11-C10-C8-C13
2	A	1101	X3R	O14-C13-C8-N7
2	A	1101	X3R	N15-C13-C8-N7
2	A	1101	X3R	N21-C22-N24-C25

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	898/1080 (83%)	0.14	49 (5%)	30 26	24, 59, 95, 137	40 (4%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	246	TYR	7.8
1	A	348	ILE	7.4
1	A	247	GLN	6.1
1	A	524	ARG	5.8
1	A	309	ARG	5.7
1	A	975	ARG	5.4
1	A	230	ARG	5.4
1	A	450	HIS	5.2
1	A	176	GLU	5.2
1	A	970	GLU	5.2
1	A	522	GLU	5.1
1	A	521	ASN	4.0
1	A	226	ARG	3.8
1	A	888	GLU	3.8
1	A	249	LYS	3.7
1	A	201	ASN	3.6
1	A	733	LYS	3.6
1	A	542	GLU	3.6
1	A	374	GLN	3.6
1	A	331	ASN	3.4
1	A	378	CYS	3.4
1	A	525	GLU	3.4
1	A	132	LYS	3.3
1	A	203	ASP	3.3
1	A	179	LYS	3.3
1	A	109	GLU	3.2
1	A	418	GLU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	440	LYS	3.1
1	A	111	LYS	2.9
1	A	532	LYS	2.9
1	A	231	SER	2.9
1	A	301	CYS	2.6
1	A	588	LYS	2.6
1	A	175	PRO	2.6
1	A	278	MET	2.5
1	A	502	ARG	2.5
1	A	725	ASP	2.5
1	A	187	LYS	2.5
1	A	199	SER	2.4
1	A	300	ASP	2.4
1	A	641	TYR	2.3
1	A	377	PRO	2.3
1	A	863	LYS	2.2
1	A	384	ASN	2.2
1	A	1022	ILE	2.2
1	A	604	CYS	2.2
1	A	479	TRP	2.1
1	A	1024	LYS	2.0
1	A	861	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	X3R	A	1101	32/32	0.94	0.07	37,48,53,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.