



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:15 PM UTC

PDB ID : 5EYP / pdb_00005eyp
Title : TUBULIN-DARPIN COMPLEX
Authors : Ahmad, S.; Knossow, M.; Gigant, B.
Deposited on : 2015-11-25
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	
2	B	445	
3	F	169	

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 8501 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	428	Total	C	N	O	S	0	1	0
			3328	2109	566	631	22			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	SER	GLY	variant	UNP D0VWZ0
A	340	SER	THR	variant	UNP D0VWZ0

- Molecule 2 is a protein called Tubulin beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	425	Total	C	N	O	S	0	12	0
			3405	2140	581	657	27			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	203	CYS	SER	variant	UNP D0VWY9
B	318	ILE	VAL	variant	UNP D0VWY9

- Molecule 3 is a protein called DESIGNED ANKYRIN REPEAT PROTEIN (DARPIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	130	Total	C	N	O	S	0	1	0
			960	603	171	183	3			

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 6 3 3	0	0
7	A	1	Total C O 6 3 3	0	0
7	A	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0
7	F	1	Total C O 6 3 3	0	0
7	F	1	Total C O 6 3 3	0	0
7	F	1	Total C O 6 3 3	0	0

- Molecule 8 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	281	Total 281	O 281	0	0
10	B	281	Total 281	O 281	0	0
10	F	93	Total 93	O 93	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha chain



- Molecule 2: Tubulin beta chain



- Molecule 3: DESIGNED ANKYRIN REPEAT PROTEIN (DARPIN)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.57Å 71.76Å 93.05Å 90.00° 99.85° 90.00°	Depositor
Resolution (Å)	49.74 – 1.90 49.74 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.74-1.90) 100.0 (49.74-1.90)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.90Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.173 , 0.208 0.174 , 0.207	Depositor DCC
R_{free} test set	4531 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8501	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, GDP, GTP, SO4, MG, LOC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.85	1/3404 (0.0%)	1.27	15/4623 (0.3%)
2	B	0.88	0/3509	1.22	8/4752 (0.2%)
3	F	0.87	0/976	1.38	2/1327 (0.2%)
All	All	0.87	1/7889 (0.0%)	1.26	25/10702 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	413	MET	SD-CE	-5.11	1.66	1.79

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	44	ASP	CA-CB-CG	7.37	119.97	112.60
1	A	205	ASP	CA-CB-CG	6.93	119.53	112.60
1	A	98	ASP	CA-CB-CG	6.39	119.00	112.60
2	B	220	THR	N-CA-C	-6.33	105.72	113.50
2	B	178	SER	N-CA-C	6.32	119.24	109.07
2	B	75	MET	CA-C-N	6.17	128.46	120.44
2	B	75	MET	C-N-CA	6.17	128.46	120.44
3	F	77	ASP	CA-CB-CG	6.10	118.70	112.60
1	A	109	THR	N-CA-C	6.07	118.39	111.11
2	B	97	SER	N-CA-C	6.02	120.69	113.23
2	B	250	ALA	N-CA-C	6.01	118.42	108.99
2	B	197	ASN	N-CA-C	5.97	121.30	113.30
1	A	306	ASP	CA-CB-CG	5.86	118.46	112.60
1	A	73	THR	N-CA-C	5.84	118.12	111.11
1	A	348	PRO	N-CA-C	5.71	124.22	112.47
1	A	181	VAL	N-CA-C	5.38	116.76	110.62
1	A	147	SER	CA-C-N	5.32	125.74	119.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	147	SER	C-N-CA	5.32	125.74	119.94
2	B	115	VAL	N-CA-C	5.23	116.58	110.62
1	A	162	GLY	N-CA-C	5.19	119.16	112.83
1	A	357	TYR	N-CA-C	5.18	119.22	113.01
1	A	81	GLY	CA-C-N	5.12	127.14	120.28
1	A	81	GLY	C-N-CA	5.12	127.14	120.28
1	A	353	VAL	N-CA-C	5.12	115.28	108.11
1	A	263	PRO	N-CA-C	5.03	121.25	113.75

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3328	0	3240	22	0
2	B	3405	0	3296	28	0
3	F	960	0	979	4	0
4	A	32	0	12	0	0
5	A	1	0	0	0	0
6	A	10	0	0	0	0
6	B	5	0	0	0	0
7	A	18	0	24	1	0
7	B	12	0	16	0	0
7	F	18	0	24	1	0
8	B	28	0	12	0	0
9	B	29	0	25	1	0
10	A	281	0	0	3	0
10	B	281	0	0	4	0
10	F	93	0	0	0	0
All	All	8501	0	7628	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:ASN:HD22	2:B:258:ASN:HD21	1.47	0.63
2:B:318:ILE:HD11	9:B:502:LOC:H4	1.84	0.60
2:B:57:THR:HG22	10:B:764:HOH:O	2.00	0.59
1:A:270:ALA:HB3	1:A:302[A]:MET:HE2	1.84	0.58
2:B:106:GLY:O	2:B:111:GLY:HA3	2.03	0.58
2:B:345:GLU:OE1	10:B:601:HOH:O	2.17	0.56
2:B:147[B]:SER:HG	2:B:190:SER:HG	1.53	0.56
2:B:400:ARG:HD2	3:F:112:TRP:NE1	2.22	0.54
2:B:295:MET:HG2	2:B:377:PHE:HB2	1.89	0.54
3:F:79:MET:HE2	3:F:79:MET:HA	1.91	0.53
1:A:221:ARG:NE	2:B:325:MET:HB3	2.24	0.52
2:B:147[B]:SER:OG	2:B:190:SER:OG	2.27	0.51
1:A:406:HIS:CG	2:B:263:PRO:HD3	2.47	0.50
2:B:214:PHE:O	2:B:218:LYS:HA	2.13	0.49
1:A:28:HIS:CE1	1:A:49:PHE:HB3	2.48	0.49
3:F:92:HIS:HE1	7:F:203:GOL:H32	1.77	0.48
1:A:266:HIS:HE1	10:A:675:HOH:O	1.96	0.48
1:A:413:MET:HE3	1:A:417:GLU:HB3	1.96	0.48
2:B:75:MET:HG3	2:B:94:PHE:CD1	2.49	0.47
2:B:192:HIS:CD2	2:B:424[A]:ASN:HD22	2.32	0.47
1:A:307:PRO:HA	1:A:383:ALA:HB2	1.96	0.46
2:B:83:PHE:O	2:B:86:ILE:HG22	2.14	0.46
2:B:269:MET:HG3	2:B:303:ALA:HB3	1.96	0.46
2:B:294:GLN:HE21	2:B:300:ASN:HD21	1.63	0.45
2:B:396:THR:O	2:B:400:ARG:HG3	2.15	0.45
1:A:8:HIS:HE1	10:A:836:HOH:O	1.99	0.45
2:B:172:MET:HG3	2:B:387:LEU:HD21	1.99	0.45
1:A:210:TYR:CZ	1:A:214:ARG:HD2	2.52	0.45
2:B:181:VAL:HG21	3:F:90:ILE:HG23	1.99	0.45
2:B:395:PHE:CE1	2:B:422[B]:GLU:HB2	2.52	0.45
1:A:175:PRO:HA	1:A:179:THR:HG22	1.98	0.45
1:A:320:ARG:HA	1:A:356:ASN:O	2.17	0.44
1:A:406:HIS:HB3	7:A:506:GOL:H32	1.98	0.44
2:B:66:ILE:HG12	2:B:121:VAL:HG12	2.00	0.44
2:B:275:LEU:HD12	2:B:275:LEU:HA	1.87	0.43
2:B:400:ARG:HD3	10:B:844:HOH:O	2.18	0.43
1:A:167:LEU:HG	1:A:200:CYS:HB3	2.02	0.42
1:A:406:HIS:HD2	10:A:849:HOH:O	2.02	0.42
2:B:215[B]:ARG:HH22	2:B:275:LEU:HD23	1.83	0.42
1:A:238:ILE:HG12	1:A:378:LEU:HD21	2.02	0.42
2:B:247:GLN:NE2	10:B:603:HOH:O	2.47	0.42
1:A:307:PRO:HA	1:A:383:ALA:CB	2.50	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:GLN:HE21	1:A:37:PRO:HD3	1.84	0.42
2:B:308:ARG:HG3	2:B:342:TYR:CZ	2.55	0.42
1:A:263:PRO:O	1:A:266:HIS:HD2	2.04	0.41
1:A:270:ALA:CB	1:A:302[A]:MET:HE2	2.50	0.41
1:A:192:HIS:CG	1:A:421:ALA:HA	2.56	0.40
1:A:71:GLU:HB3	1:A:98:ASP:HB3	2.03	0.40
2:B:298:SER:H	2:B:308:ARG:HH22	1.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	425/451 (94%)	408 (96%)	17 (4%)	0	100	100
2	B	433/445 (97%)	422 (98%)	10 (2%)	1 (0%)	43	36
3	F	129/169 (76%)	128 (99%)	1 (1%)	0	100	100
All	All	987/1065 (93%)	958 (97%)	28 (3%)	1 (0%)	48	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	276	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/379 (94%)	346 (97%)	10 (3%)	38	32
2	B	377/383 (98%)	361 (96%)	16 (4%)	26	19
3	F	99/131 (76%)	95 (96%)	4 (4%)	28	20
All	All	832/893 (93%)	802 (96%)	30 (4%)	32	23

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	84	ARG
1	A	181	VAL
1	A	188	ILE
1	A	280	LYS
1	A	297	GLU
1	A	302[A]	MET
1	A	302[B]	MET
1	A	335	ILE
1	A	349	THR
2	B	48	ARG
2	B	57	THR
2	B	75	MET
2	B	133	GLN
2	B	158	ARG
2	B	221	THR
2	B	247	GLN
2	B	275	LEU
2	B	276	THR
2	B	278	ARG
2	B	295	MET
2	B	325	MET
2	B	338	LYS
2	B	345	GLU
2	B	371	LEU
2	B	405	LEU
3	F	20	GLU
3	F	65	VAL
3	F	79	MET
3	F	127	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	31	GLN
1	A	101	ASN
1	A	107	HIS
1	A	266	HIS
1	A	283	HIS
1	A	301	GLN
1	A	372	GLN
1	A	393	HIS
2	B	8	GLN
2	B	14	ASN
2	B	91	ASN
2	B	136	GLN
2	B	167	ASN
2	B	247	GLN
2	B	300	ASN
2	B	331	GLN
2	B	336	GLN
3	F	36	ASN
3	F	92	HIS
3	F	142	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 1 is monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SO4	B	503	-	4,4,4	0.26	0	6,6,6	0.11	0
7	GOL	A	505	-	5,5,5	0.09	0	5,5,5	0.18	0
7	GOL	F	203	-	5,5,5	0.07	0	5,5,5	0.13	0
7	GOL	A	506	-	5,5,5	0.13	0	5,5,5	0.30	0
9	LOC	B	502	-	31,31,31	0.64	1 (3%)	44,44,44	0.73	2 (4%)
6	SO4	A	503	-	4,4,4	0.24	0	6,6,6	0.09	0
7	GOL	F	201	-	5,5,5	0.06	0	5,5,5	0.29	0
4	GTP	A	501	5	33,34,34	2.17	2 (6%)	50,54,54	0.53	0
7	GOL	F	202	-	5,5,5	0.08	0	5,5,5	0.13	0
8	GDP	B	501	-	29,30,30	0.47	0	45,47,47	0.47	0
7	GOL	B	505	-	5,5,5	0.09	0	5,5,5	0.60	0
6	SO4	A	504	-	4,4,4	0.31	0	6,6,6	0.09	0
7	GOL	B	504	-	5,5,5	0.08	0	5,5,5	0.30	0
7	GOL	A	507	-	5,5,5	0.06	0	5,5,5	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	505	-	-	0/4/4/4	-
7	GOL	F	203	-	-	0/4/4/4	-
7	GOL	A	506	-	-	0/4/4/4	-
9	LOC	B	502	-	-	0/12/25/25	0/3/3/3
7	GOL	F	201	-	-	0/4/4/4	-
4	GTP	A	501	5	-	7/22/38/38	0/3/3/3
7	GOL	F	202	-	-	2/4/4/4	-
8	GDP	B	501	-	-	5/16/32/32	0/3/3/3
7	GOL	B	505	-	-	0/4/4/4	-
7	GOL	B	504	-	-	0/4/4/4	-
7	GOL	A	507	-	-	1/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	GTP	PB-O3B	-8.77	1.50	1.59
4	A	501	GTP	PA-O3A	-8.22	1.50	1.59
9	B	502	LOC	C11-C14	2.66	1.57	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	B	502	LOC	C14-C11-N1	2.42	116.22	114.31
9	B	502	LOC	C11-C14-C15	-2.35	114.98	117.06

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	501	GTP	PB-O3B-PG-O2G
4	A	501	GTP	C5'-O5'-PA-O3A
4	A	501	GTP	C5'-O5'-PA-O1A
8	B	501	GDP	C5'-O5'-PA-O3A
8	B	501	GDP	C5'-O5'-PA-O1A
8	B	501	GDP	C5'-O5'-PA-O2A
4	A	501	GTP	C5'-O5'-PA-O2A
4	A	501	GTP	PB-O3A-PA-O2A
7	A	507	GOL	C1-C2-C3-O3
7	F	202	GOL	O1-C1-C2-C3
7	F	202	GOL	O1-C1-C2-O2
8	B	501	GDP	PB-O3A-PA-O1A
4	A	501	GTP	PB-O3B-PG-O3G
8	B	501	GDP	PB-O3A-PA-O2A
4	A	501	GTP	PB-O3A-PA-O1A


There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	F	203	GOL	1	0
7	A	506	GOL	1	0
9	B	502	LOC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/451 (94%)	0.22	18 (4%) 40 43	25, 49, 86, 106	1 (0%)
2	B	425/445 (95%)	0.13	19 (4%) 38 40	17, 40, 67, 120	12 (2%)
3	F	130/169 (76%)	0.23	5 (3%) 44 47	31, 45, 83, 123	1 (0%)
All	All	983/1065 (92%)	0.18	42 (4%) 40 42	17, 44, 81, 123	14 (1%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	437	VAL	5.2
2	B	276	THR	4.6
1	A	346	TRP	4.5
2	B	277	SER	3.6
2	B	94	PHE	3.4
2	B	278	ARG	3.2
2	B	286	LEU	3.1
3	F	141	ALA	3.1
2	B	75	MET	3.0
2	B	57	THR	3.0
2	B	247	GLN	2.9
2	B	441	ASP	2.9
2	B	74	THR	2.8
1	A	1	MET	2.8
1	A	335	ILE	2.7
1	A	309	HIS	2.5
1	A	37	PRO	2.5
2	B	285	ALA	2.5
2	B	250	ALA	2.5
1	A	341	ILE	2.4
2	B	370	GLY	2.4
1	A	282	TYR	2.4
2	B	325	MET	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	281	ALA	2.4
3	F	139	VAL	2.4
1	A	348	PRO	2.3
2	B	248	LEU	2.3
1	A	338	LYS	2.3
1	A	435	VAL	2.3
1	A	283	HIS	2.2
3	F	14	LEU	2.2
1	A	436	GLY	2.2
3	F	142	GLN	2.2
2	B	439	THR	2.2
2	B	275	LEU	2.2
2	B	440	ALA	2.2
3	F	36	ASN	2.1
1	A	349	THR	2.1
1	A	246	GLY	2.1
1	A	316	CYS	2.0
1	A	344	VAL	2.0
2	B	177	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	SO4	B	503	5/5	0.75	0.16	121,122,122,122	0
7	GOL	A	506	6/6	0.75	0.19	65,69,70,71	0
7	GOL	F	203	6/6	0.79	0.17	83,84,84,85	0
6	SO4	A	504	5/5	0.80	0.18	116,117,117,118	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	GOL	B	505	6/6	0.82	0.17	69,78,80,82	0
7	GOL	A	507	6/6	0.82	0.16	76,80,81,82	0
7	GOL	B	504	6/6	0.83	0.16	65,68,69,69	0
7	GOL	A	505	6/6	0.85	0.14	49,53,56,60	0
7	GOL	F	202	6/6	0.86	0.13	50,59,62,64	0
6	SO4	A	503	5/5	0.92	0.07	79,80,81,81	0
5	MG	A	502	1/1	0.94	0.08	68,68,68,68	0
7	GOL	F	201	6/6	0.94	0.09	45,46,48,51	0
4	GTP	A	501	32/32	0.98	0.05	31,35,41,43	0
9	LOC	B	502	29/29	0.98	0.05	28,33,38,39	0
8	GDP	B	501	28/28	0.99	0.04	27,31,32,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around LOC B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.