



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:59 PM UTC

PDB ID : 6EYN / pdb_00006eyn
Title : Structure of the 8D6 (anti-IgE) Fab
Authors : Chen, J.B.; Ramadani, F.; Pang, M.O.Y.; Beavil, R.L.; Holdom, M.D.;
Mitropoulou, A.N.; Beavil, A.J.; Gould, H.J.; Chang, T.W.; Sutton, B.J.;
McDonnell, J.M.; Davies, A.M.
Deposited on : 2017-11-13
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	218	8% 94% • •
1	C	218	15% 88% 7% 5%
1	L	218	% 94% 5%
2	B	227	9% 84% 7% 9%
2	D	227	13% 81% • 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	227	 A horizontal bar chart showing the quality of chain H. The bar is divided into three segments: a small red segment at the beginning labeled '4%', a large green segment in the middle labeled '90%', and a small yellow segment at the end labeled '7%'. The total length of the bar represents 100%.

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 8D6 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	217	Total	C	N	O	S	0	1	0
			1629	1026	269	329	5			
1	A	214	Total	C	N	O	S	0	1	0
			1562	987	258	312	5			
1	C	207	Total	C	N	O	S	0	1	0
			1477	924	247	301	5			

- Molecule 2 is a protein called 8D6 Fab heavy chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	211	Total	C	N	O	S	0	4	0
			1576	1002	258	308	8			
2	B	207	Total	C	N	O	S	0	1	0
			1507	967	249	284	7			
2	D	194	Total	C	N	O	S	0	1	0
			1401	888	236	270	7			

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	L	1	Total	C	O	0	0
			6	4	2		
3	L	1	Total	C	O	0	0
			6	4	2		
3	L	1	Total	C	O	0	0
			6	4	2		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total C O 4 2 2	0	0
4	H	1	Total C O 4 2 2	0	0
4	H	1	Total C O 4 2 2	0	0
4	H	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total O S 5 4 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	79	Total 79	O 79	0	0
6	H	51	Total 51	O 51	0	0
6	A	39	Total 39	O 39	0	0
6	B	21	Total 21	O 21	0	0
6	C	36	Total 36	O 36	0	0
6	D	23	Total 23	O 23	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: 8D6 Fab light chain





- Molecule 1: 8D6 Fab light chain




- Molecule 1: 8D6 Fab light chain




- Molecule 2: 8D6 Fab heavy chain






- Molecule 2: 8D6 Fab heavy chain





● Molecule 2: 8D6 Fab heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	157.96Å 157.96Å 77.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.55 – 2.40 43.55 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.7 (43.55-2.40) 98.7 (43.55-2.40)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 2.40Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.181 , 0.212 0.183 , 0.212	Depositor DCC
R_{free} test set	3695 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	51.5	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9460	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SO4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.15	0/1601	0.36	0/2186
1	C	0.16	0/1515	0.37	0/2066
1	L	0.20	0/1670	0.40	0/2280
2	B	0.13	0/1551	0.33	0/2125
2	D	0.14	0/1444	0.33	0/1973
2	H	0.18	0/1630	0.37	0/2229
All	All	0.16	0/9411	0.36	0/12859

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1562	0	1415	5	0
1	C	1477	0	1276	6	0
1	L	1629	0	1507	7	0
2	B	1507	0	1370	8	0
2	D	1401	0	1193	5	0
2	H	1576	0	1474	0	0
3	L	18	0	21	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	8	0	12	1	0
4	C	8	0	12	0	0
4	H	20	0	30	1	0
5	D	5	0	0	0	0
6	A	39	0	0	1	0
6	B	21	0	0	0	0
6	C	36	0	0	0	0
6	D	23	0	0	0	0
6	H	51	0	0	1	0
6	L	79	0	0	0	0
All	All	9460	0	8310	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:18:VAL:HG22	2:B:86:LEU:HD11	1.64	0.78
2:D:18:VAL:HG22	2:D:86:LEU:HD11	1.71	0.71
1:L:65[B]:ARG:NH2	1:L:86:ASP:OD1	2.35	0.59
2:D:4:LEU:HD22	2:D:34:MET:HE1	1.88	0.55
1:C:14:SER:HA	1:C:111:LYS:HB2	1.88	0.55
1:A:65[B]:ARG:NH1	1:A:86:ASP:OD2	2.39	0.54
1:L:120:PHE:HD2	4:H:302:EDO:H11	1.74	0.52
1:A:65[B]:ARG:HH12	1:A:86:ASP:CG	2.20	0.48
6:H:420:HOH:O	2:B:13:LYS:HE3	2.12	0.48
1:A:203:GLN:NE2	6:A:303:HOH:O	2.48	0.47
1:L:84:GLU:HG2	3:L:301:PEG:H32	1.96	0.47
2:B:11:LEU:HB2	2:B:153:PRO:HG3	1.95	0.47
2:D:177:GLN:HG3	2:D:181:LEU:O	2.17	0.45
1:L:41:HIS:HB3	1:L:51:LEU:HD11	1.99	0.44
2:D:152:PHE:HA	2:D:153:PRO:HA	1.82	0.44
2:B:203:ASN:ND2	2:B:214:ASP:OD1	2.50	0.43
1:C:80:HIS:HA	1:C:81:PRO:HA	1.83	0.43
2:B:38:LYS:HB2	2:B:48:ILE:HD11	2.01	0.43
2:D:22:CYS:SG	2:D:96[B]:CYS:HB3	2.58	0.43
1:A:147:GLU:H	1:A:147:GLU:CD	2.27	0.43
1:C:16:GLY:HA2	1:C:81:PRO:HB2	2.01	0.43
1:L:13:VAL:HG21	1:L:19:ALA:HB2	2.00	0.42
1:L:147:GLU:H	1:L:147:GLU:CD	2.27	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:22:CYS:SG	2:B:96[A]:CYS:HB3	2.58	0.42
1:C:13:VAL:HG21	1:C:19:ALA:HB2	2.00	0.42
1:A:41:HIS:HB3	1:A:51:LEU:HD11	2.01	0.42
2:B:20:LEU:HD22	2:B:113:THR:HG21	2.02	0.41
1:L:129:LEU:HD12	1:L:129:LEU:HA	1.89	0.41
1:C:119:VAL:HA	1:C:139:LEU:O	2.20	0.41
2:B:33:TRP:HE1	4:B:302:EDO:H21	1.86	0.41
1:C:37:MET:HE2	1:C:37:MET:HB2	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/218 (97%)	202 (96%)	9 (4%)	0	100	100
1	C	204/218 (94%)	196 (96%)	8 (4%)	0	100	100
1	L	216/218 (99%)	210 (97%)	6 (3%)	0	100	100
2	B	202/227 (89%)	197 (98%)	5 (2%)	0	100	100
2	D	189/227 (83%)	180 (95%)	9 (5%)	0	100	100
2	H	209/227 (92%)	204 (98%)	4 (2%)	1 (0%)	24	37
All	All	1231/1335 (92%)	1189 (97%)	41 (3%)	1 (0%)	48	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	132	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/188 (83%)	153 (98%)	3 (2%)	50	71
1	C	142/188 (76%)	137 (96%)	5 (4%)	32	53
1	L	170/188 (90%)	168 (99%)	2 (1%)	63	81
2	B	144/192 (75%)	140 (97%)	4 (3%)	38	60
2	D	125/192 (65%)	123 (98%)	2 (2%)	55	76
2	H	166/192 (86%)	161 (97%)	5 (3%)	36	58
All	All	903/1140 (79%)	882 (98%)	21 (2%)	44	66

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	41	HIS
1	L	169	GLU
2	H	4	LEU
2	H	18	VAL
2	H	71	THR
2	H	102	ARG
2	H	144	LEU
1	A	41	HIS
1	A	82	VAL
1	A	195	VAL
2	B	4	LEU
2	B	18	VAL
2	B	111	GLN
2	B	156	VAL
1	C	32	ASP
1	C	41	HIS
1	C	82	VAL
1	C	164	GLN
1	C	167	VAL
2	D	18	VAL
2	D	204	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	42	GLN
1	L	57	ASN
1	L	80	HIS
2	H	39	GLN
2	H	170	HIS
1	A	80	HIS
1	C	42	GLN
2	D	5	GLN
2	D	39	GLN
2	D	170	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	C	302	-	3,3,3	0.43	0	2,2,2	0.35	0
4	EDO	H	302	-	3,3,3	0.43	0	2,2,2	0.29	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	D	301	-	4,4,4	0.24	0	6,6,6	0.10	0
3	PEG	L	302	-	5,5,6	0.87	0	4,4,5	0.26	0
4	EDO	H	301	-	3,3,3	0.51	0	2,2,2	0.23	0
4	EDO	H	304	-	3,3,3	0.42	0	2,2,2	0.36	0
4	EDO	C	301	-	3,3,3	0.45	0	2,2,2	0.33	0
4	EDO	H	305	-	3,3,3	0.44	0	2,2,2	0.35	0
3	PEG	L	301	-	5,5,6	0.88	0	4,4,5	0.27	0
4	EDO	B	301	-	3,3,3	0.41	0	2,2,2	0.43	0
4	EDO	B	302	-	3,3,3	0.42	0	2,2,2	0.38	0
4	EDO	H	303	-	3,3,3	0.39	0	2,2,2	0.46	0
3	PEG	L	303	-	5,5,6	0.90	0	4,4,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	302	-	-	0/1/1/1	-
4	EDO	H	302	-	-	1/1/1/1	-
3	PEG	L	302	-	-	2/3/3/4	-
4	EDO	H	301	-	-	0/1/1/1	-
4	EDO	H	304	-	-	0/1/1/1	-
4	EDO	C	301	-	-	0/1/1/1	-
4	EDO	H	305	-	-	0/1/1/1	-
3	PEG	L	301	-	-	1/3/3/4	-
4	EDO	B	301	-	-	0/1/1/1	-
4	EDO	B	302	-	-	0/1/1/1	-
4	EDO	H	303	-	-	0/1/1/1	-
3	PEG	L	303	-	-	3/3/3/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	L	301	PEG	O2-C3-C4-O4
3	L	303	PEG	C1-C2-O2-C3
3	L	302	PEG	O2-C3-C4-O4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	H	302	EDO	O1-C1-C2-O2
3	L	302	PEG	C4-C3-O2-C2
3	L	303	PEG	O2-C3-C4-O4
3	L	303	PEG	C4-C3-O2-C2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	302	EDO	1	0
3	L	301	PEG	1	0
4	B	302	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/218 (98%)	0.40	18 (8%) 17 14	31, 68, 115, 129	1 (0%)
1	C	207/218 (94%)	0.67	33 (15%) 5 4	30, 73, 147, 173	1 (0%)
1	L	217/218 (99%)	-0.17	2 (0%) 81 78	27, 49, 81, 106	1 (0%)
2	B	207/227 (91%)	0.54	21 (10%) 12 9	44, 73, 119, 144	1 (0%)
2	D	194/227 (85%)	0.75	30 (15%) 5 4	37, 82, 127, 147	1 (0%)
2	H	211/227 (92%)	0.05	9 (4%) 40 36	25, 53, 101, 115	4 (1%)
All	All	1250/1335 (93%)	0.36	113 (9%) 15 12	25, 64, 123, 173	9 (0%)

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	211	THR	6.0
2	B	198	GLN	5.8
1	A	31	TYR	5.6
2	H	2	VAL	5.2
2	B	197	THR	5.1
1	C	197	ALA	4.9
2	B	29	PHE	4.7
2	B	193	SER	4.2
1	A	126	ASP	4.1
1	A	34	ASP	4.1
2	D	165	LEU	3.9
1	A	131	SER	3.9
1	C	31	TYR	3.8
1	C	133	THR	3.7
1	C	212	SER	3.7
1	C	191	GLU	3.7
1	C	186	SER	3.6
1	C	158	LEU	3.6
1	C	125	SER	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	43	GLN	3.5
1	C	207	SER	3.5
2	D	210	ASN	3.5
1	L	206	SER	3.5
1	C	188	ALA	3.5
2	D	127	VAL	3.3
2	D	163	GLY	3.3
1	A	30	ASP	3.2
1	C	195	VAL	3.2
1	C	136	VAL	3.2
2	B	2	VAL	3.1
1	C	154	VAL	3.1
2	D	184	LEU	3.1
1	C	182	THR	3.1
2	B	201	ILE	3.1
2	D	209	SER	3.0
2	D	145	GLY	3.0
2	D	128	PHE	2.9
1	C	183	LEU	2.9
1	L	32	ASP	2.9
2	B	41	PRO	2.8
1	C	160	SER	2.8
2	B	199	THR	2.8
2	D	121	SER	2.7
2	B	169	VAL	2.7
2	D	28	THR	2.7
2	D	141	THR	2.7
2	D	178	SER	2.7
1	A	188	ALA	2.7
2	B	141	THR	2.7
2	B	30	ASN	2.6
1	A	133	THR	2.6
2	B	194	SER	2.6
2	B	25	SER	2.6
1	A	191	GLU	2.5
1	A	214	ASN	2.5
2	D	168	GLY	2.5
1	A	1	ASP	2.5
2	H	221	SER	2.5
1	C	157	ALA	2.4
1	A	97	GLU	2.4
1	C	34	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	44	ASP	2.4
1	A	125	SER	2.4
1	A	72	GLY	2.4
1	C	152	TRP	2.4
2	D	126	SER	2.3
2	H	75	SER	2.3
1	C	121	ILE	2.3
2	D	177	GLN	2.3
1	C	124	PRO	2.3
2	B	131	ALA	2.3
2	B	42	GLY	2.3
2	B	196	GLY	2.3
2	D	44	ASP	2.3
1	C	153	LYS	2.3
2	D	43	GLN	2.3
1	C	126	ASP	2.3
2	H	29	PHE	2.3
2	D	190	VAL	2.2
1	C	45	GLY	2.2
2	H	196	GLY	2.2
1	A	74	ASP	2.2
2	D	129	PRO	2.2
2	D	157	THR	2.2
1	A	129	LEU	2.2
1	A	215	ARG	2.2
2	D	29	PHE	2.2
2	D	132	PRO	2.2
1	A	198	CYS	2.2
2	B	167	SER	2.2
1	C	123	PRO	2.2
2	D	26	GLY	2.1
2	B	162	SER	2.1
2	B	132	PRO	2.1
2	D	181	LEU	2.1
1	C	194	LYS	2.1
2	H	41	PRO	2.1
2	D	142	ALA	2.1
1	C	156	ASN	2.1
1	C	122	PHE	2.1
1	C	185	LEU	2.1
1	C	184	THR	2.1
1	A	207	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	203	ASN	2.0
2	D	166	THR	2.0
2	H	132	PRO	2.0
2	H	194	SER	2.0
1	C	193	HIS	2.0
1	C	46	GLN	2.0
2	B	43	GLN	2.0
1	C	127	GLU	2.0
2	D	161	ASN	2.0
2	D	207	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	D	301	5/5	0.66	0.18	143,144,145,147	0
3	PEG	L	302	6/7	0.77	0.33	81,92,101,103	0
3	PEG	L	303	6/7	0.78	0.17	59,78,82,83	0
3	PEG	L	301	6/7	0.78	0.21	71,80,85,90	0
4	EDO	C	301	4/4	0.82	0.18	85,89,92,95	0
4	EDO	H	305	4/4	0.86	0.14	89,90,91,91	0
4	EDO	B	301	4/4	0.88	0.17	79,81,85,87	0
4	EDO	H	301	4/4	0.91	0.15	53,55,58,60	0
4	EDO	C	302	4/4	0.91	0.10	70,71,77,81	0
4	EDO	H	302	4/4	0.91	0.22	66,68,71,79	0
4	EDO	H	303	4/4	0.92	0.10	67,70,71,81	0
4	EDO	H	304	4/4	0.92	0.11	88,88,88,92	0
4	EDO	B	302	4/4	0.92	0.13	67,71,72,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.