



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 12:16 AM UTC

PDB ID : 6EYV / pdb_00006eyv
Title : Crystal structure of the pyoverdine maturation protein PvdP in complex with the mock substrates L-tyrosine and zinc.
Authors : Poppe, J.; Blankenfeldt, W.
Deposited on : 2017-11-13
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	536	
1	B	536	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14338 atoms, of which 6951 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PvdP.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	457	Total	C	H	N	O	S	1	0	0
			7116	2344	3457	665	643	7			
1	A	459	Total	C	H	N	O	S	0	0	0
			7146	2352	3472	668	647	7			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	9	HIS	-	expression tag	UNP A0A0H2ZBG1
B	10	HIS	-	expression tag	UNP A0A0H2ZBG1
B	11	HIS	-	expression tag	UNP A0A0H2ZBG1
B	12	HIS	-	expression tag	UNP A0A0H2ZBG1
B	13	HIS	-	expression tag	UNP A0A0H2ZBG1
B	14	HIS	-	expression tag	UNP A0A0H2ZBG1
B	15	SER	-	expression tag	UNP A0A0H2ZBG1
B	16	SER	-	expression tag	UNP A0A0H2ZBG1
B	17	GLY	-	expression tag	UNP A0A0H2ZBG1
B	18	LEU	-	expression tag	UNP A0A0H2ZBG1
B	19	GLU	-	expression tag	UNP A0A0H2ZBG1
B	20	VAL	-	expression tag	UNP A0A0H2ZBG1
B	21	LEU	-	expression tag	UNP A0A0H2ZBG1
B	22	PHE	-	expression tag	UNP A0A0H2ZBG1
B	23	GLN	-	expression tag	UNP A0A0H2ZBG1
B	24	GLY	-	expression tag	UNP A0A0H2ZBG1
B	25	THR	-	expression tag	UNP A0A0H2ZBG1
A	9	HIS	-	expression tag	UNP A0A0H2ZBG1
A	10	HIS	-	expression tag	UNP A0A0H2ZBG1
A	11	HIS	-	expression tag	UNP A0A0H2ZBG1
A	12	HIS	-	expression tag	UNP A0A0H2ZBG1
A	13	HIS	-	expression tag	UNP A0A0H2ZBG1
A	14	HIS	-	expression tag	UNP A0A0H2ZBG1
A	15	SER	-	expression tag	UNP A0A0H2ZBG1
A	16	SER	-	expression tag	UNP A0A0H2ZBG1

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	GLY	-	expression tag	UNP A0A0H2ZBG1
A	18	LEU	-	expression tag	UNP A0A0H2ZBG1
A	19	GLU	-	expression tag	UNP A0A0H2ZBG1
A	20	VAL	-	expression tag	UNP A0A0H2ZBG1
A	21	LEU	-	expression tag	UNP A0A0H2ZBG1
A	22	PHE	-	expression tag	UNP A0A0H2ZBG1
A	23	GLN	-	expression tag	UNP A0A0H2ZBG1
A	24	GLY	-	expression tag	UNP A0A0H2ZBG1
A	25	THR	-	expression tag	UNP A0A0H2ZBG1

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is TYROSINE (CCD ID: TYR) (formula: C₉H₁₁NO₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	H	N	O	0	0
			24	9	11	1	3		
3	A	1	Total	C	H	N	O	0	0
			24	9	11	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	13	Total 13	O 13	0	0
4	A	11	Total 11	O 11	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PvdP



• Molecule 1: PvdP



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.33Å 109.14Å 82.51Å 90.00° 95.66° 90.00°	Depositor
Resolution (Å)	48.09 – 2.70 48.09 – 2.70	Depositor EDS
% Data completeness (in resolution range)	61.4 (48.09-2.70) 61.4 (48.09-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 2.69Å)	Xtriage
Refinement program	PHENIX (dev_2875)	Depositor
R, R_{free}	0.205 , 0.262 0.204 , 0.255	Depositor DCC
R_{free} test set	1991 reflections (5.32%)	wwPDB-VP
Wilson B-factor (Å ²)	39.3	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 29.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	14338	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.11	0/3797	0.26	0/5173
1	B	0.11	0/3782	0.26	0/5154
All	All	0.11	0/7579	0.26	0/10327

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3674	3472	3471	5	0
1	B	3659	3457	3457	7	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	13	11	8	0	0
3	B	13	11	8	0	0
4	A	11	0	0	0	0
4	B	13	0	0	0	0
All	All	7387	6951	6944	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:ASP:OD1	1:A:182:ARG:NH2	2.33	0.61
1:A:475:ASP:N	1:A:475:ASP:OD1	2.48	0.47
1:B:258:LYS:HD2	1:B:372:LEU:HD21	1.98	0.46
1:B:475:ASP:N	1:B:475:ASP:OD1	2.48	0.45
1:B:500:ASP:OD2	1:B:503:THR:N	2.49	0.43
1:A:258:LYS:HD2	1:A:372:LEU:HD21	1.99	0.43
1:B:106:ALA:O	1:B:301:ARG:NH2	2.47	0.42
1:A:371:GLU:HA	1:A:375:HIS:HB3	2.02	0.41
1:A:362:THR:OG1	1:A:472:GLU:OE2	2.30	0.41
1:B:223:ARG:NH1	1:B:371:GLU:OE1	2.54	0.41
1:B:204:LEU:HD23	1:B:204:LEU:C	2.46	0.40
1:B:371:GLU:HA	1:B:375:HIS:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	453/536 (84%)	436 (96%)	17 (4%)	0	100	100
1	B	451/536 (84%)	437 (97%)	14 (3%)	0	100	100
All	All	904/1072 (84%)	873 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/454 (82%)	369 (100%)	2 (0%)	81	92
1	B	368/454 (81%)	366 (100%)	2 (0%)	81	92
All	All	739/908 (81%)	735 (100%)	4 (0%)	76	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	129	ARG
1	B	182	ARG
1	A	129	ARG
1	A	182	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	37	GLN
1	B	189	HIS
1	A	37	GLN
1	A	189	HIS
1	A	455	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TYR	A	603	-	12,13,13	0.10	0	13,17,17	0.14	0
3	TYR	B	902	-	12,13,13	0.09	0	13,17,17	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TYR	A	603	-	-	4/8/8/8	0/1/1/1
3	TYR	B	902	-	-	6/8/8/8	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	902	TYR	N-CA-CB-CG
3	B	902	TYR	OXT-C-CA-N
3	B	902	TYR	O-C-CA-CB
3	B	902	TYR	O-C-CA-N
3	B	902	TYR	OXT-C-CA-CB
3	A	603	TYR	O-C-CA-CB
3	A	603	TYR	OXT-C-CA-CB
3	B	902	TYR	C-CA-CB-CG
3	A	603	TYR	N-CA-CB-CG
3	A	603	TYR	OXT-C-CA-N

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	459/536 (85%)	0.36	24 (5%) 33 29	16, 36, 62, 84	0
1	B	457/536 (85%)	0.28	19 (4%) 40 37	14, 35, 58, 85	2 (0%)
All	All	916/1072 (85%)	0.32	43 (4%) 36 33	14, 35, 60, 85	2 (0%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	388	ASP	6.4
1	A	388	ASP	5.5
1	A	404	PHE	4.0
1	A	175	LEU	3.9
1	A	484	GLY	3.6
1	B	396	MET	3.3
1	B	391	ASN	3.3
1	B	484	GLY	3.1
1	A	392	GLY	2.8
1	A	389	PRO	2.8
1	A	387	ARG	2.8
1	A	36	PRO	2.7
1	B	384	SER	2.7
1	A	159	GLY	2.6
1	B	36	PRO	2.6
1	B	175	LEU	2.6
1	A	390	SER	2.5
1	A	394	PRO	2.5
1	B	510	ILE	2.5
1	B	404	PHE	2.5
1	A	393	ALA	2.4
1	A	499	LEU	2.4
1	A	505	LYS	2.4
1	A	513	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	392	GLY	2.3
1	B	389	PRO	2.3
1	B	153	ALA	2.3
1	A	429	TRP	2.2
1	A	406	PRO	2.2
1	B	76	LEU	2.2
1	A	150	GLU	2.2
1	A	482	THR	2.2
1	A	211	GLN	2.2
1	A	395	VAL	2.2
1	B	206	TRP	2.1
1	A	206	TRP	2.1
1	B	387	ARG	2.1
1	B	211	GLN	2.1
1	B	324	ASP	2.1
1	A	511	ILE	2.1
1	B	503	THR	2.1
1	B	277	THR	2.0
1	A	174	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TYR	B	902	13/13	0.88	0.18	22,37,46,53	0
3	TYR	A	603	13/13	0.88	0.17	25,37,48,55	0
2	ZN	A	602	1/1	0.97	0.11	55,55,55,55	0
2	ZN	B	901	1/1	0.97	0.10	53,53,53,53	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	A	601	1/1	0.97	0.10	37,37,37,37	0
2	ZN	B	900	1/1	0.99	0.06	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.