



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 24, 2026 – 01:40 AM UTC

PDB ID : 8EZW / pdb_00008ezw
Title : Structure of Apo ZrgA deletion 124-184 from *Vibrio cholerae*
Authors : Valencia, D.W.; Yukl, E.T.
Deposited on : 2022-11-01
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	72% 11% 18%
1	B	160	2% 65% 15% • 18%
1	C	160	2% 64% 15% • 17%
1	D	160	• 71% 15% • 14%
1	E	160	• 72% 7% • 18%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	160	 A horizontal bar chart showing the quality of chain F. The bar is divided into four segments: a small red segment at the beginning labeled '4%', followed by a green segment labeled '54%', a yellow segment labeled '23%', and a grey segment at the end labeled '18%'. A small black dot is visible on the yellow segment.

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6763 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Zinc-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	132	Total	C	N	O	S	0	0	0
			1051	665	179	205	2			
1	B	132	Total	C	N	O	S	0	2	0
			1069	676	182	209	2			
1	C	133	Total	C	N	O	S	0	0	0
			1055	667	180	206	2			
1	D	138	Total	C	N	O	S	0	0	0
			1094	691	185	216	2			
1	E	131	Total	C	N	O	S	0	1	0
			1056	669	180	205	2			
1	F	131	Total	C	N	O	S	0	1	0
			1056	668	180	206	2			

- Molecule 2 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	E	1	Total	C	O	0	0
			10	6	4		
2	F	1	Total	C	O	0	0
			10	6	4		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	82	Total	O	0	0
			82	82		
3	B	69	Total	O	0	0
			69	69		
3	C	53	Total	O	0	0
			53	53		
3	D	64	Total	O	0	0
			64	64		
3	E	52	Total	O	0	0
			52	52		
3	F	42	Total	O	0	0
			42	42		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Zinc-binding protein

Chain A: 




- Molecule 1: Zinc-binding protein

Chain B: 




- Molecule 1: Zinc-binding protein

Chain C: 




- Molecule 1: Zinc-binding protein

Chain D: 



● Molecule 1: Zinc-binding protein



● Molecule 1: Zinc-binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.06Å 65.44Å 88.73Å 90.00° 98.77° 90.00°	Depositor
Resolution (Å)	45.88 – 2.00 45.88 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.1 (45.88-2.00) 99.1 (45.88-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 2.00Å)	Xtriage
Refinement program	REFMAC 7.1.018	Depositor
R, R_{free}	0.192 , 0.236 0.201 , 0.243	Depositor DCC
R_{free} test set	3083 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	30.9	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6763	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.21	3/1069 (0.3%)	1.37	0/1443
1	B	1.29	5/1087 (0.5%)	1.36	5/1466 (0.3%)
1	C	1.30	6/1073 (0.6%)	1.42	7/1448 (0.5%)
1	D	1.24	0/1113	1.36	4/1502 (0.3%)
1	E	1.27	3/1074 (0.3%)	1.31	1/1449 (0.1%)
1	F	1.20	3/1074 (0.3%)	1.34	1/1450 (0.1%)
All	All	1.25	20/6490 (0.3%)	1.36	18/8758 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	F	0	1
All	All	0	2

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	189	ALA	C-O	9.07	1.34	1.24
1	B	218	ILE	C-O	8.37	1.33	1.24
1	C	92	PHE	C-O	-6.81	1.16	1.23
1	B	188	THR	C-O	6.66	1.32	1.24
1	B	206	THR	C-O	6.58	1.31	1.23

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	65	PHE	CA-C-O	-6.54	114.45	121.45
1	C	69	PRO	CA-C-O	-6.40	114.14	121.56
1	D	90	LYS	N-CA-CB	-6.27	100.66	110.44
1	D	43	ASN	CA-CB-CG	-5.99	106.61	112.60
1	B	84	THR	CA-CB-OG1	-5.94	100.69	109.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	226	LYS	Peptide
1	F	97	LYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1051	0	1034	13	0
1	B	1069	0	1051	15	0
1	C	1055	0	1037	24	0
1	D	1094	0	1066	15	0
1	E	1056	0	1043	8	0
1	F	1056	0	1038	31	0
2	E	10	0	14	0	0
2	F	10	0	14	2	0
3	A	82	0	0	3	0
3	B	69	0	0	4	0
3	C	53	0	0	4	0
3	D	64	0	0	4	0
3	E	52	0	0	1	0
3	F	42	0	0	4	0
All	All	6763	0	6297	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 100 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:226:LYS:HD3	1:F:47:ASP:OD2	1.69	0.92
1:C:97:LYS:HD2	1:C:200:GLN:HG2	1.61	0.81
1:D:86:HIS:CE1	1:D:107:ILE:HD12	2.17	0.79
1:A:226:LYS:CD	1:F:47:ASP:OD2	2.31	0.77
1:B:95:SER:OG	3:B:301:HOH:O	1.65	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/160 (80%)	124 (97%)	4 (3%)	0	100	100
1	B	130/160 (81%)	127 (98%)	3 (2%)	0	100	100
1	C	129/160 (81%)	126 (98%)	3 (2%)	0	100	100
1	D	134/160 (84%)	127 (95%)	7 (5%)	0	100	100
1	E	128/160 (80%)	125 (98%)	3 (2%)	0	100	100
1	F	128/160 (80%)	123 (96%)	4 (3%)	1 (1%)	16	11
All	All	777/960 (81%)	752 (97%)	24 (3%)	1 (0%)	48	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	95	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/136 (83%)	113 (100%)	0	100	100
1	B	115/136 (85%)	107 (93%)	8 (7%)	14	10
1	C	113/136 (83%)	109 (96%)	4 (4%)	32	32
1	D	116/136 (85%)	113 (97%)	3 (3%)	40	44
1	E	114/136 (84%)	110 (96%)	4 (4%)	32	32
1	F	114/136 (84%)	105 (92%)	9 (8%)	11	8
All	All	685/816 (84%)	657 (96%)	28 (4%)	28	26

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	198	VAL
1	F	231	LEU
1	E	236[B]	LYS
1	F	214	SER
1	E	236[A]	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	76	GLN
1	F	70	GLN
1	F	219	GLN
1	F	192	GLN
1	E	200	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PGE	F	301	-	9,9,9	0.24	0	8,8,8	0.15	0
2	PGE	E	301	-	9,9,9	0.29	0	8,8,8	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PGE	F	301	-	-	4/7/7/7	-
2	PGE	E	301	-	-	1/7/7/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	301	PGE	O2-C3-C4-O3
2	F	301	PGE	O1-C1-C2-O2
2	F	301	PGE	O2-C3-C4-O3
2	F	301	PGE	O3-C5-C6-O4
2	F	301	PGE	C1-C2-O2-C3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	301	PGE	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	132/160 (82%)	-0.16	0 100 100	20, 32, 52, 60	0
1	B	132/160 (82%)	0.05	3 (2%) 61 60	17, 34, 62, 76	2 (1%)
1	C	133/160 (83%)	0.17	4 (3%) 52 51	22, 37, 65, 86	0
1	D	138/160 (86%)	0.15	1 (0%) 84 84	23, 37, 58, 71	0
1	E	131/160 (81%)	0.00	1 (0%) 82 82	14, 33, 56, 67	1 (0%)
1	F	131/160 (81%)	0.44	7 (5%) 32 31	27, 41, 74, 92	1 (0%)
All	All	797/960 (83%)	0.11	16 (2%) 65 64	14, 36, 63, 92	4 (0%)

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	68	ALA	3.4
1	F	227[A]	GLN	3.3
1	D	121	ALA	3.0
1	C	226	LYS	2.9
1	E	111	LEU	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PGE	E	301	10/10	0.77	0.19	69,79,87,88	0
2	PGE	F	301	10/10	0.86	0.14	69,71,74,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.