



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 03:22 PM UTC

PDB ID : 9EZJ / pdb_00009ezj
Title : Apo human TDO in complex with a bound inhibitor (Cpd-4)
Authors : Wicki, M.; Mac Sweeney, A.
Deposited on : 2024-04-12
Resolution : 2.61 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4951 (2.64-2.60)
Clashscore	190562	5303 (2.64-2.60)
Ramachandran outliers	187476	5217 (2.64-2.60)
Sidechain outliers	187428	5217 (2.64-2.60)
RSRZ outliers	180081	4950 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	403	14% 67% 10% 22%
1	B	403	14% 69% 9% 21%
1	C	403	21% 69% 8% 22%
1	D	403	14% 67% 9% 23%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10283 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan 2,3-dioxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	1	0
			2523	1640	428	445	10			
1	B	319	Total	C	N	O	S	0	1	0
			2539	1648	429	451	11			
1	C	315	Total	C	N	O	S	0	1	0
			2506	1627	424	445	10			
1	D	312	Total	C	N	O	S	0	0	0
			2487	1616	423	439	9			

There are 60 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	4	MET	-	initiating methionine	UNP P48775
A	5	ARG	-	expression tag	UNP P48775
A	6	GLY	-	expression tag	UNP P48775
A	7	SER	-	expression tag	UNP P48775
A	8	HIS	-	expression tag	UNP P48775
A	9	HIS	-	expression tag	UNP P48775
A	10	HIS	-	expression tag	UNP P48775
A	11	HIS	-	expression tag	UNP P48775
A	12	HIS	-	expression tag	UNP P48775
A	13	HIS	-	expression tag	UNP P48775
A	14	GLY	-	expression tag	UNP P48775
A	15	ILE	-	expression tag	UNP P48775
A	16	ASP	-	expression tag	UNP P48775
A	17	HIS	-	expression tag	UNP P48775
A	18	MET	-	expression tag	UNP P48775
B	4	MET	-	initiating methionine	UNP P48775
B	5	ARG	-	expression tag	UNP P48775
B	6	GLY	-	expression tag	UNP P48775
B	7	SER	-	expression tag	UNP P48775
B	8	HIS	-	expression tag	UNP P48775
B	9	HIS	-	expression tag	UNP P48775

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	10	HIS	-	expression tag	UNP P48775
B	11	HIS	-	expression tag	UNP P48775
B	12	HIS	-	expression tag	UNP P48775
B	13	HIS	-	expression tag	UNP P48775
B	14	GLY	-	expression tag	UNP P48775
B	15	ILE	-	expression tag	UNP P48775
B	16	ASP	-	expression tag	UNP P48775
B	17	HIS	-	expression tag	UNP P48775
B	18	MET	-	expression tag	UNP P48775
C	4	MET	-	initiating methionine	UNP P48775
C	5	ARG	-	expression tag	UNP P48775
C	6	GLY	-	expression tag	UNP P48775
C	7	SER	-	expression tag	UNP P48775
C	8	HIS	-	expression tag	UNP P48775
C	9	HIS	-	expression tag	UNP P48775
C	10	HIS	-	expression tag	UNP P48775
C	11	HIS	-	expression tag	UNP P48775
C	12	HIS	-	expression tag	UNP P48775
C	13	HIS	-	expression tag	UNP P48775
C	14	GLY	-	expression tag	UNP P48775
C	15	ILE	-	expression tag	UNP P48775
C	16	ASP	-	expression tag	UNP P48775
C	17	HIS	-	expression tag	UNP P48775
C	18	MET	-	expression tag	UNP P48775
D	4	MET	-	initiating methionine	UNP P48775
D	5	ARG	-	expression tag	UNP P48775
D	6	GLY	-	expression tag	UNP P48775
D	7	SER	-	expression tag	UNP P48775
D	8	HIS	-	expression tag	UNP P48775
D	9	HIS	-	expression tag	UNP P48775
D	10	HIS	-	expression tag	UNP P48775
D	11	HIS	-	expression tag	UNP P48775
D	12	HIS	-	expression tag	UNP P48775
D	13	HIS	-	expression tag	UNP P48775
D	14	GLY	-	expression tag	UNP P48775
D	15	ILE	-	expression tag	UNP P48775
D	16	ASP	-	expression tag	UNP P48775
D	17	HIS	-	expression tag	UNP P48775
D	18	MET	-	expression tag	UNP P48775

- Molecule 2 is alpha-methyl-L-tryptophan (CCD ID: ZIQ) (formula: C₁₂H₁₄N₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			16	12	2	2		
2	B	1	Total	C	N	O	0	0
			16	12	2	2		
2	C	1	Total	C	N	O	0	0
			16	12	2	2		
2	D	1	Total	C	N	O	0	0
			16	12	2	2		

- Molecule 3 is ethyl (9 {R})-2-methoxy-4-oxidanilidene-9-[[1 {S}]-1-phenylethyl]-[(2-prop an-2-ylphenyl)carbamoyl]amino]-6,7,8,9-tetrahydropyrido[1,2-a]pyrimidine-3-carboxylate (CCD ID: A1H8G) (formula: C₃₀H₃₆N₄O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			39	30	4	5		
3	B	1	Total	C	N	O	0	0
			39	30	4	5		
3	C	1	Total	C	N	O	0	0
			39	30	4	5		
3	D	1	Total	C	N	O	0	0
			39	30	4	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	O	0	0
			3	3		
4	B	2	Total	O	0	0
			2	2		
4	C	2	Total	O	0	0
			2	2		
4	D	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tryptophan 2,3-dioxygenase



- Molecule 1: Tryptophan 2,3-dioxygenase



- Molecule 1: Tryptophan 2,3-dioxygenase

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	133.79Å 156.16Å 90.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.33 – 2.61 45.33 – 2.61	Depositor EDS
% Data completeness (in resolution range)	74.7 (45.33-2.61) 74.6 (45.33-2.61)	Depositor EDS
R_{merge}	0.27	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 2.61Å)	Xtriage
Refinement program	BUSTER 2.11.8	Depositor
R, R_{free}	0.298 , 0.330 0.292 , 0.294	Depositor DCC
R_{free} test set	2199 reflections (3.77%)	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	10283	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.77 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.6072e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1H8G, ZIQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/2581	1.07	3/3494 (0.1%)
1	B	0.68	1/2597 (0.0%)	1.08	2/3518 (0.1%)
1	C	0.67	0/2561	1.07	1/3467 (0.0%)
1	D	0.66	0/2542	1.08	1/3441 (0.0%)
All	All	0.67	1/10281 (0.0%)	1.08	7/13920 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	148	SER	CA-C	5.11	1.59	1.52

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	143	PHE	CA-CB-CG	5.56	119.36	113.80
1	B	143	PHE	CA-CB-CG	5.39	119.19	113.80
1	A	99	ASN	CA-CB-CG	5.32	117.92	112.60
1	C	143	PHE	CA-CB-CG	5.21	119.01	113.80
1	B	99	ASN	CA-CB-CG	5.20	117.80	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2523	0	2393	25	0
1	B	2539	0	2379	21	0
1	C	2506	0	2351	21	0
1	D	2487	0	2345	23	0
2	A	16	0	0	0	0
2	B	16	0	0	0	0
2	C	16	0	0	0	0
2	D	16	0	0	0	0
3	A	39	0	0	0	0
3	B	39	0	0	0	0
3	C	39	0	0	0	0
3	D	39	0	0	0	0
4	A	3	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
4	D	1	0	0	0	0
All	All	10283	0	9468	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 80 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:334:ARG:HE	1:A:335[A]:MET:HE3	1.22	1.02
1:B:334:ARG:HE	1:B:335[A]:MET:HE3	1.23	1.01
1:C:334:ARG:HE	1:C:335[A]:MET:HE3	1.25	0.99
1:D:334:ARG:HE	1:D:335:MET:HE3	1.52	0.75
1:C:334:ARG:NE	1:C:335[A]:MET:HE3	2.04	0.70

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/403 (76%)	298 (97%)	6 (2%)	3 (1%)	12	26
1	B	314/403 (78%)	305 (97%)	7 (2%)	2 (1%)	21	39
1	C	308/403 (76%)	301 (98%)	7 (2%)	0	100	100
1	D	304/403 (75%)	296 (97%)	8 (3%)	0	100	100
All	All	1233/1612 (76%)	1200 (97%)	28 (2%)	5 (0%)	30	49

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	148	SER
1	B	148	SER
1	A	152	GLY
1	B	152	GLY
1	A	337	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/368 (69%)	242 (96%)	11 (4%)	26	49
1	B	250/368 (68%)	238 (95%)	12 (5%)	23	45
1	C	247/368 (67%)	237 (96%)	10 (4%)	28	53
1	D	246/368 (67%)	235 (96%)	11 (4%)	24	48
All	All	996/1472 (68%)	952 (96%)	44 (4%)	25	48

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	260	GLN
1	D	201	LEU
1	C	303	ARG
1	C	367	ASN
1	D	260	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	253	GLN
1	C	386	HIS
1	D	386	HIS
1	C	309	GLN
1	D	64	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	A1H8G	A	502	-	41,42,42	0.53	1 (2%)	49,59,59	0.47	0
3	A1H8G	B	502	-	41,42,42	0.51	1 (2%)	49,59,59	0.46	0
3	A1H8G	C	502	-	41,42,42	0.48	1 (2%)	49,59,59	0.46	0
2	ZIQ	A	501	-	14,17,17	1.40	1 (7%)	20,25,25	1.50	4 (20%)
2	ZIQ	B	501	-	14,17,17	1.46	1 (7%)	20,25,25	1.46	3 (15%)
3	A1H8G	D	502	-	41,42,42	0.46	1 (2%)	49,59,59	0.45	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ZIQ	C	501	-	14,17,17	1.37	2 (14%)	20,25,25	1.31	4 (20%)
2	ZIQ	D	501	-	14,17,17	1.48	2 (14%)	20,25,25	1.42	4 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1H8G	A	502	-	-	3/30/43/43	0/4/4/4
3	A1H8G	B	502	-	-	3/30/43/43	0/4/4/4
3	A1H8G	C	502	-	-	3/30/43/43	0/4/4/4
2	ZIQ	A	501	-	-	4/11/11/11	0/2/2/2
2	ZIQ	B	501	-	-	4/11/11/11	0/2/2/2
3	A1H8G	D	502	-	-	3/30/43/43	0/4/4/4
2	ZIQ	C	501	-	-	4/11/11/11	0/2/2/2
2	ZIQ	D	501	-	-	4/11/11/11	0/2/2/2

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	ZIQ	CD2-CG	-3.19	1.38	1.44
2	D	501	ZIQ	CD2-CG	-3.18	1.38	1.44
2	A	501	ZIQ	CD2-CG	-3.00	1.39	1.44
2	C	501	ZIQ	CD2-CG	-2.84	1.39	1.44
3	A	502	A1H8G	C27-N	2.73	1.42	1.38

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	ZIQ	CZ2-CE2-CD2	-3.74	118.57	122.19
2	B	501	ZIQ	CE3-CD2-CE2	3.28	122.24	118.84
2	B	501	ZIQ	CZ2-CE2-CD2	-3.23	119.06	122.19
2	D	501	ZIQ	CZ2-CE2-CD2	-3.16	119.13	122.19
2	D	501	ZIQ	CE3-CD2-CE2	3.14	122.10	118.84

There are no chirality outliers.

5 of 28 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	ZIQ	OXT-C-CA-C1
2	B	501	ZIQ	OXT-C-CA-C1
2	C	501	ZIQ	OXT-C-CA-C1
2	D	501	ZIQ	OXT-C-CA-C1
2	A	501	ZIQ	O-C-CA-C1

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1H8G A 502




Bond lengths



Bond angles




Torsions




Rings


Ligand A1H8G B 502




Bond lengths



Bond angles




Torsions




Rings


Ligand A1H8G C 502




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/403 (77%)	1.00	55 (17%) 4 3	18, 33, 70, 88	1 (0%)
1	B	319/403 (79%)	0.91	57 (17%) 3 3	17, 34, 97, 115	1 (0%)
1	C	315/403 (78%)	1.42	84 (26%) 1 1	18, 39, 67, 81	1 (0%)
1	D	312/403 (77%)	1.00	57 (18%) 3 3	16, 38, 62, 77	0
All	All	1260/1612 (78%)	1.08	253 (20%) 3 2	16, 36, 70, 115	3 (0%)

The worst 5 of 253 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	359	TYR	6.6
1	C	369	SER	5.5
1	B	246	GLU	5.4
1	B	240	ARG	5.4
1	C	357	ASP	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZIQ	A	501	16/16	0.86	0.13	30,31,34,34	0
3	A1H8G	C	502	39/39	0.89	0.13	36,38,41,42	0
3	A1H8G	B	502	39/39	0.90	0.12	33,34,37,37	0
3	A1H8G	D	502	39/39	0.90	0.12	34,36,40,41	0
2	ZIQ	C	501	16/16	0.91	0.11	30,31,32,32	0
2	ZIQ	D	501	16/16	0.92	0.09	27,27,29,29	0
3	A1H8G	A	502	39/39	0.93	0.10	27,27,29,30	0
2	ZIQ	B	501	16/16	0.95	0.07	26,27,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A1H8G C 502:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1H8G B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1H8G D 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A1H8G A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.