



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 09:48 PM UTC

PDB ID : 5F0Q / pdb_00005f0q
Title : Crystal structure of C-terminal domain of the human DNA primase large subunit with bound DNA template/RNA primer
Authors : Tahirov, T.H.; Baranovskiy, A.G.; Babayeva, N.D.
Deposited on : 2015-11-28
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)
RNA backbone	3983	1098 (2.54-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	191	6% 66% 27% • •
1	B	191	5% 60% 36% • •
2	C	6	50% 50%
2	E	6	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	12	17% 33% 17% 33% 17%
3	F	12	17% 42% 17% 8% 17%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3838 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA primase large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	187	Total	C	N	O	S	0	0	0
			1526	974	270	272	10			
1	B	185	Total	C	N	O	S	0	0	0
			1508	964	266	268	10			

- Molecule 2 is a RNA chain called RNA (5'-R(P*GP*GP*CP*GP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	P	0	0	0
			141	58	26	49	8			
2	E	6	Total	C	N	O	P	0	0	0
			141	58	26	49	8			

- Molecule 3 is a DNA chain called DNA (5'-D(*GP*CP*CP*GP*CP*CP*AP*AP*CP*AP*TP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	12	Total	C	N	O	P	0	0	0
			240	115	47	67	11			
3	F	10	Total	C	N	O	P	0	0	0
			199	95	40	55	9			

- Molecule 4 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Fe	S	0	0
			8	4	4		
4	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	Mg	0	0
			1	1		
5	E	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	24	Total	O	0	0
			24	24		
6	B	19	Total	O	0	0
			19	19		
6	C	7	Total	O	0	0
			7	7		
6	D	8	Total	O	0	0
			8	8		
6	E	4	Total	O	0	0
			4	4		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	F	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA primase large subunit



- Molecule 1: DNA primase large subunit



- Molecule 2: RNA (5'-R(P*GP*GP*CP*GP*GP*C)-3')





- Molecule 2: RNA (5'-R(P*GP*GP*CP*GP*GP*C)-3')



There are no outlier residues recorded for this chain.

- Molecule 3: DNA (5'-D(*GP*CP*CP*GP*CP*CP*AP*AP*CP*AP*TP*A)-3')



● Molecule 3: DNA (5'-D(*GP*CP*CP*GP*CP*CP*AP*AP*CP*AP*TP*A)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	122.51Å 126.00Å 83.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.48 – 2.21 49.48 – 2.21	Depositor EDS
% Data completeness (in resolution range)	93.3 (49.48-2.21) 93.7 (49.48-2.21)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.223 , 0.258 0.216 , 0.246	Depositor DCC
R_{free} test set	1552 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	51.8	Xtriage
Anisotropy	0.357	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.023 for -k,-h,-l 0.013 for -1/2*h-1/2*k+l,-1/2*h-1/2*k-l,1/2 *h-1/2*k 0.017 for -1/2*h-1/2*k-l,-1/2*h-1/2*k+l,-1/ 2*h+1/2*k 0.015 for -1/2*h+1/2*k-l,1/2*h-1/2*k-l,-1/2 *h-1/2*k 0.002 for -1/2*h+1/2*k+l,1/2*h-1/2*k+l,1/ 2*h+1/2*k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3838	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SF4, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/1567	0.98	6/2103 (0.3%)
1	B	0.43	0/1549	0.96	4/2081 (0.2%)
2	C	0.36	0/121	0.52	0/187
2	E	0.36	0/121	0.54	0/187
3	D	0.35	0/269	1.12	6/412 (1.5%)
3	F	0.30	0/223	0.96	2/341 (0.6%)
All	All	0.42	0/3850	0.96	18/5311 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	D	0	2
3	F	0	2
All	All	0	4

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	345	TYR	N-CA-C	9.01	122.23	111.33
3	D	3	DC	C2'-C3'-O3'	-7.91	99.63	111.50
1	B	345	TYR	N-CA-C	7.80	120.47	111.11
3	D	11	DT	N1-C1'-C2'	7.22	124.33	113.50
3	F	3	DC	C2'-C3'-O3'	-6.74	101.39	111.50
1	A	273	LEU	N-CA-C	6.42	117.94	111.07
1	A	334	GLY	N-CA-C	-5.87	106.40	115.73
3	F	2	DC	C2'-C3'-O3'	-5.78	102.83	111.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	306	ARG	N-CA-C	-5.76	105.09	111.71
1	A	446	GLN	N-CA-C	-5.73	105.12	111.36
1	A	425	GLN	N-CA-C	-5.61	105.17	111.28
3	D	8	DA	C4'-C3'-O3'	5.55	118.32	110.00
3	D	1	DG	C2'-C3'-O3'	-5.53	103.21	111.50
1	B	303	HIS	N-CA-C	5.52	117.73	111.11
1	A	276	ILE	N-CA-C	5.47	115.67	110.42
1	B	276	ILE	N-CA-C	5.37	115.57	110.42
3	D	2	DC	C2'-C3'-O3'	-5.37	103.45	111.50
3	D	7	DA	N9-C1'-C2'	5.09	121.14	113.50

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	D	3	DC	Sidechain
3	D	8	DA	Sidechain
3	F	3	DC	Sidechain
3	F	8	DA	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1526	0	1500	46	0
1	B	1508	0	1481	46	0
2	C	141	0	67	2	0
2	E	141	0	67	0	0
3	D	240	0	135	13	0
3	F	199	0	112	5	0
4	A	8	0	0	0	0
4	B	8	0	0	0	0
5	C	1	0	0	0	0
5	E	1	0	0	0	0
6	A	24	0	0	1	0
6	B	19	0	0	1	0
6	C	7	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	D	8	0	0	0	0
6	E	4	0	0	0	0
6	F	3	0	0	0	0
All	All	3838	0	3362	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:407:GLY:HA3	1:A:430:MET:HE1	1.39	1.01
1:B:337:ASP:HB2	1:B:340:LYS:HB2	1.57	0.86
1:A:278:LEU:HD11	1:A:282:LYS:HE3	1.64	0.78
3:D:10:DA:H2''	3:D:11:DT:H5'	1.65	0.76
3:D:11:DT:H4'	3:D:12:DA:OP1	1.84	0.76
3:F:3:DC:H2''	3:F:4:DG:H5'	1.67	0.75
1:A:334:GLY:O	1:A:336:MET:HE2	1.88	0.73
1:A:422:VAL:HA	1:A:425:GLN:HE21	1.54	0.72
1:A:407:GLY:HA3	1:A:430:MET:CE	2.18	0.72
1:A:302:ARG:HH11	1:A:302:ARG:HG3	1.55	0.69
1:A:407:GLY:CA	1:A:430:MET:HE1	2.22	0.68
1:A:358:LYS:HD2	1:A:360:THR:OG1	1.95	0.67
1:B:393:LEU:HG	1:B:397:LYS:HE3	1.77	0.67
1:A:307:MET:HA	1:A:307:MET:HE3	1.76	0.66
3:D:10:DA:C2'	3:D:11:DT:H5'	2.27	0.65
1:B:314:LYS:HG3	1:B:353:PHE:CE2	2.32	0.64
3:F:7:DA:H2''	3:F:8:DA:H5'	1.78	0.64
1:A:280:SER:HA	1:A:284:PHE:CG	2.32	0.64
1:B:412:LEU:O	1:B:416:LYS:HG3	1.97	0.64
1:A:337:ASP:HB2	1:A:340:LYS:HB2	1.77	0.64
3:D:10:DA:H2''	3:D:11:DT:H3'	1.79	0.64
3:D:7:DA:H2''	3:D:8:DA:H5'	1.81	0.63
3:D:11:DT:O4'	3:D:12:DA:H5'	2.03	0.58
1:B:364:PRO:HB2	1:B:443:HIS:HE1	1.68	0.57
1:A:396:GLN:NE2	1:B:278:LEU:HD11	2.19	0.57
1:B:395:LYS:O	1:B:399:GLN:HG3	2.05	0.57
1:B:290:GLN:HE22	1:B:397:LYS:HZ2	1.53	0.56
1:B:320:LEU:HD11	1:B:350:ARG:HG3	1.88	0.55
1:B:319:THR:OG1	1:B:322:GLN:HG3	2.07	0.55
3:D:10:DA:H1'	3:D:11:DT:H5'	1.88	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:426:LYS:O	1:A:430:MET:HG3	2.08	0.54
1:A:295:LEU:HG	1:A:330:GLU:HG2	1.89	0.54
1:B:437:CYS:HB2	1:B:439:PHE:CE2	2.42	0.54
1:A:286:PRO:HG3	1:A:386:PHE:CE2	2.43	0.54
1:B:280:SER:HA	1:B:284:PHE:CG	2.43	0.53
1:B:331:PHE:HB3	6:B:619:HOH:O	2.09	0.53
3:D:10:DA:H2''	3:D:11:DT:C3'	2.39	0.53
1:A:403:ILE:HG23	1:A:430:MET:HE3	1.91	0.53
1:B:421:GLN:HE21	1:B:442:ASN:HA	1.74	0.53
1:A:443:HIS:O	1:A:446:GLN:HB3	2.09	0.53
1:B:357:GLY:O	1:B:359:ARG:HG3	2.08	0.53
1:A:403:ILE:CG2	1:A:430:MET:HE3	2.39	0.52
1:A:411:ILE:HG23	1:A:423:ALA:HB1	1.92	0.52
1:A:273:LEU:C	1:A:275:GLN:H	2.18	0.52
1:B:425:GLN:O	1:B:429:GLU:HG3	2.10	0.51
1:B:443:HIS:O	1:B:446:GLN:HB3	2.11	0.50
1:B:293:LYS:O	1:B:297:GLU:HG3	2.11	0.50
1:A:280:SER:HA	1:A:284:PHE:CD2	2.46	0.49
1:B:337:ASP:OD1	1:B:337:ASP:N	2.37	0.49
1:B:411:ILE:HG23	1:B:423:ALA:HB1	1.93	0.49
1:B:441:LEU:HA	1:B:446:GLN:OE1	2.13	0.49
1:A:410:GLN:OE1	1:A:426:LYS:HE2	2.13	0.48
1:A:408:ILE:O	1:A:412:LEU:HG	2.13	0.48
1:B:290:GLN:NE2	1:B:397:LYS:HZ2	2.10	0.48
1:A:286:PRO:HG3	1:A:386:PHE:HE2	1.78	0.48
1:B:313:LEU:HB3	1:B:318:LEU:HD12	1.95	0.48
1:B:333:LYS:C	1:B:335:LYS:H	2.21	0.48
1:A:337:ASP:C	1:A:339:ASP:H	2.20	0.48
1:B:273:LEU:HD11	1:B:326:PHE:HB2	1.95	0.48
1:A:298:ASN:O	1:A:299:HIS:HB2	2.14	0.48
1:A:278:LEU:HD21	1:A:282:LYS:NZ	2.29	0.47
1:A:314:LYS:HG3	1:A:353:PHE:CE2	2.49	0.47
1:A:337:ASP:C	1:A:339:ASP:N	2.70	0.47
1:B:429:GLU:OE2	1:B:436:ASP:HA	2.15	0.47
3:D:1:DG:H2'	3:D:2:DC:C6	2.49	0.47
1:A:302:ARG:HG3	1:A:302:ARG:NH1	2.27	0.47
1:B:337:ASP:CB	1:B:340:LYS:HB2	2.39	0.46
3:F:8:DA:H2''	3:F:9:DC:C6	2.50	0.46
1:A:388:HIS:O	1:B:402:LYS:HE2	2.15	0.46
1:B:272:SER:HB3	1:B:275:GLN:OE1	2.16	0.46
1:A:435:ASP:OD1	1:A:436:ASP:HB2	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:393:LEU:O	1:B:397:LYS:HG3	2.15	0.46
1:A:429:GLU:OE2	1:A:436:ASP:HA	2.16	0.45
1:B:287:CYS:SG	1:B:288:MET:N	2.90	0.45
1:B:286:PRO:HD3	1:B:427:TYR:CE1	2.51	0.45
1:A:427:TYR:O	1:A:431:ILE:HG23	2.16	0.45
6:A:610:HOH:O	3:D:11:DT:H73	2.16	0.45
1:B:385:PRO:O	1:B:389:SER:HB2	2.16	0.45
1:B:321:GLU:H	1:B:321:GLU:CD	2.23	0.45
1:A:281:THR:HG23	1:A:289:ARG:NH2	2.32	0.45
1:B:358:LYS:HB2	3:F:6:DC:OP1	2.16	0.45
1:A:320:LEU:HD21	1:A:350:ARG:CG	2.48	0.44
1:B:439:PHE:CE2	1:B:450:GLU:HG3	2.52	0.44
1:B:454:ILE:HG22	1:B:455:LEU:HD23	1.99	0.44
1:A:373:SER:O	1:A:374:ASN:C	2.60	0.44
1:A:287:CYS:SG	1:A:288:MET:N	2.91	0.44
1:B:339:ASP:C	1:B:341:PHE:N	2.75	0.44
1:B:364:PRO:HB2	1:B:443:HIS:CE1	2.51	0.44
1:B:290:GLN:NE2	1:B:397:LYS:NZ	2.66	0.43
1:A:337:ASP:HB3	1:A:340:LYS:H	1.82	0.43
2:C:2:G:H2'	2:C:3:C:C6	2.53	0.43
1:A:286:PRO:CG	1:A:386:PHE:HE2	2.32	0.43
3:D:10:DA:C1'	3:D:11:DT:H5'	2.48	0.43
1:A:347:TYR:CD2	1:A:347:TYR:C	2.96	0.43
1:A:407:GLY:O	1:A:411:ILE:HG13	2.19	0.42
2:C:1:GTP:O2A	2:C:1:GTP:O1B	2.36	0.42
1:A:321:GLU:H	1:A:321:GLU:CD	2.27	0.42
1:B:410:GLN:NE2	1:B:426:LYS:NZ	2.67	0.42
1:B:410:GLN:NE2	1:B:426:LYS:HE3	2.34	0.42
3:F:1:DG:H2'	3:F:2:DC:C6	2.55	0.42
1:A:343:LYS:HG3	3:D:3:DC:OP2	2.19	0.42
1:A:320:LEU:HD21	1:A:350:ARG:HG3	2.02	0.41
1:A:374:ASN:N	1:A:375:PRO:HD3	2.35	0.41
3:D:11:DT:O4'	3:D:12:DA:C5'	2.67	0.41
1:B:280:SER:HA	1:B:284:PHE:CD2	2.56	0.41
1:B:452:GLN:HA	1:B:452:GLN:OE1	2.21	0.41
1:B:288:MET:HE1	1:B:308:GLN:HG2	2.02	0.40
1:B:393:LEU:CG	1:B:397:LYS:HE3	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/191 (97%)	174 (94%)	10 (5%)	1 (0%)	24	26
1	B	183/191 (96%)	176 (96%)	7 (4%)	0	100	100
All	All	368/382 (96%)	350 (95%)	17 (5%)	1 (0%)	36	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	391	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/171 (99%)	163 (96%)	6 (4%)	31	40
1	B	167/171 (98%)	162 (97%)	5 (3%)	36	47
All	All	336/342 (98%)	325 (97%)	11 (3%)	33	44

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	307	MET
1	A	329	GLN
1	A	337	ASP
1	A	405	PRO
1	A	430	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	435	ASP
1	B	337	ASP
1	B	369	LYS
1	B	391	PRO
1	B	404	SER
1	B	440	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	298	ASN
1	A	329	GLN
1	A	378	GLN
1	A	396	GLN
1	A	399	GLN
1	A	425	GLN
1	A	456	ASN
1	B	290	GLN
1	B	298	ASN
1	B	303	HIS
1	B	325	GLN
1	B	374	ASN
1	B	378	GLN
1	B	396	GLN
1	B	410	GLN
1	B	421	GLN
1	B	445	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	4/6 (66%)	0	0
2	E	4/6 (66%)	0	0
All	All	8/12 (66%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SF4	B	501	1	0,12,12	-	-	-		
4	SF4	A	501	1	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	B	501	1	-	-	0/6/5/5
4	SF4	A	501	1	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	187/191 (97%)	0.52	12 (6%) 25 22	39, 63, 97, 110	0
1	B	185/191 (96%)	0.59	10 (5%) 31 28	39, 65, 92, 104	0
2	C	5/6 (83%)	-1.13	0 100 100	44, 45, 49, 50	0
2	E	5/6 (83%)	-1.10	0 100 100	48, 50, 52, 55	0
3	D	12/12 (100%)	-0.14	2 (16%) 4 3	43, 52, 87, 97	0
3	F	10/12 (83%)	-0.47	0 100 100	44, 51, 62, 104	0
All	All	404/418 (96%)	0.47	24 (5%) 28 25	39, 63, 94, 110	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	336	MET	4.6
1	B	271	ILE	4.2
1	A	343	LYS	3.5
1	B	331	PHE	3.4
1	A	331	PHE	3.3
1	A	345	TYR	3.3
1	B	455	LEU	3.1
1	A	344	GLY	2.9
1	A	336	MET	2.9
1	B	332	ILE	2.8
1	A	435	ASP	2.7
1	A	341	PHE	2.6
1	B	335	LYS	2.5
3	D	11	DT	2.5
1	A	270	LYS	2.5
1	B	337	ASP	2.3
1	B	338	PRO	2.3
1	A	373	SER	2.2
1	A	332	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	D	12	DA	2.1
1	B	282	LYS	2.1
1	B	275	GLN	2.1
1	A	342	ASP	2.0
1	A	340	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MG	C	101	1/1	0.85	0.11	78,78,78,78	0
5	MG	E	101	1/1	0.88	0.10	88,88,88,88	0
4	SF4	A	501	8/8	0.99	0.03	47,49,54,54	0
4	SF4	B	501	8/8	0.99	0.03	44,47,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.