



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:29 AM UTC

PDB ID : 6F0A / pdb_00006f0a
Title : Crystal structure of human indoleamine 2,3-dioxygenase bound to a triazole inhibitor and alanine molecule.
Authors : Swan, M.K.; Latchem, M.
Deposited on : 2017-11-17
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	3% 87% 10% ..
1	C	393	4% 85% 11% ..

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6378 atoms, of which 14 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Indoleamine 2,3-dioxygenase 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	383	Total	C	N	O	S	0	0	0
			3016	1935	514	550	17			
1	C	381	Total	C	N	O	S	0	0	0
			3007	1931	514	545	17			

- Molecule 2 is {N}-(4-chlorophenyl)-1 {H}-1,2,3-triazol-5-amine (CCD ID: C82) (formula: C₈H₇ClN₄) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Cl	H	N	0
			20	8	1	7	4	
2	C	1	Total	C	Cl	H	N	0
			20	8	1	7	4	

- Molecule 3 is ALANINE (CCD ID: ALA) (formula: C₃H₇NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			6	3	1	2		
3	C	1	Total	C	N	O	0	0
			6	3	1	2		

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	107	Total	O	0	0
			107	107		
5	C	110	Total	O	0	0
			110	110		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Indoleamine 2,3-dioxygenase 1



• Molecule 1: Indoleamine 2,3-dioxygenase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.48Å 96.89Å 132.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.78 – 2.26 71.78 – 2.26	Depositor EDS
% Data completeness (in resolution range)	98.4 (71.78-2.26) 98.4 (71.78-2.26)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 2.27Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.191 , 0.225 0.199 , 0.233	Depositor DCC
R_{free} test set	2600 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	45.2	Xtriage
Anisotropy	0.692	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 24.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6378	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, C82

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	1/3085 (0.0%)	1.32	16/4173 (0.4%)
1	C	0.75	0/3076	1.34	11/4160 (0.3%)
All	All	0.77	1/6161 (0.0%)	1.33	27/8333 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	349	ILE	CG1-CD1	-5.77	1.29	1.51

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	265	GLY	N-CA-C	-6.98	105.71	114.16
1	A	107	ILE	CB-CA-C	6.75	117.52	110.91
1	A	401	LYS	CA-C-N	6.48	133.92	121.54
1	A	401	LYS	C-N-CA	6.48	133.92	121.54
1	A	214	PHE	CA-CB-CG	6.43	120.23	113.80
1	A	383	ASP	N-CA-C	-6.42	97.36	108.52
1	C	383	ASP	CA-C-N	6.21	133.40	121.54
1	C	383	ASP	C-N-CA	6.21	133.40	121.54
1	A	356	ILE	N-CA-CB	5.57	114.80	110.45
1	A	243	LEU	CA-C-N	5.48	127.88	120.38
1	A	243	LEU	C-N-CA	5.48	127.88	120.38
1	A	383	ASP	CA-C-N	5.44	127.83	120.38
1	A	383	ASP	C-N-CA	5.44	127.83	120.38
1	A	145	TYR	CA-C-N	5.43	128.09	120.28
1	A	145	TYR	C-N-CA	5.43	128.09	120.28
1	C	303	HIS	CA-C-N	5.39	127.50	120.28
1	C	303	HIS	C-N-CA	5.39	127.50	120.28
1	C	128	ASP	CA-CB-CG	5.31	117.91	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	TYR	CA-C-N	5.30	127.91	120.28
1	A	126	TYR	C-N-CA	5.30	127.91	120.28
1	C	158	ASP	CA-CB-CG	5.12	117.72	112.60
1	C	188	MET	CA-C-N	5.12	127.10	120.44
1	C	188	MET	C-N-CA	5.12	127.10	120.44
1	A	125	VAL	CA-C-N	5.03	127.33	120.54
1	A	125	VAL	C-N-CA	5.03	127.33	120.54
1	C	125	VAL	CA-C-N	5.00	127.29	120.54
1	C	125	VAL	C-N-CA	5.00	127.29	120.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3016	0	3010	17	0
1	C	3007	0	3014	17	0
2	A	13	7	0	0	0
2	C	13	7	0	1	0
3	A	6	0	4	0	0
3	C	6	0	4	1	0
4	A	43	0	30	1	0
4	C	43	0	30	3	0
5	A	107	0	0	0	0
5	C	110	0	0	0	0
All	All	6364	14	6092	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:CYS:HG	1:C:308:CYS:HG	0.94	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:178:ILE:HD11	1:C:269:VAL:HG23	1.58	0.84
1:C:175:ALA:HA	1:C:178:ILE:HD12	1.64	0.80
1:A:178:ILE:HD11	1:A:269:VAL:HG23	1.69	0.73
1:C:264:ALA:HB3	4:C:503:HEM:C1D	2.30	0.66
1:A:264:ALA:HB3	4:A:503:HEM:C1D	2.36	0.61
1:A:11:ILE:HG13	1:A:12:SER:H	1.72	0.54
1:C:177:ALA:HB2	1:C:206:CYS:HB2	1.91	0.52
1:C:264:ALA:O	1:C:267:SER:HB3	2.09	0.52
1:A:345:TYR:CE2	1:A:349:ILE:HD11	2.45	0.51
1:A:24:ALA:HA	1:A:131:LEU:HB3	1.93	0.49
1:A:188:MET:HE3	1:A:319:PHE:HB2	1.96	0.48
1:A:12:SER:HB2	1:A:17:ILE:O	2.13	0.48
1:A:178:ILE:CD1	1:A:269:VAL:HG23	2.41	0.47
1:A:160:SER:HB2	1:A:164:PHE:CE2	2.50	0.47
1:A:175:ALA:HA	1:A:178:ILE:CD1	2.48	0.44
1:C:80:ARG:HG3	1:C:114:LEU:HD13	1.99	0.44
1:C:226:PHE:CE2	4:C:503:HEM:HMB3	2.52	0.44
1:C:160:SER:HB2	1:C:164:PHE:CE2	2.52	0.43
1:C:88:MET:HE2	1:C:124:LEU:HB3	2.00	0.43
1:C:296:ARG:O	1:C:304:ARG:HD3	2.19	0.43
1:A:345:TYR:CE2	1:A:349:ILE:CD1	3.02	0.43
1:A:226:PHE:O	1:A:231:ARG:HG3	2.18	0.42
1:C:347:LEU:O	1:C:351:THR:HG23	2.19	0.42
1:A:175:ALA:HA	1:A:178:ILE:HD12	2.01	0.42
1:C:126:TYR:HD1	2:C:501:C82:C	2.32	0.41
1:C:24:ALA:HA	1:C:131:LEU:HB3	2.00	0.41
1:C:354:ILE:HD13	1:C:384:LEU:HD11	2.02	0.41
3:C:502:ALA:HA	4:C:503:HEM:CHB	2.50	0.41
1:A:76:GLN:HB3	1:A:118:LEU:HD11	2.03	0.41
1:A:126:TYR:OH	1:A:171:GLU:HG2	2.21	0.41
1:C:281:GLN:HE22	1:C:391:VAL:HA	1.86	0.41
1:A:264:ALA:O	1:A:267:SER:HB3	2.21	0.40
1:C:17:ILE:HG21	1:C:178:ILE:HG21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/393 (96%)	371 (98%)	7 (2%)	1 (0%)	36	40
1	C	377/393 (96%)	367 (97%)	9 (2%)	1 (0%)	36	40
All	All	756/786 (96%)	738 (98%)	16 (2%)	2 (0%)	36	40

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	402	GLU
1	C	384	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/340 (96%)	314 (96%)	12 (4%)	30	37
1	C	326/340 (96%)	310 (95%)	16 (5%)	22	24
All	All	652/680 (96%)	624 (96%)	28 (4%)	26	31

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	GLU
1	A	101	LYS
1	A	107	ILE
1	A	114	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	178	ILE
1	A	189	GLN
1	A	209	LYS
1	A	212	GLN
1	A	254	GLU
1	A	280	GLN
1	A	337	LYS
1	A	342	LEU
1	C	61	LYS
1	C	62	LEU
1	C	99	VAL
1	C	107	ILE
1	C	114	LEU
1	C	212	GLN
1	C	243	LEU
1	C	254	GLU
1	C	272	CYS
1	C	308	CYS
1	C	318	GLU
1	C	337	LYS
1	C	351	THR
1	C	361	GLN
1	C	398	SER
1	C	401	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	GLN
1	A	191	GLN
1	A	216	GLN
1	A	220	HIS
1	A	313	ASN
1	A	360	GLN
1	C	106	ASN
1	C	281	GLN
1	C	290	GLN
1	C	303	HIS
1	C	313	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	C82	A	501	4	13,14,14	0.16	0	14,18,18	0.53	0
2	C82	C	501	4	13,14,14	0.15	0	14,18,18	0.37	0
4	HEM	A	503	2,1	50,50,50	1.40	5 (10%)	67,82,82	1.39	10 (14%)
4	HEM	C	503	2,1	50,50,50	1.47	8 (16%)	67,82,82	1.41	12 (17%)
3	ALA	C	502	-	5,5,5	1.17	0	6,6,6	1.08	0
3	ALA	A	502	-	5,5,5	1.10	0	6,6,6	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C82	A	501	4	-	0/4/4/4	0/2/2/2
2	C82	C	501	4	-	2/4/4/4	0/2/2/2
4	HEM	A	503	2,1	-	6/14/54/54	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEM	C	503	2,1	-	7/14/54/54	-
3	ALA	C	502	-	-	2/4/4/4	-
3	ALA	A	502	-	-	0/4/4/4	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	503	HEM	C1B-NB	-5.31	1.31	1.40
4	A	503	HEM	C1B-NB	-4.47	1.32	1.40
4	A	503	HEM	O2D-CGD	3.05	1.40	1.30
4	C	503	HEM	CBD-CGD	2.81	1.57	1.50
4	C	503	HEM	C1D-ND	-2.73	1.33	1.38
4	A	503	HEM	C3B-C4B	2.39	1.49	1.44
4	A	503	HEM	C1D-ND	-2.37	1.34	1.38
4	C	503	HEM	C4D-ND	-2.34	1.36	1.40
4	C	503	HEM	CHA-C1A	-2.30	1.34	1.39
4	C	503	HEM	C4D-C3D	2.28	1.48	1.45
4	A	503	HEM	CHD-C4C	-2.19	1.34	1.38
4	C	503	HEM	C3B-C4B	2.13	1.49	1.44
4	C	503	HEM	FE-ND	2.01	2.01	1.94

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	HEM	C4B-C3B-C2B	-4.41	103.23	107.28
4	C	503	HEM	C4B-C3B-C2B	-4.12	103.49	107.28
4	A	503	HEM	C3B-C4B-NB	4.05	112.38	109.47
4	C	503	HEM	C3B-C4B-NB	2.67	111.39	109.47
4	C	503	HEM	CHD-C1D-C2D	-2.60	120.93	125.03
4	A	503	HEM	CHD-C1D-ND	2.51	127.13	124.42
4	C	503	HEM	O2A-CGA-O1A	-2.51	116.88	123.33
4	C	503	HEM	C2B-C1B-NB	2.51	112.72	109.84
4	A	503	HEM	C1A-CHA-C4D	-2.45	120.49	126.25
4	C	503	HEM	C1A-CHA-C4D	-2.44	120.50	126.25
4	A	503	HEM	C1C-CHC-C4B	-2.44	120.83	126.02
4	A	503	HEM	CHA-C4D-ND	2.38	127.31	124.37
4	C	503	HEM	O2D-CGD-O1D	-2.34	117.30	123.33
4	C	503	HEM	O2D-CGD-CBD	2.32	121.33	114.00
4	A	503	HEM	C2B-C1B-NB	2.31	112.50	109.84
4	C	503	HEM	CHA-C4D-C3D	-2.26	121.06	125.23
4	C	503	HEM	C2D-C1D-ND	2.25	112.50	109.90
4	C	503	HEM	C4A-NA-C1A	-2.23	102.19	105.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	503	HEM	C2A-C1A-NA	2.13	112.52	110.15
4	A	503	HEM	CHD-C1D-C2D	-2.10	121.71	125.03
4	A	503	HEM	CHA-C4D-C3D	-2.08	121.38	125.23
4	A	503	HEM	CHC-C1C-NC	2.04	126.67	124.45

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	503	HEM	C2C-C3C-CAC-CBC
4	A	503	HEM	C4C-C3C-CAC-CBC
4	C	503	HEM	C2C-C3C-CAC-CBC
4	C	503	HEM	C2A-CAA-CBA-CGA
3	C	502	ALA	O-C-CA-CB
3	C	502	ALA	OXT-C-CA-CB
4	C	503	HEM	C4C-C3C-CAC-CBC
4	A	503	HEM	CAD-CBD-CGD-O1D
4	A	503	HEM	CAD-CBD-CGD-O2D
4	C	503	HEM	CAA-CBA-CGA-O2A
4	C	503	HEM	CAD-CBD-CGD-O1D
4	C	503	HEM	CAA-CBA-CGA-O1A
4	A	503	HEM	CAA-CBA-CGA-O2A
4	A	503	HEM	CAA-CBA-CGA-O1A
2	C	501	C82	C7-C6-N-C5
2	C	501	C82	N3-C6-N-C5
4	C	503	HEM	CAD-CBD-CGD-O2D


There are no ring outliers.

4 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	501	C82	1	0
4	A	503	HEM	1	0
4	C	503	HEM	3	0
3	C	502	ALA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be


highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand HEM A 503




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/393 (97%)	0.17	12 (3%) 51 51	35, 51, 76, 119	0
1	C	381/393 (96%)	0.29	14 (3%) 45 45	39, 54, 82, 108	0
All	All	764/786 (97%)	0.23	26 (3%) 48 48	35, 53, 81, 119	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	11	ILE	8.3
1	C	11	ILE	7.1
1	C	374	LEU	4.4
1	A	403	GLY	4.2
1	A	363	LYS	4.0
1	A	285	GLY	3.8
1	C	402	GLU	3.7
1	C	362	PRO	3.6
1	A	272	CYS	3.4
1	A	374	LEU	3.2
1	A	12	SER	3.1
1	C	67	ILE	3.1
1	A	362	PRO	2.8
1	A	282	THR	2.7
1	C	12	SER	2.6
1	C	69	HIS	2.6
1	A	96	HIS	2.6
1	C	61	LYS	2.5
1	C	41	PHE	2.5
1	A	174	ALA	2.4
1	A	43	ALA	2.2
1	C	273	PHE	2.1
1	C	118	LEU	2.1
1	C	355	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	107	ILE	2.0
1	C	13	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	C82	C	501	13/13	0.94	0.10	47,56,60,60	0
2	C82	A	501	13/13	0.95	0.09	48,50,56,57	0
3	ALA	C	502	6/6	0.95	0.12	44,47,49,65	0
3	ALA	A	502	6/6	0.98	0.06	38,43,45,49	0
4	HEM	A	503	43/43	0.98	0.07	38,40,45,46	0
4	HEM	C	503	43/43	0.98	0.07	39,40,44,45	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around C82 C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around C82 A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM C 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.